GRBS FOLLOWED-UP BY THE BOOTES NETWORK

S. Guziy^{1,17}, A. Castro-Tirado¹, M. Jelínek¹, J. Gorosabel^{1,21,22},
P. Kubánek¹, R. Cunniffe¹, O. Lara-Gil¹, O. Rabaza-Castillo¹,
A. de Ugarte Postigo¹, R. Sánchez-Ramírez¹, J. Tello¹,
C. Pérez del Pulgar³, S. Castillo-Carrión⁴, J. Castro Cerón⁵,
T. de J. Mateo Sanguino⁶, R. Hudec⁷, S. Vitek⁸,
B. de la Morena Carretero⁹, J. Díaz Andreu⁹, R. Fernández-Muñoz¹⁰,
D. Pérez-Ramírez¹¹, P. Yock¹², W. Allen¹³, I. Bond¹⁴, I. Kheifets¹⁷,
G. Christie¹⁵, L. Sabau-Graziati¹⁶, C. Cui¹⁸, Y. Fan¹⁹ and I.H. Park²⁰

¹ Instituto de Astrofísica de Andalucía, IAA-CSIC, PO Box 03004, 18080 Granada, Spain

² Image Processing Laboratory, Univ. de Valencia, Burjassot, Valencia, Spain

³ Departamento de Ingeniería de Sistemas y Automática, Univ. de Málaga, Spain

⁴ Servicio Central de Informática, Universidad de Málaga, Spain

⁵ European Space Astronomy Centre, ESAC, Villafranca del Castillo Madrid, Spain

⁶ Departamento de Ingeniería de Sistemas y Automática, Universidad de Huelva, E.P.S. de La Rábida, Huelva, Spain

⁷ Astronomical Institute, Academy of Sciences of the Czech Republic, 251 65 Ondřejov, Czech Republic

⁸ Czech Technical University, Faculty of Electronic Engineering, Dep. of Radioelectronics, Czech Republic

⁹ Estación de Sondeos Atmosféricos, ESAt de El Arenosillo, CEDEA-INTA, Mazagón, Huelva, Spain

¹⁰ Estación Experimental de La Mayora, EELM-CSIC, Algarrobo Costa, Málaga, Spain

¹¹ Universidad de Jaén, Campus Las Lagunillas, Jaén, Spain

¹² University of Auckland, New Zealand

¹³ Vintage Lane Observatory, Blenheim, New Zealand

¹⁴ Massey University, New Zealand

¹⁵ Stardome Observatory, Auckland, New Zealand

¹⁶ División de Ciencias del Espacio, INTA, Torrejón de Ardoz, Madrid, Spain

¹⁷ Astronomical Observatory, Nikolaev National University, Nikolaev, Ukraine

¹⁸ National Astronomical Observatory, Chinese Academy of Sciences, Beijing 100012, China

¹⁹ Yunnan Astronomical Observatory, CAS Kunming, 650011 Yunnan, China

²⁰ Ewha Womans University, Daehyundong 11-1, Seodaemungu, 120-750 Seoul, South Korea

²¹ Unidad Asociada Grupo Ciencia Planetarias UPV/EHU-IAA/CSIC, Departamento de Física Aplicada I, E.T.S. Ingeniería, Universidad del País Vasco UPV/EHU, Alameda de Urquijo s/n, 48013 Bilbao, Spain

²² Ikerbasque, Basque Foundation for Science, Alameda de Urquijo 36-5, 48008 Bilbao, Spain

Abstract. The Burst Observer and Optical Transient Exploring System (BOOTES), is a global robotic observatory network, which started in 1998 with Spanish leadership devoted to study optical emissions from gamma ray bursts (GRBs) that occur in the Universe. We present shot history and current status of BOOTES network. The Network philosophy, science and some details of 117 GRBs followed-up are discussed.

1 Introduction

The Burst Observer and Optical Transient Exploring System (BOOTES), started in 1998 as a Spanish-Czech collaboration devoted to study optical emissions from gamma ray bursts (GRBs).

The first BOOTES robotic astronomical station was located at INTA's Estación de Sondeos Atmosféricos in Centro de Experimentación de El Arenosillo, a dark-sky site near Mazagón (Huelva), center owned by the Instituto Nacional de Técnica Aeroespacial (INTA). The second observing station was opened in 2001 and it is located at the Estación Experimental de La Mayora (dubbed BOOTES-2), 240 km apart. The latter is run by the Consejo Superior de Investigaciones Científicas (CSIC). In 2009 BOOTES expanded abroad, with the third station (BOOTES-3) being installed in Blenheim (South Island, New Zealand). The fourth one (BOOTES-4) has been deployed in 2011 at the Lijiang Astronomical Observatory (unnan, China).

2 Location

Four BOOTES station are located in the three countries (Spain, New Zealand and China) that have different geographic coordinates. In our dreams and plans an installation of three new station in the other geographic locations (countries). A general view of BOOTES stations in the world is shown in Figure 1.

3 Science and goals

The observation of the GRB error box simultaneously to the GRB occurrence Although the first detected optical counterparts were not brighter than 19th mag few hours after the burst, there have been several GRBs for which optical transient emission has been detected simultaneously to the gamma-ray event, with magnitudes in the range 5–10. The faint transient emission that has been detected few hours after the event is a consequence of the expanding remnant that the GRB leaves behind it. This provides information about the surrounding medium, but not about the burster itself. The fast slewing 0.6 m BOOTES telescopes should produce important results in this field.

The detection of optical flashes (OTs) of cosmic origin that could be unrelated to GRBs and constitute a new type of different astrophysical phenomenon

Fig. 1. The telescopes locations in the world.

(perhaps associated to QSOs/AGNs). If some of them are related to GRBs, the most recent GRB models predict that there should be a large number of bursting sources in which only transient X-ray/optical emission should be observed, but no gamma-ray emission. The latter would be confined in a jet-like structure and pointing towards us only in a few cases.

The monitoring of high-energy targets in different optical, as ground-based support for the ESA's International Gamma-Ray Laboratory (INTEGRAL) in which Spain had, for the first time, the leadership in one of the instruments, the Optical Transient Camera (OMC). This included test of technologies, methods, data processing, ground-based observational network, etc.. *INTEGRAL* was launched in 2002.

The monitoring of several objects (bright AGNs/QSOs, old GRB positions, etc.) looking for recurrent optical transient optical emission arising from these sources. There are hints that sudden and rapid flares occurs, though of smaller amplitude. This will be achieved by means of the 0.6-m network of BOOTES telescopes.

4 GRBs with BOOTES network

GRBs are indeed one of the main scientific goals of BOOTES. We know that GRBs arise at cosmological distances (with mean redshift $z \sim 2.5$ and redshifts in the range ~ 0.01 to ~ 10), with huge isotropic equivalent radiated energy, and small timescales (in the range few ms to 10^2 s), thus implying a small emitting region. The spectrum is non-thermal and relativistic outflows ($\Gamma > 100$) are involved. A frequent assumption is that short and long GRBs (with the short ones representing 1/3 of the overall GRB population) are due to different progenitors leading to the same succession of events: formation of a compact object and ejection of

a relativistic outflow which produces the (long-lasting) afterglow at other wavelengths. Main program for BOOTES network system is observations optical counterparts for gamma ray burst: open and monitoring OT in different filters. The BOOTES network philosophy is: identical telescopes spaced around the Earth, identical filter sets: g'r'i'ZY, identical CCD cameras, impact on several scientific fields and public outreach, Castro-Tirado *et al.* (2012).

Some summary of GRB response at the BOOTES stations and examples of observations we are present below:

- **BOOTES-1/0.3 m.** Around 50 real-time follow-ups in 2004-2012, with 10 detections of the OA and 27 observations with upper limit, 12 publications (GCN circulars and other), Jelínek *et al.* (2010);
- **BOOTES-2/0.3 m.** 24 real-time follow-ups, with 9 upper limit of the OA and 5 publications;
- **BOOTES-2/0.6 m – TELMA.** 8 real-time follow-ups with 3 detections of the OA, 5 observations with upper limit, 7 publications;
- **BOOTES-3/0.6 m – YA.** 23 real-time follow-ups with 6 detections of the OA, 10 observations with upper limit, 18 publications;
- **BOOTES-4/0.6 m – MET.** 12 real-time follow-ups with 1 detection of the OA, 5 observations with upper limit, 2 publications (GCN circulars).

5 Discussion

We have shown the advances in establishing the worldwide network of BOOTES telescopes in different locations around the Earth. BOOTES has played a significant role in the gamma-ray burst field over the last decade. Multiwavelength observations (photometry, spectroscopy, polarimetry) are ideal to better understand the GRB diversity. As of Sep. 2012, the number of GRBs followed-up at the four BOOTES stations is 117, with 20 optical counterpart detections and 56 upper limits reported (the rest being too crowd fields or unusable due to dew, low airmass, unfocused images, ...), altogether leading to 44 publications.

Installing the remaining BOOTES stations will help in continuous monitoring for some celestial sources, building more precise light curve for the targets. BOOTES contributes significantly to the GLORIA Network (EU-FP7) too. More detailed information about the BOOTES network can be seen in <http://bootes.iaa.es>

We acknowledge the support of the Spanish Ministry Projects AYA 2009-14000-C03-01 and AYA 2012-39727-C03-01.

References

- Castro-Tirado, A.J., Jelínek, M., Gorosabel, J., *et al.*, 2012, BASI, in press
 Jelínek, M., Castro-Tirado, A.J., de Ugarte Postigo, A., *et al.*, 2010, AdAst, id. 432172