ORTHO AMINO CYANO AROMATIC COMPOUNDS AS PRECURSOR FOR THE SYNTHESIS OF SOME NOVEL HETEROCYCLIC COMPOUNDS

V.S.Rao* ^a, S.V.S. Arun Kumar Gupta ^a and B.S.Reddy ^b

a: Department of Chemistry, Birla Institute of Technology & Science, Pilani, Rajasthan, India.

B: Suven Pharmaceuticals, Hyderabad.

Abstract: Pyrimidine-4(1H)-ones from 2-amino-3-cyano aromatic compounds were prepared in the presence of formic acid under strong acid catalyst. The intermediates and the final products structures were confirmed by spectral and elemental data.

Introduction:

2-amino-3-cyano aromatic compounds are used as a building blocks for the synthesis of fused pyrimidine-4(1H)-ones, which are intermediates in the synthesis of antipyretics, diuretics, anticonvulsants, sedatives, coronary dilators and anti-cancer agents ¹⁻⁴. Historically quinazolines-4(1H)-ones have been prepared by Neimantowski reaction of anthranilic acids ⁵. Other literature methods were cyclisation of ortho esters ⁶, Gold's reagents ⁷, ethoxymethylenemalononitrile with 6-amino benzaxamides ⁸, amination of benzoxazines ⁹, synthesis of quinazoline derivatives via, palladium catalyzed reductive cyclisation methods ¹⁰. Literature reveals that ortho cyano amino compounds are often more readily accessible synthetic targets than the corresponding anthranilic acids

Synthesis and Results:

For the first time, we are reporting a new cyclisation method on 2-amino-3-cyano pyridines, pyrazolines and pyridones systems, which affords good yields of pyrido-(2,3-d)-pyrimidine-4(1H)-ones (II), Pyrazolo-(3,4-d)-pyrimidine-4(7H)-ones (IV) and pyrido-(2,3-d)-pyrimidine-4,7-(1H)-diones (VI) respectively. The advantage of this method is simpler experimental conditions easy isolation procedure, shorter reaction times and delivery of the desired product in good purity with quantitative yields ¹⁴.

Cyclisation of 2-cyano-3-amino-5,7-diaryl pyridines (**I**), 2-cyano-3-amino-5-aryl pyrazoles (**III**), 2,5-dicyano-3-amino-4-aryl-pyridine-6-ones (**V**) with formic acid in the presence of a trace amount of sulfuric acid at 110 °C for 25 min, gave compound **II**, **IV**, **VI** respectively. The structure of the compounds were established on the basis of IR, PMR and Mass spectral data and the result of elemental analysis (see table-I). The PMR spectra of compounds **II**, **IV** and **VI** are quite interesting in that the protons **N-H** and **vinyl protons** are observed in the range of 7.5 and 8.3 ppm respectively¹⁵.

The synthesis of the starting compounds **I**, **III** and **V** was carried out according to the literature procedure. The reaction is believed to proceed via the formation of an amide intermediate. This intermediate is isolated and synthesized on a different route. Compounds on further cyclisation under the same experimental conditions gave **II**, **IV** and **VI**, which was found to be identical with compounds obtained earlier.

SCHEME-II

TABLE-1: Physical , C,H,N analysis

Compd. No.	R	R'	Yield (%)	M.P. (C)	Mol. for (Mol. wt)	Analysis found (Calcd.) (%)		
						C	H	N
II a.	CH ₃	H	76	230-31	C ₈ H ₇ N ₃ O (161)	59.60 (59.62)	4.34 (4.29)	26.08 (26.10)
b.	C ₆ H ₅	H	78	199-202	C ₁₃ H ₉ N ₃ O (223)	69.89 (69.95)	4.03 (3.99)	18.81 (18.84)
c.	C ₆ H ₅	C ₆ H ₅	75	210-12	C ₁₉ H ₁₃ N ₃ O (301)	75.69 (75.74)	4.29 (4.31)	13.87 (13.95)
d.	4-ClC ₆ H ₄	C ₆ H ₅	78	216-18	C ₁₉ H ₁₂ N ₃ OCI (335.5)	67.89 (67.94)	3.50 (3.57)	12.48 (12.51)
e.	C ₆ H ₅	2-Cl-C ₆ H ₅	86	240-42	C ₁₉ H ₁₂ N ₃ OCI (335.5)	67.90 (67.94)	3.56 (3.57)	12.44 (12.51)
IV a.	H	C ₆ H ₅	77	216-18	C ₁₇ H ₁₂ N ₄ O (288)	70.80 (70.82)	4.15 (4.16)	19.40 (19.44)
b.	H	4-ClC ₆ H ₄	79	232-34	C ₁₇ H ₁₁ N ₄ O (322.5)	63.20 (63.24)	3.45 (3.41)	17.30 (17.36)
c.	H	3-NO ₂ C ₆ H ₄	71	240-42	C ₁₇ H ₁₁ N ₆ O ₃ (328)	62.15 (62.17)	3.30 (3.35)	19.55 (19.50)
d.	H	2-ClC ₆ H ₄	81	222-23	C ₁₇ H ₁₁ N ₄ OCI (330)	61.79 (61.81)	3.29 (3.33)	16.91 (16.96)
e.	H	4-CH ₃ C ₆ H ₄	80	198-99	C ₁₈ H ₁₄ N ₄ O (310)	69.65 (69.66)	4.45 (4.51)	18.10 (18.06)
VI a.	H	C ₆ H ₅	75	211-12	C ₁₃ H ₉ N ₃ O ₂ (239)	65.19 (65.22)	3.72 (3.76)	17.60 (17.56)
b.	H	2-ClC ₆ H ₄	61	205-06	C ₁₃ H ₈ N ₃ O ₂ Cl (273.5)	56.91 (57.03)	2.90 (2.92)	15.28 (15.35)
c.	H	3-NO ₂ C ₆ H ₄	65	234-36	C ₁₃ H ₈ N ₄ O ₄ (284)	54.96 (54.92)	2.77 (2.81)	19.80 (19.71)
d.	H	4-CH ₃ C ₆ H ₄	67	217-19	C ₁₄ H ₁₁ N ₃ O ₂ (253)	66.35 (66.39)	4.30 (4.34)	16.51 (16.59)
e.	H	4-BrC ₆ H ₄	70	254-56	C ₁₃ H ₈ N ₂ O ₂ Br (318)	49.00 (49.04)	2.46 (2.51)	13.16 (13.20)

TABLE - 2: Spectral data.

Compound No.	IR data (max)	PMR (ppm)
II a.	1645(C=O), 1590 (C=N) 3450 (N-H)	2.53(s,3H); 7.4(d, J=7,1H); 7.68(d, J=7,1H); 7.28(b, N-H); 8.12 (s, 1H)
c.	1640(C=O), 1580 (C=N) 3454(N-H)	6.95-7.15 (m, Ar-H, 10H); 8.15 (s, 1H, N-H); 7.22 (s, 1H), 7.56 (s, 1H, Ar-H)
d.	1640(C=O), 1575(C=N) 3380 (N-H)	7.2 (d, 2H, Ar-H); 7.50 (d, 2H, Ar-H); 8.31 (s, 1H), 7.95 (b, 1H, N-H) 7.05 (s, 5H, Ar-H).
IV a.	1630 (C=C), 1680 (C=O) 1590 (C=N), 3350 (N-H)	7.15 (s, 5H, Ar-H), 7.0 (s, 5H, Ar-H), 7.25 (b, 1H, NH), 8.18 (s, 1H),
b.	1615 (C=C), 1685 (C=O) 1600 (C=N) 3340 (N-H)	7.43 (m, 2H, 3'-H, 5'-H), 7.76 (m, 2H, 2'-H, 6'-H); 6.95 (s, 5H, Ar-H), 7.15 (s, 1H, N-H) 8.20 (s, 1H)
VI a.	1690 (C=O), 1620 (C=C) 1580(C=N), 3300-3400 (N-H)	7.90 (b, 1H, N-H), 8.20 (b, 1H, N-H), 8.25 (s, 1H), 8.15 (s, 1H), 6.95 (s, 5H, Ar-H).
d.	1685(C=C), 1690 (C=O) 1590 (C=N), 3350-3400 (N-H)	2.80 (s, 3H CH ₃), 7.25 (d, 2H, 3'-H, 5'-H), 7.45 (d, 2H, 2'-H, 6'-H), 7.35 (s, 1H), 8.15 (s, 1H), 8.70 (s, 1H), 7.13 (b, 1H, N-H), 7.23 (b, 1H, N-H).

Experimental Section:

Melting points were determined on Mel.Temp and are uncorrected. IR spectra were recorded in KBr on Perkin-Elmer spectrometer, PMR were recorded in DMSO-d₆ and CDCl₃ on Varian-Gemeini 200 spectrometer.

General procedure for the preparation of fused Pyrimidine-4(1H)-ones (II, IV & VI):

A mixture of 2-amino-3-cyano -aromatic compound (2 mmoles) was added in portion wise over 1 hr to a mildly reflecting mixture of 88% formic acid and sulfuric acid (1.0 gm) . After 15 min, the mixture was allowed to cool to 60 C , poured into crushed ice (100 gms) and allowed to stand for 15 min. The resulting precipitate was collected and washed well with water. Drying to constant weight provided **II, IV & VI** as an off white solid (80%).

Acknowledgments: One of the authors (SVSAKG) is thankful to Prof.Venkateswaran, Director, BITS, Pilani (Raj) , India, for providing the necessary research facilities.

References:

1. S.S.F.Burni, A.Castanzo, G.Guerrini, P.M.Ajello, G.Iavrsone and C.Martini, Eur.J.Med.Chem. 27, 985 (1992).
2. M.F.Hasan, A.M.Mounir,, I.Sallem,J.M.Abdul Rahman and A.E.Mohammed, Hetero cycles, 38 57 (1994).
3. N.Singhal, I.S.Gupta and P.C.Bhansal, J.Indian Chem.Soc, 61, 690 (1984).
4. A.K.Sharma, A.K.Yadav and Lalit Prakash, Ind.J.Chemistry, 34B, 740 (1990).
5. V.Neimentoskii, J.Pract.Chem, 51, 564 (1895); W.L.F.Armarego, J.Appl.Chem, 11, 70 (1961).
6. M.K.Mckee, R.L.McKee and R.W.Bost, J.Am.Chem.Soc, 69, 184 (1947).
7. K.Nagahara, K.Takagi and T.Ueda, Chem. Pharm.Bull, 24, 310 (1976).
8. J.T.Gupton, K.F.Correia, and B.S.Foster, Synth.Comm, 16, 365, (1986).
9. M.T.Bogert and W.F.Hard, J.Am.Chem.Soc,28, 94 (1906).
10. M.Akazome, J.Yamamoto, T.Kordo and Y.Wantanhe, J.Organomet.Chem, 494, (1995).
11. M.Shah, P.Patel.S.Korgaokar and H.Parekh, Indian.J.Chem.SOc, 35B, 1282 (1996).
12. U.Hanefeld, V.W.Rees, A.J.P.White and D.J.Williams, J.Chem>Soc, Per. Tran-I,1546, (1996).
13. A.H.H.Elghandour, M.K.Abrahim, F.M.M.Ali and S.M.M.Elshikh, Indian.J.Chem, 36B, 79 (1997).
14. G.A.Roth and J.J.Tai, J.Hetrtocyclic.Chem, 33, 2051 (1996).
15. U.T.Bhale Rao and A.Krishnaiah, Indian.J.Chem, 34B, 587 (1995).

Received on December 20, 1998