Reactions of 1-(2-benzothiazolyl)-4-(dicyanomethylene)-3-methyl-2-pyrazolin-5-one towards Amines

Aly H. Atta

Chemistry Department; Faculty of Education (Suez); Suez Canal University; Suez; Egypt

ABSTRACT: The reaction of 1-(2-Benzothiazolyl)-4-(dicyanomethylene)-3-methyl-2-pyrazolin-5-one (1) with amines has been investigated. Primary and secondary amines react with (1) to yield 4-substituted(aminocyanomethylene)pyrazolone derivatives (4a-j) and (6a, b) respectively. o-substituted arylamines afford 4-azolyldene pyrazolone derivatives (8a, b). Tertiary amines (9a, b) adds to the dicyano derivative (1) by stirring in acetic acid solution and in exclusion of light to give adducts (10a, b). The latter products gave the condensation products (11a, b) by heating in DMF.

Key words : 2-pyrazolin-5-one derivatives.

INTRODUCTION

Since the last few years we are interested in the reactions of pyrazolones.¹⁻⁴⁾ Owing to the vital importance of pyrazolone derivatives due to their wide applications for industrial uses⁵⁻⁷⁾ and medicinal fields⁷⁻¹⁰⁾, I aim to synthesis some heterocyclic compounds starting with 1-(2-benzothiazolyl)-4-(dicyanomethylene)-3-methyl-2-pyrazolin-5-one (1)⁴⁾ hoping that such new compounds will have certain biological effects or other applied uses.

RESULTS AND DISCUSSION

I found that compound (1) condenses with primary amines (2a-j) by attack at C- α of the dicyanomethylene moiety¹¹⁻¹³⁾ to afford (3) which loses hydrogen cyanide yielding 4-alkyl- or aryl-(aminocyanomethylene)-3-methyl-2-pyrazolin-5-one derivatives (4a-j). In the same manner secondary amines (5a,b) reacts with 1 to give the corresponding products (6a, b). Structures of (4a-j) and (6a, b) were confirmed by spectral and elemental analyses.

o-Phenylenediamine and o-aminophenol (7a, b) reacts with the dicyano derivative (1) to give 4-azolyldene pyrazolones (8a, b) through elimination of two molecules of hydrogen cyanide.

N,N-Dimethylaniline and N,N-diethylaniline as tertiary amines adds to the dicyano derivative (1) in acetic acid solution and in exclusion of light to give the disubstituted malononitriles (10a, b). These colourless adducts (10a, b) are unstable when heated in DMF, hydrogen cyanide is eliminated and deeply coloured arylcyanomethylene pyrazolone derivatives (11a, b) are formed¹⁴⁾.

6a) X=CH₂
6b) X=O

4a-j

	R		R
4a	CH ₃	f	C ₆ H ₄ OH(4)
b	C ₂ H ₅	g	Cyclohexyl
c	C ₆ H ₅	h	benzyl
d	C ₆ H ₄ CH ₃ -(4)	i	NH ₂
e	C ₆ H ₄ OCH ₃ -(4)	j	NHph

8a) X=NH
8b) X=O

10a, b

11a, b

9-11 : a) R=Me
b) R=Et

EXPERIMENTAL

All melting points are uncorrected. Infrared spectra were recorded with a Shimadzu 408 Infrared spectrometer using KBr discs. ^1H NMR spectra were measured with a Varian XL-100 spectrometer; Chemical Shifts are reported in ppm; initial standard was tetramethylsilane (δ scale). Mass spectra were obtained at Cairo University. Microanalyses were carried out at microanalysis unit at Assiut University.

1-(2-benzothiazolyl)-4-(dicyanomethylene)-3-methyl-2-pyrazolin-5-one(1) was prepared according to ref⁴⁾.

Reactions of 1-(2-benzothiazolyl)-4-(dicyanomethylene)-3-methyl-2-pyrazolin 5-one (1) with primary, secondary amines and o-substituted arylamines : General Method :

A mixture of an equimolar amount of the dicyano derivative (1) and primary, secondary amine or o-substituted arylamine (2, 5 or 7) (1 m mol) in absolute ethanol (30 ml) was stirred for 1/2-2h. The precipitated product was collected and crystallized from the proper solvent. The results are listed as the following :

(4a) : Yield 71%, pale yellow crystals, m.p. 272 °C; (ethanol). - IR : ν = 3250 cm^{-1} (NH); 1650 (CO). - MS : m/z (%) = 297 (14.6) [M^+], 282 (28.3) [$\text{M}^+ - \text{CH}_3$], Found C, 56.42; H, 3.81; N, 23.62; S, 10.70, $\text{C}_{14}\text{H}_{11}\text{N}_5\text{OS}$ (297.3), requires C 56.55 H 3.73 N 23.55 S 10.78.

(4b) : Yield 61%, pale yellow crystals, m.p. 275 °C (ethanol). - IR : ν = 3200 cm^{-1} (NH); 1680 (CO). - ^1H NMR (DMSO) : δ = 1.2 (t, 3H, CH_3), 2.1 (s, 3H, CH_3), 3.4 (q, 2H, CH_2), 7.2-8.0 (m, 4H, aromatic H), Found C 57.81 H 4.11 N 22.52 S 10.28, $\text{C}_{15}\text{H}_{13}\text{N}_5\text{OS}$ (311.4), requires C 57.86 H 4.21 N 22.49 S 10.30.

(4c) : Yield 67%, orange crystals, m.p. 222 °C (ethanol) - IR : ν 3400 cm^{-1} (NH) 1670 (CO). -MS : m/z (%) = 359 (91.4) [M^+], 267 (13.6) [$\text{M}^+ - (\text{HN} - \text{C}_6\text{H}_5)$], Found C 63.52 H 3.60 N 19.38, $\text{C}_{19}\text{H}_{13}\text{N}_5\text{OS}$ (359.4), requires C 63.50 H 3.65 N 19.49.

(4d) : Yield 69%, orange crystals, m.p. 219 °C (benzene) IR : ν 1670 cm^{-1} (CO). - ^1H NMR (CDCl_3) : δ = 2.2 (s, 3H, CH_3), 2.4 (s, 3H, CH_3), 7.2-8.1 (m, 8H, aromatic H), 12.0 (s, 1H, NH), Found C 64.42 H 4.11 N 18.67, $\text{C}_{20}\text{H}_{15}\text{N}_5\text{OS}$ (373.4), requires C 64.33 H 4.05 N 18.75.

(4e) : Yield 82%, orange crystals, m.p. 210 °C (ethanol). - IR : ν 1650 cm^{-1} ^1H NMR (CDCl_3) : δ = 2.6 (s, 3H, CH_3), 3.85 (s, 3H, OCH_3), 7.0 - 8.1 (m, 8H, aromatic H), 11.9 (s, 1H, NH). - MS : m/z (%) = 389 (56.6) [M^+], 374(11) [$\text{M}^+ - \text{CH}_3$], 358(1.2) [$\text{M}^+ - \text{OCH}_3$], Found C 61.62 H 3.92 N 17.82 S 8.09, $\text{C}_{20}\text{H}_{15}\text{N}_5\text{O}_2\text{S}$ (389.4), requires C 61.68 H 3.88 N 17.98 S 8.23.

(4f) : Yield 83%, red crystals, m.p. 296 °C (ethanol). - IR : ν 3400 cm^{-1} (NH); 1670 (CO). - ^1H NMR (DMSO) : δ = 2.4 (s, 3H, CH_3), 6.8 - 8.2 (m, 9H, 8 aromatic H + NH), 10.0 (s, 1H, OH). - MS : m/z (%) = 375 (70.5) [M^+], 358 (3.1) [$\text{M}^+ - \text{OH}$], 367(13.6) [$\text{M}^+ - \text{HNC}_6\text{H}_4\text{OH}(4)$], Found C 60.71 H 3.53 N 18.55 S 8.42, $\text{C}_{19}\text{H}_{13}\text{N}_5\text{O}_2\text{S}$ (375.4), requires C 60.79 H 3.49 N 18.66 S 8.54.

(4g) : Yield 81%, cream crystals, m.p. 224 °C (ethanol) IR : ν 1660 cm^{-1} (CO). - ^1H NMR (CDCl_3) : δ = 1.2 - 2.5 (m, 10 H, cyclohexyl H), 2.5 (s, 3H, CH_3), 7.4 - 8.2 (m, 4H, aromatic H), 10.6 (s, 1H, NH), Found C 62.52 H 5.17 N 19.07, $\text{C}_{19}\text{H}_{19}\text{N}_5\text{OS}$ (365.5), requires C 62.45 H 5.24 N 19.16.

(4h) : Yield 89%, yellow crystals, m.p. 225 °C (ethanol). IR : ν 3200 cm^{-1} (NH); 1775 (CO) - ^1H NMR (CDCl_3) : δ = 2.5 (s, 3H, CH_3), 4.8 (s, 2H, CH_2), 7.2 - 8.1 (m, 9H, aromatic H), 10.4 (s, 1H, NH), Found C 64.39 H 4.09 N 18.81 S 8.65, $\text{C}_{20}\text{H}_{15}\text{N}_5\text{OS}$ (373.4), requires C 64.33 H 4.05 N 18.75 S 8.59.

(4i) : Yield 71%, brownish yellow crystals, m.p. 189 °C (ethanol). - IR : ν = 3350, 3250 cm^{-1} (NH_2), 2200 (CN). - MS : m/z (%) = 298 (100) [M^+], 282(13) [$\text{M}^+ - \text{NH}_2$], Found C 52.51 H 3.29 N 28.07, $\text{C}_{13}\text{H}_{10}\text{N}_6\text{OS}$ (298.3), requires C 52.34 H 3.38 N 28.17.

(4j) : Yield 67%, cream crystals, m.p. 168 °C (ethanol). - IR : ν = 3400 cm^{-1} (NH), 2210 (CN). - $^1\text{H-NMR}$ (CDCl_3) : δ = 2.4 (s, 3H, CH_3), 7.05 - 7.95 (m, 11H, 9 aromatic H + 2 NH), Found C 60.91 H 3.67 N 22.36., $\text{C}_{19}\text{H}_{14}\text{N}_6\text{OS}$ (374.4), requires C 60.95 H 3.77 N 22.45.

(6a) : Yield 61%, cream crystals, m.p. 215 °C (methanol). - IR : ν 2220 cm^{-1} (CN), 1680 (CO). - MS : m/z (%) = 351 (62.4) [M^+], Found C 61.41 H 4.73 N 19.81, $\text{C}_{18}\text{H}_{17}\text{N}_5\text{OS}$ (351.4), requires C 61.52 H 4.88 N 19.93.

(6b) : Yield 57%, cream crystals, m.p. 197 °C (ethanol) IR : ν 1680 cm^{-1} (CO). - $^1\text{H NMR}$ (CDCl_3) : δ = 2.5 (s, 3H, CH_3), 3.95 (s, 8H, 4 CH_2 of morpholine), 7.2 - 8.0 (m, 4H, aromatic H), Found C 57.82 H 4.22 N 19.71, $\text{C}_{17}\text{H}_{15}\text{N}_5\text{O}_2\text{S}$ (353.4), requires C 57.78 H 4.28 N 19.82.

(8a) : Yield 71%, reddish crystals, m.p. 322 °C (ethanol) IR : ν 1640 (CO). - $^1\text{H NMR}$ (DMSO) : δ = 2.4 (s, 3H, CH_3), 7.3 - 8 (m, 8H, aromatic H). - MS : m/z (%) 347 (100). [M^+], Found C 62.26 H 3.71 N 20.02 S 9.18, $\text{C}_{18}\text{H}_{13}\text{N}_5\text{OS}$ (347.4), requires C 62.23 H 3.77 N 20.16 S 9.23.

(8b) : Yield 79%, orange crystals, m.p. 220 °C (ethanol) IR : ν 3100 cm^{-1} (NH), 1660 (CO). - MS : m/z (%) 348 (100) [M^+], Found C 62.15 H 3.39 N 16.12 S 9.11, $\text{C}_{18}\text{H}_{12}\text{N}_4\text{O}_2\text{S}$ (348.4), requires C 62.06 H 3.47 N 16.08 S 9.20.

Preparation of adducts (10a,b)-General Method : N,N-dialkyl aniline (9a, b) (10 m mol) was added slowly to a solution of 10 m mol of 1-(2-benzothiazolyl)-4-(dicyanomethylene)-3-methyl-2-pyrazolin-5-one (1) in acetic acid (15 ml) with magnetic stirrer at room temperature with the exclusion of light. Stirring was continued 2h. The precipitated product by dilution was collected, washed with water, purified by dissolving in acetic acid and reprecipitated by addition of water. The products were colourless powders, they were dried at room temperature in vacuo and in the dark. $^1\text{H NMR}$ and mass spectra could not be recorded due to the instability of the products.

2-[1-(2-Benzothiazolyl)-5-hydroxy-3-methyl-1H-pyrazol-4-yl]-N,N-dimethylanilin-4-yl)-malononitrile (10a) : Yield, 89%, colourless powder, m.p. 163 -165 °C. IR : ν = 3450 cm^{-1} (OH), Found C 63.71 H 4.31 N 20.17 S 7.59, $\text{C}_{22}\text{H}_{18}\text{N}_6\text{OS}$ (414.5), requires C 63.75 H 4.38 N 20.28 S 7.74.

2-[1-(2-Benzothiazolyl)-5-hydroxy-3-methyl-1H-pyrazol-4-yl]-N,N-diethylanilin-4-yl)-malononitrile (10b) : Yield 86%, colourless powder, m.p. 174 - 176 °C. IR : ν = 3400 cm^{-1} (OH). Found C 65.07 H 5.09 N 19.12 S 7.13, $\text{C}_{24}\text{H}_{22}\text{N}_6\text{OS}$ (442.5), requires C 65.14 H 5.01 N 18.99 S 7.25.

Preparation of the condensation products (11a, b) : A solution of (10a, b) in DMF (1.5 gm in 10 ml) was heated at 100 °C for 1/2 h. After cooling the precipitated product by dilution with water was collected and crystallized from dilute DMF as deep violet crystals.

(11a) : Yield 92%, m.p. 158 °C. - IR : ν = 1670 cm^{-1} (C=O, pyrazolone). $^1\text{H NMR}$ (DMSO) : δ = 2.5 (s, 3H, CH_3), 3.4 (s, 6H, $\text{N}(\text{CH}_3)_2$), 7.0 - 8.2 (m, 8H, aromatic H). - MS : m/z (%) = 387 (75.9%) [M^+], 372(17.7) [$\text{M}^+ - \text{CH}_3$], 343(8.1) [$\text{M}^+ - \text{N}(\text{CH}_3)_2$], Found C 65.21 H 4.37 N 18.13, $\text{C}_{21}\text{H}_{17}\text{N}_5\text{OS}$ (387.5), requires C 65.10 H 4.42 N 18.07.

(11b) : Yield 90%, m.p. 165 °C. IR : ν 1680 cm^{-1} (C=O, pyrazolone). - ^1H NMR (DMSO) : δ = 1.1 (t, 6H, 2 CH_3CH_2), 2.5 (s, 3H, CH_3); 3.4 (q, 4H, 2 CH_2CH_3), 6.8 - 8.2 (m, 8H, aromatic H). - MS : m/z (%); 415 (44.8) [M^+], 400(100) [M^+-CH_3], 386(1.6) [$\text{M}^+-\text{C}_2\text{H}_5$], 434(3.5) [$\text{M}^+-\text{N}(\text{C}_2\text{H}_5)_2$], Found C 66.39 H 5.12 N 16.79, $\text{C}_{23}\text{H}_{21}\text{N}_5\text{OS}$ (415.5), requires C 66.48 H 5.09 N 16.85.

ACKNOWLEDGEMENT

The author thanks prof. S.A. Metwally, Chemistry Department, Faculty of Science, Assiut University, for his kind help. Thanks are also to microanalysis unit of Assiut University for microanalytical data and also to mass spectrum unit of Cairo University for mass spectral data.

REFERENCES

- 1) M.I. Younes, S.A. Metwally, A.H. Atta, *synthesis* 704(1990).
- 2) M.I. Younes, A.H. Atta, S.A. Metwally, *J. of the Chinese Chem. Soc.* **37**, 617(1990).
- 3) M.E. Abd Elmonem, M.I. Younes, S.A. Metwally, A.H. Atta, *Collec Czech. Chem. Commun.* **56**, 1977(1991).
- 4) A.H. Atta, R.F. Fandy, M.I. Younes, S.A. Metwally, *Bull. J. of Assiut*, **25**, 35 (1996)
- 5) C. Prakash, G.S. Saharia, H.R. Sharma, *J. Indian Chem. Soc.* **54**, 381(1977).
- 6) G. Tacconi, G. Marinoni, P.P. Righetti, G. Desimoni, *J. Prakt. Chem.* **322**, 674(1980).
- 7) A.I. Koraiem, *J. Prakt. Chem.* **326**, 695-, 811- (1984).
- 8) A. Nayak, A.S. Mittra, *J. Indian Chem. Soc.* **57**, 643(1980).
- 9) A.M. Ismaiel, M.Y. Yousif, M.A. Metwally, M. El-Kerdawy, *Indian J. Chem.* **25 B** (12), 1238(1986).
- 10) F.A. Adam, M.T. El-Haty, A.I. Koraiem, A.H. Amrallah, *J. Indian Chem. Soc.* **66**, 157(1989).
- 11) H. Junek, H. Aigner, H. Fisher-Colbrie, *Monatsh. Chem.* **103** 639. (1972)
- 12) H. Junek, H. Fisher-Colbrie, H. Sterk, *Chem. Ber.* **110**, 2276. (1977)
- 13) S.A.M. Metwally, G.M. El Naggar, M.I. Younis, T.I. El-Emary, M.H. Elnagdi, *Liebigs Ann. Chem.*, 1037 (1989).
- 14) R. Dworczak, W.M.F. Fabian, H. Sterk, C. Kratky, H. Junek, *Liebigs Ann. Chem.*, 7. (1992).

Received on February 20, 1999