SYNTHESIS OF 6-[(4-CHROMON-3-YL-BENZOPYRANO[4,3-b] PYRIDIN-2-YL)]-2H-[1,4]-BENZOXAZIN-3(4H)-ONES

G. Jagath Reddy *, D. Latha, C. Thirupathaiah and K. Srinivasa Rao

R & D Laboratories, Dr. Jagath Reddy's Heterocyclics, 81, S.V.Co-op Industrial Estate, Balanagar, Hyderabad – 500 037, India. e-mail-jagathreddy@usa.net; Fax #91-40-23773487.

and

Md. Khalilullah

Department of Chemistry, Jawaharlal Nehru Technological University, Hyderabad-500072, India.

Abstract:

A series of new 6-[(4-chromon-3-yl-benzopyrano[4,3-b]pyridin-2-yl)]-2H-1,4-benzoxazin-3(4H)-ones (5a-l) have been prepared.

Introduction:

A variety of heterocyclic substituted benzoxazinones have been reported to possess interesting biological activities. For example 6-tetrahydrophthalimido benzoxazinone is a herbicide¹. α -Adrenoceptor activities have been associated with imidazolylaminobenzoxazinones². Several imidazolylmethyl³ and thiazolyl benzoxazinones⁴ exhibited significant antifungal and antiinflammatory activities. Bemoradan⁵ is a pyridazinylbenzoxazinone derivative, which exhibited hypotensive, blood platelet aggregation inhibition, positive chronotropic and ionotropic activities. Furthermore, benzopyranopyridines⁶ and chromones⁷ are compounds of considerable interest because of their pharmacological properties. In view of this, and in continuation of our work on chromone based heterocycles⁸ it was considered of interest to synthesize the title compounds in which all the three pharmacophores are present.

1,2,3,5 R₁ = H, CH₃, Cl, F and R₂=H, CH₃, F

SCHEME - 1

Results and Discussion

The required starting materials, benzopyranomethynylbenzopyranones(3a-l) were prepared according to the method reported earlier⁹. Thus reaction of chromanones(1a-d) with appropriately substituted 3-formylchromones(2a-c) in refluxing ethanol in presence of triethylamine gave 3a-l in good yields. The other starting material 3-oxo-2H-[1,4]-benzoxazin-3,4-dihydro-6-acetylpyridinium chloride (4) was prepared by reaction of 6-chloroacetyl-2H-[1,4]-benzoxazinone⁴ with pyridine at 100°C. Reaction of compound 4 with various benzopyranomethynylbenzopyranones(3a-l) in the presence of ammonium acetate in acetic acid under Krohnke's condition¹⁰ gave the title 6-(chromonyl benzopyranopyridinyl)-benzoxazinones(5a-l) (Scheme-1) in good yields as crystalline solids. The compounds are homogeneous on TLC. The IR spectrum showed characteristic peaks at 1705 and 1645 cm⁻¹ for lactam carbonyl and pyrone carbonyl. ¹H NMR spectrum showed two singlets at δ 4.6 and δ 5.2 for -OCH₂ of pyrone and -OCH₂ of benzoxazine ring respectively.

All the compounds reported in the present work (Table-1), were characterized by their IR, NMR and Mass spectra and supported by elemental analysis.

TABLE 1- PHYSICAL DATA OF 6(CHROMONYL BENZOPYRANOPYRIDINYL)-[1,4]BENZOXAZINONES-5a-l

Compound	R ₁	R ₂	Yield %	mp
5a	H	H	63	>300 ^o
5b	H	CH ₃	57	>300 ^o
5c	H	F	58	>300 ^o
5d	CH ₃	H	61	>300 ^o
5e	CH ₃	CH ₃	45	>300 ^o
5f	CH ₃	F	67	>300 ^o
5g	Cl	H	62	>300 ^o
5h	Cl	CH ₃	51	>300 ^o
5i	Cl	F	65	>300 ^o
5j	F	H	60	>300 ^o
5k	F	CH ₃	47	>300 ^o
5l	F	F	56	>300 ^o

Experimental

Melting points were carried out in an open capillary and are uncorrected. NMR spectra were recorded on 200 Mz spectrophotometer and chemical shift values are expressed in δ (ppm).

General procedure for preparation of 3-(4-oxo-4H-1-benzopyrano-3-methynyl)-2,3-dihydro-4H-1-benzopyran-4-ones (3a-l)

A mixture of 3-formylbenzopyran-4-one (**2**, 0.1 mole) and 4H-1-benzopyran-4-one (**1**, 0.15 mole) in ethanol (100 ml) and triethylamine (50 ml) was refluxed for 3 hrs. The reaction was followed by TLC. At the end of the reaction it was cooled and the solid was filtered washed and recrystallized with ethanol to give pure **3** in 67-80% yields.

General procedure for preparation of (3-oxo-2H[1,4]-benzoxazin-6-yl)acetylpyridiniumchloride (**4**)

A mixture 6-chloroacetylbenzoxazinone (**2**, 2.5 g, 0.1 mole), pyridine (100 ml) and toluene (100 ml) was refluxed for 1 hr. The separated solid was filtered and washed with toluene to give pure **4** as white crystalline solid (27.4 g, 90%, mp>300°C).

6-[(4-Chromon-3-yl)-benzopyrano[4,3-b]pyridin-2-yl]-2H-[1,4]-benzoxazin-3(4H)-one (**5a**)

A mixture of **4** (3.04 g, 0.01 mole), **3a** (3.0 gm, 0.01 mole) and ammonium acetate (4.62 gm, 0.06 mole) in glacial acetic acid (30 ml) was refluxed under stirring for 2 hrs. The separated solid was filtered, washed with water, ethanol, dried and recrystallized from DMSO to give pure **5a** (3.0 g, 63%) mp>300°C IR: 1705, 1645 cm⁻¹; NMR(DMSO-d₆): δ 4.6(s, 2H, OCH₂), 5.2(s, 2H, OCH₂), 6.9-8.0(m, 12H, ArH), 8.5(s, 1H, chromone H), 10.6(s, 1H, NH). Mass m/z at (M⁺) 474. Anal. Calcd for C₂₉H₁₈N₂O₅: C, 73.41; H, 3.79; N, 5.90. Found: C, 73.37; H, 3.82; N, 5.92

References:

1. T. Hoga, H. Naguno, M. Sato and K. Morita J.P. 6251, 685, *Chem. Abstr*; **108**, 5611g **1988**.
2. C.B. Chapreo, R.C.M.Batler, D.England, P.L.Myen, AG. Rnach, C.F.C. Smith, M.R. Stillings and I.F. Talloch, *J. Med. Chem* **32**, 1627 **1989**.
3. C.V. Reddy Sastry, K. Srinivasa Rao, K. Rastogi, M.L. Jain and G.S. Reddy, *Indian J. Chem*; **28B**, 1096 **1989**.
4. C.V.Reddy Sastry, O.P. Bansal, K. Srinivasa Rao, P. Pulla Rao, M.L. Jain and D.R. Shridhar., *Indian J. Chem*; **26B**, 662, **1987**.
5. D.W. Combs, MS. Rampulla, S.C.Bell, D.H.Klanbost, A.J. Tobia, R. Farotica, B. Heartlein, CL.Weiss and J.B. Moora, *J. Med. Chem*, **33**, 380 **1990**.
6. A. Nohara, K. Ishiguru, K.Ukawa, H. Sugihara, Y. Maki and Y. Sanno, *J. Med. Chem* **28**, 559 **1985**
7. K.A. Thakoor and C.H. Gill, *J. Ind. Chem. Soc*, **LX**, 668 **1983**.
8. G. Jagath Reddy, D. Latha and C. Thirupathaiah, *Heterocyclic. Commun.* (in press) **2003**.
9. K.V. Reddy, G. Sabitha and A.V. Subba Rao., *Org. Prep. Proced. Int* **28** (3), 325 **1996**.
10. F. Krohnke, *Synthesis*, **1**, **1976**.

Received on March 12, 2003