SYNTHESIS OF A PERHYDROINDOLIC ANALOG OF KAINIC ACID

Mathieu Danel, Akram Hijazi, Roland Barret*

Laboratoire de Chimie Thérapeutique, EA 3741, Faculté de Pharmacie, Université Claude Bernard, 8 Avenue Rockefeller, F69373 Lyon Cedex 08

Abstract : Synthesis of perhydroindolyl-2-carboxy-3-acetic acid, a strained analog of kainic acid, is described. This compound was obtained in 7 steps by the use of the Barco procedure from nitrocyclohexene.

Introduction

In the recent years there has been a growing interest in the L-glutamate receptors in the central nervous system. This neurotransmitter plays a role of utmost importance in many physiological processes such as neuronal plasticity, memory and learning (1) but an excessive L-glutamate release can result in neuronal cell death by excitotoxicity.

Among the different sub-types of receptors of L-glutamate (AMPA, kainic acid, MND, metabotropic) (2) kainic acid **1** is an agonist of kainic acid receptor and kaitocephalin **2** (3) is an antagonist but the selectivity is low and these compounds have affinity in particular for the AMPA receptor.

With the aim to obtain a selective affinity for the kainic acid receptor, we had projected to synthesize the analog **3** of kainic acid.

Result and discussion :

The precursor **5** of **4** is obtained by the use of Barco's process (4) with two Michael reactions with 1-nitro-cyclohexene and the amino-alcohol-ester **4**. Some chemical modifications of **5** give **3**.

-Synthesis of the amino-alcohol-ester 4

The ester **4** was obtained from serine in seven steps.

Scheme 1 – 1- CH_3OH , CH_3COCl , 0°C , then reflux for 3h. 2- a- Et_3N , $\text{C}_6\text{H}_5\text{CHO}$ –b- NaBH_4 . 3- $(\text{Boc})_2\text{O}$, Et_3N , THF. 4- TBDMSiCl , Et_3N , DMAP, CH_2Cl_2 .

D-serine gave the ester **6** as hydrochloride quantitatively. After reductive alkylation (PhCHO - NaBH_4) the secondary amine **7** was transformed in the carbamate **8** and then alcohol function was protected as silyl ether **9**.

To obtain aldehyde function, ester **9** was reduced in alcohol by the use of lithium aluminium hydride at -50°C and then oxidized in aldehyde by the Swern reaction. During the reduction step a few quantity of aldehyde **11** was isolated.

The aldehyde **11** was not purified and directly reacted with (carboethoxymethylene) triphenylphosphorane to give **12** (trans isomer) in a good yield.

Scheme 2 – 1- LiAlH_4 , THF. 2- Et_3N , DMSO, $(\text{COCl})_2$. 3- $(\text{C}_6\text{H}_5)_3\text{PCHCO}_2\text{C}_2\text{H}_5$, CH_2Cl_2 .

The aldehyde obtained during the reduction step gave **12** to increase yield. Thus **12** was obtained with a good yield (93% from **9**). At last, the protective group was cleaved with trifluoroacetic acid to give **4**.

Scheme 3 – $\text{CF}_3\text{CO}_2\text{H}$, CH_2Cl_2

The global yield of the synthesis of **4** from serine in 7 steps was 24%.

-Condensation of **4** with 1-nitrocyclohexene.

Nitrocyclohexene and **4** were stirred for seven days at room temperature. After this time, **13** was obtained with a 65% yield. All modifications of these conditions (time, temperature, solvent....) lead to low yields.

This reaction is a double-Michael addition:

The nitro group was eliminated by reflux into toluene with tributyltin hydride in the presence of catalytic quantity of AIBN to give **14**.

Debenzylation succeeded with Pd-C 5% in alcohol under hydrogen pressure (25 bars) on the condition that one or two drops of 10% hydrochloric acid be added. In the absence of acid no reaction took place.

It's known that in case of amino-alcohol the amino group must be protected with *tert*-butoxycarbamate function during the oxidation of the alcohol function into acid (4,5); then, after debenzylation and protection of amine with *tert*-butoxycarbonyl group, the oxidation with PDC gave acid **16**.

Scheme 4 – 1- Bu_3SnH , AIBN, toluene, reflux. 2- H_2 25 bars, Pd.C, $\text{C}_2\text{H}_5\text{OH}$, HCl 10% 2 drops. 3- $(\text{Boc})_2\text{O}$, Et_3N , THF. 4- PDC, CH_2Cl_2 .

This acid-ester **16** was hydrolysed by LiOH and deprotected with trifluoroacetic acid.

Scheme 4- 1- LiOH, H_2O , THF. 2- $\text{CF}_3\text{CO}_2\text{H}$, CH_2Cl_2 .

Conclusion

The compound **4** was obtained with a global yield of 11% from nitrocyclohexene. Now, we synthesize other compounds by this strategy and we have to test these ones with different L-glutamate sub-type receptors

Aknowledgements

The authors thank ANVAR and EZUS for their financial support.

References

- (1) N.S. Nakanishi, *Science* **258**,31 (1992); Z.A. Bartolotto, Z.I. Bashir, C.H. Davies, G.L. Collingridge, *Nature* **368**, 740 (1994).
- (2) M.D. Cunningham, J.W. Ferkany, S.J. Enna, *Life Sciences* **54**, 135 (1994). M. Hollmann, S. Heinemann, *Annu. Rev. Neurosc.* **17**, 31 (1994).
- (3) K. Shin-Ya, J.S. Kim, K. Furibata, Y. Hayakawa, H. Seto, *Tetrahedron Lett.* **38**, 7079 (1997). D. Ma, J. Yang, *J. Amer. Chem. Soc.* **123**, 9706 (2001).
- (4) A. Barco, S. Benetti, A. Casolari, G.P. Pollini, G. Spalluto, *Tetrahedron Lett.* **31**, 4917 (1990). A. Barco, S. Benetti, G. Spalluto, *J. Org. Chem.* **57**, 6279 (1992).
- (5) F.J. Sardina, M.H. Howard, A.M.P. Koshinen, H. Rapoport, *J. Org. Chem.* **54**, 4654 (1989).
- (6) Analytical data **4**: yellow oil, $^1\text{H-NMR}$ (CDCl_3 , 200MHz) δ (ppm) : 1.05-2.45 (m, 9H, H-3a and $(\text{CH}_2)_4$), 2.45-3.00 (m, 3H, H-3, CH_2CO_2), 3.80-4.50 (m, 2H, H-2 and H-7a), 10.16 and 12.01 (2s, 4H, 2 COOH, 2NH). MS (EI): 228 (M^+).

Received on October 4, 2004