Research Article

Nils Waterstraat

A remark on bifurcation of Fredholm maps

DOI: 10.1515/anona-2016-0067

Received March 20, 2016; revised June 4, 2016; accepted June 5, 2016

Abstract: We modify an argument for multiparameter bifurcation of Fredholm maps by Fitzpatrick and Pejsachowicz to strengthen results on the topology of the bifurcation set. Furthermore, we discuss an application to families of differential equations parametrised by Grassmannians.

Keywords: Bifurcation, Fredholm maps, index bundle

MSC 2010: Primary 58E07; secondary 58J20, 19L20

1 Introduction

Let X and Y be Banach spaces, $\mathcal{O} \subset X$ an open neighbourhood of $0 \in X$ and Λ a compact topological space. A family of C^1 -Fredholm maps is a map $f: \Lambda \times \mathcal{O} \rightarrow Y$ such that each $f_\lambda := f(\lambda, \cdot): \mathcal{O} \rightarrow Y$ is C^1 and the Fréchet derivatives $D_u f_\lambda \in \mathcal{L}(X, Y)$ are Fredholm operators of index 0, which depend continuously on $(\lambda, u) \in \Lambda \times \mathcal{O}$ with respect to the operator norm on the space of bounded linear operators $\mathcal{L}(X, Y)$. In what follows, we assume that $f(\lambda, 0) = 0$ for all $\lambda \in \Lambda$, and we say that $\lambda^* \in \Lambda$ is a *bifurcation point* if every neighbourhood of $(\lambda^*, 0)$ in $\Lambda \times \mathcal{O}$ contains an element (λ, u) such that $u \neq 0$ and $f(\lambda, u) = 0$. By the implicit function theorem, the linear operator

$$L_{\lambda^*} := D_0 f_{\lambda^*}: X \rightarrow Y$$

is non-invertible if λ^* is a bifurcation point, and so a non-trivial kernel of some $L_\lambda = D_0 f_\lambda$ is a necessary assumption for the existence of a bifurcation point. Lots of efforts have been made to obtain sufficient criteria for the existence of bifurcation points by topological methods (cf., e.g., [1, 4, 5, 9]). Here we focus on the articles [16] and [17] of Pejsachowicz, which are the state of the art of a line of research on which he has worked for at least three decades (cf. [7, 13–15, 18]). In order to explain these results briefly, let us recall that every family $\{L_\lambda\}_{\lambda \in \Lambda}$ of Fredholm operators has an *index bundle*, which is a K -theory class $\text{ind } L$ in $\widetilde{KO}(\Lambda)$ that generalises the integral index for Fredholm operators to families. Moreover, Atiyah introduced in the sixties the J -group $J(\Lambda)$ and a natural epimorphism $J: \widetilde{KO}(\Lambda) \rightarrow J(\Lambda)$, which is called the generalised J -homomorphism. Pejsachowicz' bifurcation theorem [16] reads as follows.

Theorem 1.1. *Let $f: \Lambda \times \mathcal{O} \rightarrow Y$ be a family of C^1 -Fredholm maps, parametrised by a connected finite CW-complex Λ , such that $f(\lambda, 0) = 0$ for all $\lambda \in \Lambda$. If there is some $\lambda_0 \in \Lambda$ such that L_{λ_0} is invertible and $J(\text{ind } L) \neq 0$, then the family f has at least one bifurcation point from the trivial branch $\Lambda \times \{0\}$.*

Let us point out that the non-triviality of $J(\text{ind } L)$ can be obtained by characteristic classes, what paves the way for obtaining bifurcation invariants by methods from global analysis. Pejsachowicz has obtained striking results for bifurcation of families of nonlinear elliptic boundary value problems along these lines in [16].

The paper [17] deals with the question if $J(\text{ind } L)$ not only provides information about the existence of a single bifurcation point, but if we can also get results about the dimension and topology of the set of all bifurcation points $B(f) \subset \Lambda$ of f . Let us recall that the *covering dimension* $\dim(\mathcal{X})$ of a topological space $\mathcal{X}$ is the

minimal value of $n \in \mathbb{N}$ such that every finite open cover of $\mathcal{X}$ has a finite open refinement in which no point is included in more than $n + 1$ elements. In what follows, for $k \in \mathbb{N}$, we denote by $w_k(\text{ind } L) \in H^k(\Lambda; \mathbb{Z}_2)$ the Stiefel–Whitney classes and by $q_k(\text{ind } L) \in H^{2(p-1)k}(\Lambda; \mathbb{Z}_p)$ the Wu classes of the index bundle $\text{ind } L \in \widetilde{KO}(\Lambda)$, where in the latter case we assume that $\text{ind } L$ is orientable. Pejsachowicz' bifurcation theorem [17], which is based on his joint paper [7] with Fitzpatrick, reads as follows.

Theorem 1.2. *Let Λ be a compact connected topological manifold of dimension $n \geq 2$ and let $f: \Lambda \times \mathcal{O} \rightarrow Y$ be a continuous family of C^1 -Fredholm maps verifying $f(\lambda, 0) = 0$, and such that there is some $\lambda_0 \in \Lambda$ for which L_{λ_0} is invertible.*

- (i) *If Λ and $\text{ind } L$ are orientable and, for some odd prime p , there is $1 \leq k \leq \frac{n-1}{2(p-1)}$ such that $0 \neq q_k(\text{ind } L) \in H^{2(p-1)k}(\Lambda; \mathbb{Z}_p)$, then the covering dimension of the set $B(f)$ is at least $n - 2(p-1)k$.*
 - (ii) *If $0 \neq w_k(\text{ind } L) \in H^k(\Lambda; \mathbb{Z}_2)$ for some $1 \leq k \leq n-1$, then the dimension of $B(f)$ is at least $n-k$.*
- Moreover, either the set $B(f)$ disconnects Λ or it cannot be deformed in Λ into a point.*

Note that if $\text{ind } L$ is non-orientable, and so (i) cannot be applied, then $w_1(\text{ind } L) \neq 0$ and we obtain by (ii) the strongest result that Theorem 1.2 can yield. Moreover, as in [16], the characteristic classes of $\text{ind } L$ in Theorem 1.2 can be computed for families of elliptic boundary value problems by methods from global analysis.

A little downside of the final statement on the topology of $B(f)$ in Theorem 1.2 is that there is no a priori knowledge about which of the alternatives hold. The aim of this short article is to show that some simple modifications in Pejsachowicz' proof of Theorem 1.2 show that actually $B(f)$ is never a contractible topological space. Note that this is slightly weaker than the second alternative in Theorem 1.2 as every contractible subspace of Λ can also be deformed in Λ into a point.

After having stated and proved our version of Theorem 1.2 in the following section, we discuss an application to families of boundary value problems of ordinary differential equations.

2 Theorem and proof

2.1 Statement of the theorem

We denote by $\Phi(X, Y)$ the subspace of $\mathcal{L}(X, Y)$ consisting of all Fredholm operators, and so the derivatives $L_\lambda = D_0 f_\lambda: X \rightarrow Y$ define a family $L: \Lambda \rightarrow \Phi(X, Y)$ parametrised by the compact space Λ . Atiyah and Jänich introduced independently the *index bundle* for families of Fredholm operators (cf., e.g., [2, 21]). Accordingly, there is a finite dimensional subspace $V \subset Y$ such that

$$\text{im } L_\lambda + V = Y, \quad \lambda \in \Lambda.$$

Hence, if we denote by P the projection onto V , then we obtain a family of exact sequences of the form

$$X \xrightarrow{L_\lambda} Y \xrightarrow{I_Y - P} \text{im}(I_Y - P) \rightarrow 0,$$

and so a vector bundle $E(L, V)$ consisting of the union of the kernels of the maps $(I_Y - P) \circ L_\lambda$, $\lambda \in \Lambda$. As L is Fredholm of index 0, the dimensions of $E(L, V)$ and V coincide. Hence, if $\Theta(V)$ stands for the product bundle $\Lambda \times V$, then we obtain the reduced KO -theory class

$$\text{ind}(L) := [E(L, V)] - [\Theta(V)] \in \widetilde{KO}(\Lambda),$$

which is called the *index bundle* of L . Let us mention for later reference that the index bundle is natural, i.e., if Λ' is another compact topological space and $f: \Lambda' \rightarrow \Lambda$ a continuous map, then $f^*L: \Lambda' \rightarrow \Phi(X, Y)$, defined by $(f^*L)_\lambda = L_{f(\lambda)}$, is a family of Fredholm operators parametrised by Λ' and

$$\text{ind}(f^*L) = f^* \text{ind}(L). \quad (2.1)$$

Two vector bundles E and F over Λ are called *fibrewise homotopy equivalent* if there is a fibre preserving homotopy equivalence between their sphere bundles $S(E)$ and $S(F)$. Moreover, E and F are *stably fibrewise*

homotopy equivalent if $E \oplus \Theta(\mathbb{R}^m)$ and $F \oplus \Theta(\mathbb{R}^n)$ are fibrewise homotopy equivalent for some non-negative integers m, n . The quotient of $\widetilde{KO}(\Lambda)$ by the subgroup generated of all $[E] - [F]$ where E and F are stably fibrewise homotopy equivalent is denoted by $J(\Lambda)$, and the quotient map $J: \widetilde{KO}(\Lambda) \rightarrow J(\Lambda)$ is called the *generalised J -homomorphism* (cf. [2]). Very little is known about $J(\Lambda)$ as these groups are notoriously hard to compute.

Note that in order to apply Theorem 1.1 it is not necessary to know $J(\Lambda)$. All what is needed is a way to decide if $J(\text{ind } L)$ is non-trivial, which can be done by *spherical characteristic classes*. We denote by $w_k(E) \in H^k(\Lambda; \mathbb{Z}_2)$ the *Stiefel–Whitney classes* of real vector bundles E over Λ , and by $q_k(E) \in H^{2(p-1)k}(\Lambda; \mathbb{Z}_p)$ the *Wu classes* of (real) orientable vector bundles E over Λ , where p is an odd prime (cf. [12]). By the naturality of characteristic classes, if Λ' is another compact space and $f: \Lambda' \rightarrow \Lambda$ is continuous, then

$$w_k(f^*E) = f^*w_k(E), \quad q_k(f^*E) = f^*q_k(E), \quad k \in \mathbb{N}. \quad (2.2)$$

As w_k and q_k are invariant under addition of trivial bundles, they are well defined on $\widetilde{KO}(\Lambda)$. Moreover, they only depend on the stable fibrewise homotopy class of the associated sphere bundles, and so they even factorise through $J(\Lambda)$. In particular, they detect elements having a non-trivial image under the J -homomorphism, i.e., $J(\text{ind } L) \neq 0$ if $w_k(\text{ind } L)$ or $q_k(\text{ind } L)$ does not vanish for some k .

The aim of this article is to prove the following modification of Theorem 1.2.

Theorem 2.1. *Let Λ be a compact connected topological manifold of dimension $n \geq 2$, and let $f: \Lambda \times \mathcal{O} \rightarrow Y$ be a continuous family of C^1 -Fredholm maps verifying $f(\lambda, 0) = 0$. We denote by $L_\lambda := D_0 f_\lambda$ the Fréchet derivative of f_λ at $0 \in \mathcal{O}$, and we assume that there is some $\lambda_0 \in \Lambda$ for which L_{λ_0} is invertible.*

- (i) *If Λ and $\text{ind } L$ are orientable and, for some odd prime p , there is $1 \leq k \leq \frac{n-1}{2(p-1)}$ such that $0 \neq q_k(\text{ind } L) \in H^{2(p-1)k}(\Lambda; \mathbb{Z}_p)$, then the covering dimension of the set $B(f)$ is at least $n - 2(p-1)k$.*
 - (ii) *If $0 \neq w_k(\text{ind } L) \in H^k(\Lambda; \mathbb{Z}_2)$ for some $1 \leq k \leq n-1$, then the dimension of $B(f)$ is at least $n - k$.*
- Moreover, the set $B(f)$ is not contractible to a point.

Let us point out once again that the novelty in our approach to this theorem is the result that $B(f)$ is never a contractible space.

2.2 Proof of Theorem 2.1

In this section we denote by $\check{H}^k(\mathcal{X}; G)$ the Čech cohomology groups of a topological space $\mathcal{X}$ with respect to the abelian coefficient group G . Moreover, we use without further reference the fact that $\dim(\mathcal{X}) \geq k$ if $\check{H}^k(\mathcal{X}; G) \neq 0$ (cf. [8, §VIII.4.A]).

Proof of Theorem 2.1. We split the proof into two parts depending on whether or not $\Lambda \setminus B(f)$ is connected.

Case 1: $\Lambda \setminus B(f)$ is not connected. In this case the reduced homology group $\tilde{H}_0(\Lambda \setminus B(f); \mathbb{Z}_2)$ is non-trivial as its rank is one less than the number of connected components of $\Lambda \setminus B(f)$. By the long exact sequence of reduced homology (cf. [6, §IV.6])

$$\cdots \rightarrow H_1(\Lambda, \Lambda \setminus B(f); \mathbb{Z}_2) \rightarrow \tilde{H}_0(\Lambda \setminus B(f); \mathbb{Z}_2) \rightarrow \tilde{H}_0(\Lambda; \mathbb{Z}_2) = 0,$$

where we used that Λ is connected by assumption, it follows that there is a surjective map

$$H_1(\Lambda, \Lambda \setminus B(f); \mathbb{Z}_2) \rightarrow \tilde{H}_0(\Lambda \setminus B(f); \mathbb{Z}_2),$$

which shows that $H_1(\Lambda, \Lambda \setminus B(f); \mathbb{Z}_2)$ is non-trivial. As $B(f)$ is closed, we obtain by Poincaré–Lefschetz duality (cf. [6, Corollary VI.8.4]) an isomorphism

$$H_1(\Lambda, \Lambda \setminus B(f); \mathbb{Z}_2) \xrightarrow{\cong} \check{H}^{n-1}(B(f); \mathbb{Z}_2),$$

and so $\check{H}^{n-1}(B(f); \mathbb{Z}_2) \neq 0$. Consequently, as $n \geq 2$, $B(f)$ is not contractible to a point and, moreover, we obtain $\dim B(f) \geq n-1$, which is greater or equal to any of the lower bounds stated in Theorem 2.1.

Note that, as we use $\mathbb{Z}_2$ coefficients, Λ does not need to be orientable in this part of the proof and so the argument works under either of the assumptions (i) or (ii).

Case 2: $\Lambda \setminus B(f)$ is connected. We only give the proof under the assumption (i), as the argument for (ii) is very similar and even simpler, since we do not need to take orientability into account. Let us point out that this second part of the proof follows quite closely [17]. A similar argument can also be found in [19].

As $\mathbb{Z}_p$ is a field, the duality pairing

$$\langle \cdot, \cdot \rangle : H^{2(p-1)k}(\Lambda; \mathbb{Z}_p) \times H_{2(p-1)k}(\Lambda; \mathbb{Z}_p) \rightarrow \mathbb{Z}_p$$

is non-degenerate. Hence, there exists $\alpha \in H_{2(p-1)k}(\Lambda; \mathbb{Z}_p)$ such that $0 \neq \langle q_k(\text{ind } L), \alpha \rangle \in \mathbb{Z}_p$. Now we let $\eta \in H^{n-2(p-1)k}(\Lambda; \mathbb{Z}_p)$ be the Poincaré dual of α with respect to a fixed $\mathbb{Z}_p$ -orientation of Λ . According to [6, Corollary VI.8.4], there is a commutative diagram

$$\begin{array}{ccccc} & & \check{H}^{n-2(p-1)k}(\Lambda; \mathbb{Z}_p) & \xrightarrow{i^*} & \check{H}^{n-2(p-1)k}(B(f); \mathbb{Z}_p), \\ & & \uparrow & & \uparrow \\ H_{2(p-1)k}(\Lambda \setminus B(f); \mathbb{Z}_p) & \xrightarrow{j_*} & H_{2(p-1)k}(\Lambda; \mathbb{Z}_p) & \xrightarrow{\pi_*} & H_{2(p-1)k}(\Lambda, \Lambda \setminus B(f); \mathbb{Z}_p) \end{array} \quad (2.3)$$

where the vertical arrows are isomorphisms given by Poincaré–Lefschetz duality and the lower horizontal sequence is part of the long exact homology sequence of the pair $(\Lambda, \Lambda \setminus B(f))$. Because of the commutativity, the class $i^*\eta$ is dual to $\pi_*\alpha$ and we now assume by contradiction the triviality of the latter one.

Then, by exactness, there exists $\beta \in H_{2(p-1)k}(\Lambda \setminus B(f); \mathbb{Z}_p)$ such that $\alpha = j_*\beta$. Moreover, since homology is compactly supported (cf. [11, §20.4]), there exist a compact connected CW-complex P and a map $g: P \rightarrow \Lambda \setminus B(f)$ such that $\beta = g_*\gamma$ for some $\gamma \in H_{2(p-1)k}(P; \mathbb{Z}_p)$.

Let us recall that by assumption there is some $\lambda_0 \in \Lambda$ such that L_{λ_0} is invertible. As $\lambda_0 \notin B(f)$ by the implicit function theorem and $\Lambda \setminus B(f)$ is connected, we can deform g such that λ_0 belongs to its image. Clearly, this does not affect the property that $\beta = g_*\gamma$, and so we can assume without loss of generality that $\lambda_0 \in \text{im } g$. We now set $\bar{g} = j \circ g$ and consider

$$\bar{f}: P \times X \rightarrow Y, \quad \bar{f}(p, u) = f(\bar{g}(p), u).$$

Then $\bar{f}(p, 0) = 0$ for all $p \in P$ and the linearisation of $\bar{f}_p$ at $0 \in X$ is $\bar{L}_p = L_{\bar{g}(p)}$. Moreover, as λ_0 is in the image of g , there is some $p_0 \in P$ such that $\bar{L}_{p_0}$ is invertible. Clearly, $\bar{g}$ sends bifurcation points of $\bar{f}$ to bifurcation points of f , and as $\bar{g}(P) \cap B(f) = \emptyset$, we see that the family $\bar{f}$ has no bifurcation points. Consequently, $J(\text{ind } \bar{L}) = 0 \in J(P)$, by Theorem 1.1, showing that $q_k(\text{ind } \bar{L}) = 0 \in H^{2(p-1)k}(P; \mathbb{Z}_p)$. From (2.1) and (2.2), we obtain

$$\begin{aligned} 0 &= \langle q_k(\text{ind } \bar{L}), \gamma \rangle \\ &= \langle q_k(\bar{g}^*(\text{ind } L), \gamma) \rangle \\ &= \langle \bar{g}^* q_k(\text{ind } L), \gamma \rangle \\ &= \langle g^* j^* q_k(\text{ind } L), \gamma \rangle \\ &= \langle q_k(\text{ind } L), j_* g_* \gamma \rangle \\ &= \langle q_k(\text{ind } L), j_* \beta \rangle \\ &= \langle q_k(\text{ind } L), \alpha \rangle, \end{aligned}$$

which is a contradiction to the choice of α .

Consequently, $\pi_*\alpha$ is non-trivial, and so $i^*\eta \in \check{H}^{n-2(p-1)k}(B(f); \mathbb{Z}_p)$ is non-trivial, as well. Therefore, $\dim B(f) \geq n - 2(p-1)k$ and, moreover, $B(f)$ is not contractible to a point as $n - 2(p-1)k \geq 1$. $\square$

Remark 2.2. Let us point out that the estimate of the dimension of $B(f)$ in Theorem 2.1 (ii) was obtained by Bartsch in [5] for semilinear Fredholm maps f by different methods. His argument is also based on Poincaré–Lefschetz duality and diagram (2.3) for $p = 2$. However, instead of the J -homomorphism and Theorem 1.1, he used the codegree as a numerical invariant that measures the non-triviality of $\text{ind } L$ in $\widehat{KO}(\Lambda)$.

3 An example

In this section we denote by $G_n(\mathbb{R}^{2n})$ the Grassmannian manifold of all n -dimensional subspaces of $\mathbb{R}^{2n}$ (cf. [12, §5]). Let us recall that $G_n(\mathbb{R}^{2n})$ is compact and that for every $n \in \mathbb{N}$ there is a canonical vector bundle $\gamma^n(\mathbb{R}^{2n})$ over $G_n(\mathbb{R}^{2n})$ which is called the *tautological bundle* and which has as total space

$$\{(V, u) \in G_n(\mathbb{R}^{2n}) \times \mathbb{R}^{2n} : u \in V\}.$$

In what follows we denote by $I = [0, 1]$ the compact unit interval, and we let Λ be a compact manifold of dimension $m \geq 2$. We denote by $H^1(I, \mathbb{R}^n)$ the space of all absolutely continuous functions having a square integrable derivative with respect to the usual Sobolev norm

$$\|u\|_{H^1}^2 := \int_0^1 \langle u, u \rangle dt + \int_0^1 \langle u', u' \rangle dt,$$

and we let

$$\varphi : \Lambda \times I \times \mathbb{R}^n \rightarrow \mathbb{R}^n$$

be a continuous function such that each $\varphi_\lambda := \varphi(\lambda, \cdot, \cdot) : I \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ is smooth, all its derivatives depend continuously on λ , and $\varphi(\lambda, t, 0) = 0$ for all $(\lambda, t) \in \Lambda \times I$. For a continuous map $b : \Lambda \rightarrow G_n(\mathbb{R}^{2n})$, we consider the family of differential equations

$$\begin{cases} u'(t) = \varphi_\lambda(t, u(t)), & t \in [0, 1], \\ (u(0), u(1)) \in b(\lambda), \end{cases} \quad (3.1)$$

and we note that $u \equiv 0$ is a solution for all $\lambda \in \Lambda$.

Definition 3.1. A parameter value $\lambda_0 \in \Lambda$ is called a bifurcation point of (3.1) if in every neighbourhood of $(\lambda_0, 0) \in \Lambda \times H^1(I, \mathbb{R}^n)$ there is (λ, u) such that $u \neq 0$ is a solution of (3.1).

In what follows we denote by $B \subset \Lambda$ the set of all bifurcation points. The linearisation of (3.1) at 0 is

$$\begin{cases} u'(t) = (D_0 \varphi_\lambda)(t, \cdot)u(t), & t \in [0, 1], \\ (u(0), u(1)) \in b(\lambda), \end{cases} \quad (3.2)$$

where $D_0 \varphi_\lambda(t, \cdot) : \mathbb{R}^n \rightarrow \mathbb{R}^n$ denotes the derivative of $\varphi_\lambda(t, u)$ with respect to u at $0 \in \mathbb{R}^n$.

We can now state our main theorem of this section.

Theorem 3.2. *If there is some $\lambda_0 \in \Lambda$ such that (3.2) has only the trivial solution and*

$$0 \neq b^*(w_k(\gamma^n(\mathbb{R}^{2n}))) \in H^k(\Lambda; \mathbb{Z}_2)$$

for some $1 \leq k \leq m - 1$, then the dimension of B is at least $m - k$ and B is not a contractible topological space.

Let us note that $G_n(\mathbb{R}^{2n})$ is itself a compact manifold, which is orientable as Grassmannians $G_n(\mathbb{R}^l)$ are orientable if and only if l is even (cf. [3]). We obtain the following corollary for $\Lambda = G_n(\mathbb{R}^{2n})$ and b the identity.

Corollary 3.3. *If $n \geq 2$ and there is some $V_0 \in G_n(\mathbb{R}^{2n})$ such that (3.2) has only the trivial solution, then B is a non-contractible space which has at most codimension 1 in $G_n(\mathbb{R}^{2n})$.*

Let us point out that the restriction which we impose on the dimension n in Corollary 3.3 is satisfied for all equations (3.1) that are first-order reductions of second-order scalar equations.

In the remainder of this section we will be concerned with the proofs of Theorem 3.2 and Corollary 3.3. We set

$$\mathcal{H}_\lambda := \{u \in H^1(I, \mathbb{R}^n) : (u(0), u(1)) \in b(\lambda)\},$$

and we note that each one of the following maps is C^1 :

$$f_\lambda : \mathcal{H}_\lambda \rightarrow L^2(I, \mathbb{R}^n), \quad f_\lambda(u) = u' - \varphi_\lambda(\cdot, u).$$

Moreover,

$$(D_v f_\lambda)u = u' - (D_v \varphi_\lambda)u,$$

and we leave it to the reader to check that each $D_v f_\lambda: \mathcal{H}_\lambda \rightarrow L^2(I, \mathbb{R}^n)$ is a Fredholm operator of index 0.

Note that we cannot apply Theorem 2.1 to the operators f_λ as they are not defined on a single Banach space. To overcome this technical issue, we need the following lemma.

Lemma 3.4. *The set*

$$\mathcal{H} := \{(\lambda, u) \in \Lambda \times H^1(I, \mathbb{R}^n) : u \in \mathcal{H}_\lambda\}$$

is a Hilbert subbundle of the product bundle $\Lambda \times H^1(I, \mathbb{R}^n)$.

Proof. Let $\tilde{P}: G_n(\mathbb{R}^{2n}) \times \mathbb{R}^{2n} \rightarrow \mathbb{R}^{2n}$ be a family of projections in $\mathbb{R}^{2n}$ such that $\text{im}(\tilde{P}_V)$ is the fibre of $\gamma^n(\mathbb{R}^{2n})$ over V for $V \in G_n(\mathbb{R}^{2n})$. We consider the family

$$P: \Lambda \times H^1(I, \mathbb{R}^n) \rightarrow H^1(I, \mathbb{R}^n)$$

defined by

$$(P_\lambda u)(t) = u(t) - (1-t)P_1(I_{\mathbb{R}^{2n}} - \tilde{P}_{b(\lambda)})(u(0), u(1)) - tP_2(I_{\mathbb{R}^{2n}} - \tilde{P}_{b(\lambda)})(u(0), u(1)),$$

where $P_1, P_2: \mathbb{R}^{2n} \rightarrow \mathbb{R}^n$ are the projections on the first and last n components in $\mathbb{R}^{2n}$, respectively, and $I_{\mathbb{R}^{2n}}$ is the identity in $\mathbb{R}^{2n}$. The reader can easily check that $P_\lambda^2 = P_\lambda$ and $\text{im}(P_\lambda) = \mathcal{H}_\lambda$, i.e., P_λ is a projection in $H^1(I, \mathbb{R}^n)$ onto $\mathcal{H}_\lambda$. Hence,

$$\{(\lambda, u) \in \Lambda \times H^1(I, \mathbb{R}^{2n}) : u \in \text{im}(P_\lambda)\} = \mathcal{H}$$

is a subbundle of $\Lambda \times H^1(I, \mathbb{R}^{2n})$, by [10, §III.3]. □

As every Hilbert bundle over a finite CW-complex is trivial, by Kuiper's Theorem and [20, p. 54], there exists a bundle isomorphism $\psi: \Lambda \times H \rightarrow \mathcal{H}$ for some separable Hilbert space H . Moreover, by [10, Theorem VII.3.1] we can assume without loss of generality that each $\psi_\lambda: H \rightarrow \mathcal{H}_\lambda$ is orthogonal and so an isometry. Composing ψ and f , we obtain a family of C^1 maps

$$\tilde{f} := f \circ \psi: \Lambda \times H \rightarrow L^2(I, \mathbb{R}^n)$$

such that $f(\lambda, 0) = 0$ for all $\lambda \in \Lambda$, and each $\tilde{f}_\lambda: H \rightarrow L^2(I, \mathbb{R}^n)$ is Fredholm of index 0. Consequently, we can now apply Theorem 2.1 to the family $\tilde{f}$, and so we need to compute $\text{ind}(\tilde{L}) \in \widetilde{KO}(\Lambda)$, where $\tilde{L}_\lambda = D_0 \tilde{f}_\lambda$, $\lambda \in \Lambda$. By the chain rule, we get

$$\tilde{L}_\lambda = (D_{\psi_\lambda(0)} f_\lambda)(D_0 \psi_\lambda) = (D_0 f_\lambda) \psi_\lambda.$$

Consequently, if we introduce two Hilbert bundle morphisms

$$M: \Lambda \times H \rightarrow \mathcal{H}, \quad M_\lambda u = \psi_\lambda u,$$

and

$$L: \mathcal{H} \rightarrow \Lambda \times L^2(I, \mathbb{R}^n), \quad L_\lambda u = (D_0 f_\lambda)u = u' - (D_0 \varphi_\lambda)u,$$

then $\tilde{L}_\lambda = L_\lambda M_\lambda$, $\lambda \in \Lambda$.

The construction of the index bundle, which we recalled in Section 2.1, generalises almost verbatim to Fredholm morphisms of Banach bundles (cf. [21]). Accordingly, if $\mathcal{X}, \mathcal{Y}$ are Banach bundles over a compact base Λ , and $A: \mathcal{X} \rightarrow \mathcal{Y}$ is a bundle morphism such that every $A_\lambda: \mathcal{X}_\lambda \rightarrow \mathcal{Y}_\lambda$ is Fredholm, then

$$\text{ind}(A) = [E(A, \mathcal{V})] - [\mathcal{V}] \in \widetilde{KO}(\Lambda),$$

where $\mathcal{V} \subset \mathcal{Y}$ is a finite dimensional subbundle such that

$$\text{im}(A_\lambda) + \mathcal{V}_\lambda = \mathcal{Y}_\lambda, \quad \lambda \in \Lambda.$$

As the index bundle is additive under compositions of bundle morphisms, and it is trivial for bundle isomorphisms, we obtain

$$\text{ind}(\tilde{L}) = \text{ind}(LM) = \text{ind}(L) + \text{ind}(M) = \text{ind}(L) \in \widetilde{KO}(\Lambda). \quad (3.3)$$

Let us now consider the Fredholm morphism

$$\hat{L}: \mathcal{H} \rightarrow \Lambda \times L^2(I, \mathbb{R}^n), \quad \hat{L}_\lambda u = u'.$$

Then $L_\lambda - \hat{L}_\lambda: \mathcal{H}_\lambda \rightarrow L^2(I, \mathbb{R}^n)$ is compact for all $\lambda \in \Lambda$ and so, by [21, Corollary 7],

$$\text{ind}(L) = \text{ind}(\hat{L}) \in \widetilde{KO}(\Lambda) \quad (3.4)$$

Our next aim is to compute $\text{ind}(\hat{L})$. Let us denote by Y_1 the n -dimensional subspace of constant functions in $L^2(I, \mathbb{R}^n)$ and

$$Y_2 := \left\{ u \in L^2(I, \mathbb{R}^n) : \int_0^1 u(t) dt = 0 \right\}.$$

Clearly, $Y_1 \cap Y_2 = \{0\}$ and, as every function $u \in L^2(I, \mathbb{R}^n)$ can be written as

$$u(t) = \left(u(t) - \int_0^1 u(s) ds \right) + \int_0^1 u(s) ds,$$

we see that $L^2(I, \mathbb{R}^n) = Y_1 \oplus Y_2$. If $v \in Y_2$, then

$$u(t) = \int_0^t v(s) ds, \quad t \in I,$$

is an element of $\mathcal{H}_\lambda$ which is mapped to v . Consequently, $\text{im}(\hat{L}_\lambda) + Y_1 = L^2(I, \mathbb{R}^n)$ for all $\lambda \in \Lambda$, and we get that

$$\begin{aligned} E(\hat{L}, Y_1) &= \{(\lambda, u) \in \Lambda \times H^1(I, \mathbb{R}^n) : u \in \mathcal{H}_\lambda, \hat{L}_\lambda u \in Y_1\} \\ &= \{(\lambda, u) \in \Lambda \times H^1(I, \mathbb{R}^n) : u(t) = (1-t)a + tb, (a, b) \in b(\lambda)\} \\ &\cong \{(\lambda, a, b) \in \Lambda \times \mathbb{R}^{2n} : (a, b) \in b(\lambda)\}. \end{aligned}$$

As this vector bundle is isomorphic to the pullback $b^*(\gamma^n(\mathbb{R}^{2n}))$ of the tautological bundle $\gamma^n(\mathbb{R}^{2n})$ over $G_n(\mathbb{R}^{2n})$ by b , we finally obtain, by (3.3) and (3.4),

$$\text{ind}(\tilde{L}) = \text{ind}(L) = \text{ind}(\hat{L}) = [E(\hat{L}, Y_1)] - [Y_1] = [b^*(\gamma^n(\mathbb{R}^{2n}))] - [\Theta^n] \in \widetilde{KO}(\Lambda),$$

where Θ^n denotes the product bundle with fibre $\mathbb{R}^n$ over Λ . Hence,

$$w_k(\text{ind}(\tilde{L})) = w_k(b^*(\gamma^n(\mathbb{R}^{2n}))) = b^*w_k(\gamma^n(\mathbb{R}^{2n})) \neq 0 \in H^k(\Lambda; \mathbb{Z}_2),$$

by the assumption of Theorem 3.2. Moreover, as (3.2) has only the trivial solution for $\lambda = \lambda_0$, we see that the operator $L_{\lambda_0}: \mathcal{H}_{\lambda_0} \rightarrow L^2(I, \mathbb{R}^n)$ is injective and so invertible as it is Fredholm of index 0. Since $\psi_{\lambda_0}: H \rightarrow \mathcal{H}_{\lambda_0}$ is invertible, we conclude that $\tilde{L}_{\lambda_0}$ is invertible as well. Consequently, from Theorem 2.1, we obtain that the family of C^1 -Fredholm maps $\tilde{f}$ has a bifurcation point $\lambda^* \in \Lambda$, i.e., there is a sequence $\{(\lambda_n, u_n)\}_{n \in \mathbb{N}} \subset \Lambda \times H$ converging to $(\lambda^*, 0)$ such that $u_n \neq 0$ and $\tilde{f}_{\lambda_n}(u_n) = f_{\lambda_n}(\psi_{\lambda_n} u_n) = 0$ for all $n \in \mathbb{N}$. We now set $v_n := \psi_{\lambda_n} u_n \neq 0 \in \mathcal{H}_{\lambda_n}$ for $n \in \mathbb{N}$. As $\psi_{\lambda_n}: H \rightarrow \mathcal{H}_{\lambda_n}$ is an isometry, it follows that $\|v_n\|_{H^1} = \|u_n\|_H \rightarrow 0$ as $n \rightarrow \infty$, where $\|\cdot\|_H$ is the norm of H . Since $v_n \neq 0$ is a solution of (3.1) by the definition of f , we obtain that λ^* is a bifurcation point of (3.1) in the sense of Definition 3.1, and so Theorem 3.2 is proved.

Let us now focus on the proof of Corollary 3.3. By Theorem 2.1 we only need to recall the well-known fact that $0 \neq w_1(\gamma^n(\mathbb{R}^{2n})) \in H^1(G_n(\mathbb{R}^{2n}); \mathbb{Z}_2)$. Indeed, if we consider the tautological bundle $\gamma^n(\mathbb{R}^\infty)$ over the infinite Grassmannian $G_n(\mathbb{R}^\infty)$, then by [12, Theorem 5.6], for every n -dimensional vector bundle ξ , there is a bundle map $g: \xi \rightarrow \gamma^n(\mathbb{R}^\infty)$. Hence, $w_1(\xi) = \bar{g}^* w_1(\gamma^n(\mathbb{R}^\infty))$, where $\bar{g}$ is the map between the base space of ξ and $G_n(\mathbb{R}^\infty)$ that is induced by g . Hence, if $w_1(\gamma^n(\mathbb{R}^\infty)) = 0$, then $w_1(\xi)$ would be trivial for every n -dimensional bundle ξ . However, this is wrong as if we set $\xi = \gamma^1(\mathbb{R}^2) \oplus \Theta^{n-1}$, where Θ^{n-1} is the product bundle with fibre $\mathbb{R}^{n-1}$ over $G_1(\mathbb{R}^2)$, then

$$w_1(\xi) = w_1(\gamma^1(\mathbb{R}^2)) + w_1(\Theta^{n-1}) = w_1(\gamma^1(\mathbb{R}^2)),$$

and the latter class is non-trivial as $\gamma^1(\mathbb{R}^2)$ is isomorphic to the Möbius bundle over S^1 , which is non-orientable and so has a non-trivial first Stiefel–Whitney class.

If we now let $\iota: G_n(\mathbb{R}^{2n}) \hookrightarrow G_n(\mathbb{R}^\infty)$ be the canonical inclusion map, then $\iota^*(\gamma^n(\mathbb{R}^\infty))$ is isomorphic to $\gamma^n(\mathbb{R}^{2n})$ and we obtain

$$\iota^*w_1(\gamma^n(\mathbb{R}^\infty)) = w_1(\iota^*(\gamma^n(\mathbb{R}^\infty))) = w_1(\gamma^n(\mathbb{R}^{2n})) \in H^1(G_n(\mathbb{R}^{2n}); \mathbb{Z}_2).$$

As $\iota^*: H^1(G_n(\mathbb{R}^\infty); \mathbb{Z}_2) \rightarrow H^1(G_n(\mathbb{R}^{2n}); \mathbb{Z}_2)$ is an isomorphism for $n \geq 2$ by [12, Problem 6-B], we conclude that $w_1(\gamma^n(\mathbb{R}^{2n}))$ is non-trivial, and so Corollary 3.3 is proved.

References

- [1] J. C. Alexander, Bifurcation of zeroes of parametrized functions, *J. Funct. Anal.* **29** (1978), 37–53.
- [2] M. F. Atiyah, *K-Theory*, 2nd ed., Adv. Book Class., Addison-Wesley, Redwood City, 1989.
- [3] V. Bartík and J. Korbaš, Stiefel–Whitney characteristic classes and parallelizability of Grassmann manifolds, *Rend. Circ. Mat. Palermo (2) Suppl.* **6** (1984), 19–29.
- [4] T. Bartsch, The role of the J-homomorphism in multiparameter bifurcation theory, *Bull. Sci. Math. (2)* **112** (1988), 177–184.
- [5] T. Bartsch, The global structure of the zero set of a family of semilinear Fredholm maps, *Nonlinear Anal.* **17** (1991), 313–331.
- [6] G. E. Bredon, *Topology and Geometry*, Grad. Texts in Math. 139, Springer, New York, 1993.
- [7] P. M. Fitzpatrick and J. Pejsachowicz, Nonorientability of the index bundle and several-parameter bifurcation, *J. Funct. Anal.* **98** (1991), 42–58.
- [8] W. Hurewicz and H. Wallmann, *Dimension Theory*, Princeton Math. Ser. 4, Princeton University Press, Princeton, 1948.
- [9] J. Ize, Bifurcation theory for Fredholm operators, *Mem. Amer. Math. Soc.* **7** (1976), 1–128.
- [10] S. Lang, *Differential and Riemannian Manifolds*, 3rd ed., Grad. Texts in Math. 160, Springer, New York, 1995.
- [11] J. P. May, *A Concise Course in Algebraic Topology*, 2nd ed., Chicago University Press, Chicago, 1999.
- [12] J. W. Milnor and J. D. Stasheff, *Characteristic Classes*, Princeton University Press, Princeton, 1974.
- [13] J. Pejsachowicz, K-theoretic methods in bifurcation theory, in: *Fixed Point Theory and its Applications* (Berkeley 1986), Contemp. Math. 72, American Mathematical Society, Providence (1988), 193–206.
- [14] J. Pejsachowicz, Index bundle, Leray–Schauder reduction and bifurcation of solutions of nonlinear elliptic boundary value problems, *Topol. Methods Nonlinear Anal.* **18** (2001), 243–267.
- [15] J. Pejsachowicz, Topological invariants of bifurcation, in: *C*-Algebras and Elliptic Theory II*, Trends Math., Birkhäuser, Basel (2008), 239–250.
- [16] J. Pejsachowicz, Bifurcation of Fredholm maps I. The index bundle and bifurcation, *Topol. Methods Nonlinear Anal.* **38** (2011), 115–168.
- [17] J. Pejsachowicz, Bifurcation of Fredholm maps II. The dimension of the set of bifurcation points, *Topol. Methods Nonlinear Anal.* **38** (2011), 291–305.
- [18] J. Pejsachowicz, The index bundle and bifurcation from infinity of solutions of nonlinear elliptic boundary value problems, *J. Fixed Point Theory Appl.* **17** (2015), 43–64.
- [19] M. Starostka and N. Waterstraat, A remark on singular sets of vector bundle morphisms, *Eur. J. Math.* **1** (2015), 154–159.
- [20] N. Steenrod, *The Topology of Fibre Bundles*, Princeton Math. Ser. 14, Princeton University Press, Princeton, 1951.
- [21] N. Waterstraat, The index bundle for Fredholm morphisms, *Rend. Semin. Mat. Univ. Politec. Torino* **69** (2011), 299–315.