Corrigendum

Jindong Xie, Lieqian Chen, Dongqiang Huang, Weiwei Yue, Jingyu Chen*, Chunxiao Liu*

Corrigendum to “A nitric oxide-releasing prodrug promotes apoptosis in human renal carcinoma cells: Involvement of reactive oxygen species”

<https://doi.org/10.1515/chem-2021-0556>
received June 8, 2021; accepted June 10, 2021

apoptosis in human renal carcinoma cells: Involvement of reactive oxygen species. Open Chem. 2021;19(1):635–45. doi: 10.1515/chem-2020-0075.

Corrigendum to: Xie J, Chen L, Huang D, Yue W, Chen J, Liu C. A nitric oxide-releasing prodrug promotes

After publishing the article, the authors noticed that the wrong Figure 4 was used in the manuscript.

*** Corresponding author: Jingyu Chen**, Department of Urology, The First Hospital of Huizhou, No. 20, Sanxin Road, Huizhou, 516000, Guangdong, China, e-mail: hzmiwai@163.com

*** Corresponding author: Chunxiao Liu**, Department of Urology, Zhujiang Hospital, Southern Medical University, No. 253, Industrial Road, Guangzhou, 510282, Guangdong, China, e-mail: liuchx888@163.com

Jindong Xie: Department of Urology, Zhujiang Hospital, Southern Medical University, No. 253, Industrial Road, Guangzhou, 510282, Guangdong, China

Lieqian Chen, Dongqiang Huang, Weiwei Yue: Department of Urology, The First Hospital of Huizhou, No. 20, Sanxin Road, Huizhou, 516000, Guangdong, China

The correct version of Figure 4 is as follows:

Figure 4: Effect of JS-K on ROS, RNS, mitochondrial membrane potential, GSH/GSSG ratio, ATP production, and apoptotic-related proteins in renal carcinoma cells. Treatment of JS-K renal carcinoma cells (0, 1, 2, and 5 μM) for 360 min, and then detection of intracellular maximum ROS (a), RNS (b), superoxide (c), and GSH/GSSG ratio (d) levels. Upon 360 min of JS-K treatment, measurements were made for mitochondrial membrane potential (e) and ATP output (f). Western blot was used to identify concentrations of apoptotic-related proteins such as Bak, Bax, and PARP after cells were treated with JS-K (0, 1, 2, and 5 μM) for 24 h (g). For at least three separate assays, findings are reported as mean ± SD.