

Matthew Gibson and Jeffrey Streets\*

# Deformation classes in generalized Kähler geometry

<https://doi.org/10.1515/coma-2020-0101>

Received May 5, 2020; accepted October 11, 2020

**Abstract:** We describe natural deformation classes of generalized Kähler structures using the Courant symmetry group, which determine natural extensions of the notions of Kähler class and Kähler cone to generalized Kähler geometry. We show that the generalized Kähler-Ricci flow preserves this generalized Kähler cone, and the underlying real Poisson tensor.

**Keywords:** Ricci flow, generalized geometry

**MSC:** 53C55, 53D18, 53E20, 53E30

## 1 Introduction

A rudimentary notion of Kähler geometry is that of the Kähler class: given  $(M^{2n}, \omega, J)$  a Kähler manifold, the Kähler form  $\omega$  is closed, and  $[\omega] \in H_{\mathbb{R}}^{1,1}$  is the associated Kähler class. Fixing the complex structure  $J$ , the space of all Kähler classes defines an open cone in  $H_{\mathbb{R}}^{1,1}$  (the Kähler cone), and the fundamental result of Demailly-Paun [7] gives a characterization of this cone in terms of pairing against complex subvarieties. The space of Kähler metrics within a given Kähler class is an open infinite dimensional cone in  $C^\infty(M)$  using the  $\sqrt{-1}\partial\bar{\partial}$ -lemma. Thus the basic structure of the space of Kähler metrics compatible with a fixed complex structure  $J$  is fairly well understood. If we instead ask for the space of all Kähler pairs  $(g, J)$  on a given smooth manifold  $M$ , the question becomes decidedly more delicate. The global structure can be quite wild, with disconnected components of arbitrarily large dimension (cf. [6]).

Understanding the space of *generalized Kähler* structures on a given manifold  $M$  becomes even more delicate. Originally discovered by Gates-Hull-Rocek [8], a generalized Kähler structure on a smooth manifold  $M$  is a triple  $(g, b, I, J)$  consisting of a Riemannian metric  $g$  compatible with two integrable complex structures  $I, J$ , and a two-form  $b$  further satisfying

$$d_I^c \omega_I = H = H_0 + db = -d_J^c \omega_J, \quad dH = 0.$$

Later, Gualtieri [11] gave a natural description of this geometry using the language of Hitchin's generalized complex structures [13], in particular in terms of a pair of generalized complex structures  $(J_1, J_2)$  satisfying some natural conditions (cf. §2.1). A fundamental question is to understand the degrees of freedom, moduli, and topology of the space of generalized Kähler structures on a given smooth manifold.

Whereas in the Kähler setting we can roughly speaking divide the problem of understanding the space of Kähler metrics into first identifying the space of possible complex structures, and then understanding the space of compatible Kähler metrics, in the generalized Kähler setting such a decomposition is difficult to achieve (cf. [5] for recent progress). Indeed, in many settings, given two complex structures  $I, J$ , there is at most one compatible metric which defines a generalized Kähler structure. Nonetheless, many different classes of deformations of generalized Kähler structure have been constructed. Joyce gave the first examples

**Matthew Gibson:** Rowland Hall, University of California, Irvine, CA 92617, E-mail: gibsonmd@uci.edu

**\*Corresponding Author: Jeffrey Streets:** Rowland Hall, University of California, Irvine, CA 92617, E-mail: jstreets@uci.edu

of nontrivial (i.e. non-Kähler) generalized Kähler structures by deforming away from hyperKähler structures (cf. [1]), specifically using an action of diffeomorphisms which are Hamiltonian with respect to an associated holomorphic symplectic structure. Later Hitchin produced nontrivial generalized Kähler structures on del Pezzo surfaces, with a choice of holomorphic Poisson structure playing a key role [14]. Further developments on Hamiltonian deformations recently appeared in [5, 10, 12]. Also, Goto [9] has extended the stability result of Kodaira-Spencer to the generalized Kähler setting, with the restriction that one of the generalized complex structures be globally defined by a pure spinor.

Our first goal in this work is to give an elementary derivation of a class of deformations which generalizes and unifies different notions of “Kähler class” arising in the different flavors of generalized Kähler geometry. It is well-known that two-forms ( $B$ -fields) can act on generalized complex structures by conjugation, with the integrability condition being preserved if and only if  $B$  is closed. Our deformations exploit a different, and moreover infinitesimal, action of  $B$ -fields. In particular, we will say (cf. Definition 3.1) that a one-parameter family of generalized Kähler structures is a *canonical deformation* if there exists a one parameter family  $K_t \in \Lambda^2$  such that for all times  $t$  where defined, one has

$$\frac{\partial}{\partial t} \mathbf{J}_1 = [\mathbf{J}_1, e^K \mathbf{J}_1], \quad \frac{\partial}{\partial t} \mathbf{J}_2 = [\mathbf{J}_2, e^K \mathbf{J}_2].$$

As we will discuss in §3, it turns out that these deformations are special cases of the Hamiltonian deformation construction appearing in [4, 12].

Equivalence classes of canonical deformations lead to natural definitions of generalized Kähler class and generalized Kähler cone (cf. §3), which is closely related to definitions appearing in [4]. These definitions make nonobvious departures from the classical idea of Kähler class and Kähler cone. The first is the use of infinitesimal deformations as opposed to ‘large’ deformations. Whereas any two metrics in the same Kähler class admit an explicit relationship using the  $\sqrt{-1}\partial\bar{\partial}$ -lemma, we can no longer expect such an explicit relationship in general. For instance, as described above, Joyce’s construction of nontrivial GK structure uses diffeomorphisms which are Hamiltonian with respect to the associated holomorphic symplectic structure, and in general these cannot be described by a single potential function. Instead, as is typical of Hamiltonian diffeomorphisms, we expect to be able to explicitly describe their infinitesimal deformations. Moreover, given that in the Kähler setting, deformations in the Kähler class involve freezing the complex structure and varying the Kähler form, it is natural to imagine that one should deform while fixing either  $\mathbf{J}_1, \mathbf{J}_2$ . Nonetheless through careful consideration of natural variational classes of different flavors of generalized Kähler metrics it emerges that varying  $\mathbf{J}_1$  and  $\mathbf{J}_2$  simultaneously will correctly capture various existing notions of Kähler class in GK geometry.

A fundamental first step in unpacking this definition is to derive the algebraic and differential conditions which are imposed on  $K$  to preserve the compatibility and integrability conditions for the pair  $(\mathbf{J}_1, \mathbf{J}_2)$ . Through careful computations, it turns out that the answer is pleasingly simple:

**Theorem 1.1.** *Given  $M$  a smooth manifold, suppose  $(\mathbf{J}_1, \mathbf{J}_2)$  is a generalized Kähler structure. Suppose  $(\mathbf{J}_1^t, \mathbf{J}_2^t)$  is a one-parameter family of generalized almost complex structures such that*

$$\frac{\partial}{\partial t} \mathbf{J}_1 = [\mathbf{J}_1, e^K \mathbf{J}_1], \quad \frac{\partial}{\partial t} \mathbf{J}_2 = [\mathbf{J}_2, e^K \mathbf{J}_2],$$

*for some one parameter family  $K_t \in \Lambda^2$ . Then  $(\mathbf{J}_1^t, \mathbf{J}_2^t)$  is a one-parameter family of generalized Kähler structures if and only if for all  $t$  one has*

1.  $K_t \in \Lambda_{J_t}^{1,1}$ ,
2.  $dK_t = 0$ ,
3.  $\langle -\mathbf{J}_1^t \mathbf{J}_2^t, \cdot \rangle > 0$ .

*where  $J^t$  is determined via the Gualtieri map, and  $\langle \cdot, \cdot \rangle$  denotes the symmetric neutral inner product on  $T \oplus T^*$  (cf. §3.1).*

In particular, this theorem exhibits that the canonical deformations are, as is true in the Kähler setting, determined infinitesimally by a closed form which is  $(1, 1)$  with respect to  $J$ . We emphasize here that the condition that  $dK = 0$  does *not* follow from the known fact that the conjugation action of  $B$ -fields on generalized complex structures preserves integrability if and only if  $dB = 0$ . For instance, if we consider our infinitesimal action on a single generalized complex structure, the condition to preserve integrability is *strictly weaker* than  $dK = 0$  (cf. Proposition 2.3). It is only in the context of preserving the integrability conditions of generalized Kähler geometry that one derives  $dK = 0$ .

Despite the simplicity of the conditions of Theorem 1.1 and the apparent simplicity of canonical deformations from the point of view of generalized geometry, the deformations induced on the classical bihermitian data  $(g, I, J)$  are delicate. Remarkably, these canonical deformations unify all previously known instances of “Kähler class” in generalized geometry, specifically the classical notion of Kähler class, the modified Kähler classes implicit in Apostolov-Gualtieri ([2] Proposition 5, cf. also [8]) in the commuting GK case, as well as Joyce’s Hamiltonian deformation construction in the nondegenerate case. All of these are thus in turn special cases of the deformations considered in [4, 12]. We state this for emphasis (cf. §3.6 for notation):

**Proposition 1.2.** *The following hold:*

1. Given  $(M^{2n}, g, J)$  a Kähler manifold, and  $u \in C^\infty(M)$  such that  $\omega + \sqrt{-1}\partial\bar{\partial}u > 0$ , the one-parameter family

$$(\omega_I)_t = \omega + t dI du, \quad I_t = J, \quad J_t = J$$

arises as a canonical deformation of generalized Kähler structures for  $0 \leq t \leq 1$  defined by

$$K_t = dJ du.$$

2. Given  $(M^{2n}, g, I, J)$  a generalized Kähler manifold such that  $[I, J] = 0$ , and  $u \in C^\infty(M)$  such that  $\omega_I + \sqrt{-1}(\partial_+ \bar{\partial}_+ - \partial_- \bar{\partial}_-) u > 0$ , the one-parameter family

$$(\omega_I)_t = \omega_I + t \sqrt{-1}(\partial_+ \bar{\partial}_+ - \partial_- \bar{\partial}_-) u, \quad I_t = I, \quad J_t = J$$

arises as a canonical deformation of generalized Kähler structures for  $0 \leq t \leq 1$  defined by

$$K_t = dJ du.$$

3. Let  $(M^{2n}, g, I, J)$  be a generalized Kähler manifold such that the Poisson structure  $\sigma = \frac{1}{2}[I, J]g^{-1}$  is nondegenerate, with  $\Omega = \sigma^{-1}$ . Given  $u_t \in C^\infty(M)$  a family of smooth functions, let  $\phi_t$  denote the one-parameter family of  $\Omega$ -Hamiltonian diffeomorphisms generated by  $u_t$ . Then, for all  $t$  such that  $-\text{Im} \pi_{\Lambda_t^{1,1}} \Omega_J > 0$ , the one-parameter family of generalized Kähler structures determined by

$$\Omega_t = \Omega, \quad I_t = I, \quad J_t = \phi_t^* J$$

defines a canonical deformation of generalized Kähler structures determined by

$$K_t = dJ_t du_t.$$

As a final point to contextualize these deformations, we recall that various interesting deformation classes of generalized Kähler structure have been produced using holomorphic Poisson structures. Given a generalized Kähler structure  $(g, I, J)$ , there is a Poisson tensor

$$\sigma = \frac{1}{2}[I, J]g^{-1}$$

which is the real part of a holomorphic Poisson tensor with respect to both  $I$  and  $J$ . By choosing an appropriate deformation of  $\sigma$ , Hitchin [14] produced deformations of Kähler metrics on del Pezzo surfaces to strictly generalized Kähler structures. Also the deformation theory of Goto [9] changes this underlying Poisson tensor. As it turns out our deformations fix  $\sigma$  and  $I$ , so occur against a fixed background of a holomorphic Poisson structure. This feature is key to the deformations of Joyce described above, and as discussed above also occurs in special cases of the classes of Hamiltonian deformations studied in [5, 10, 12].

**Corollary 1.3.** *Given  $M$  a smooth manifold, suppose  $(\mathbf{J}_1^t, \mathbf{J}_2^t)$  is a canonical deformation of generalized Kähler structures. Then for all  $t$ ,*

$$I^t \equiv I^0, \quad \sigma^t \equiv \sigma^0.$$

As an application, we are able to express the generalized Kähler-Ricci flow in a simple way using canonical deformations. The equation is an extension of Kähler-Ricci flow to the setting of generalized Kähler geometry, introduced by the second author and Tian [18]. Recently this flow has been used to study the global topology of the (nonlinear) space of generalized Kähler structures in certain settings [3, 16]. To describe this flow, fix  $(g, I, J)$  a generalized Kähler structure. Associated to the Hermitian structure  $(g, I)$  is the Bismut connection

$$\nabla^I = D + \frac{1}{2}Hg^{-1},$$

where  $H = d_I^c \omega_I$ , and  $D$  denotes the Levi-Civita connection. This is a Hermitian connection, and if  $\Omega_I$  denotes its curvature, we obtain a representative of the first Chern class via contraction, called the Bismut-Ricci tensor:

$$\rho_I = \frac{1}{2} \text{tr} \Omega_I I.$$

From the Bianchi identity we know that  $d\rho_I = 0$ , but it is not in general true that  $\rho_I \in \Lambda_I^{1,1}$ , and we will let  $\rho_I^{1,1}$  denote its  $(1, 1)$  projection. Furthermore, associated to  $(g, I)$  we obtain the *I-Lee form*, defined by

$$\theta_I(X) = d^* \omega_I(IX).$$

Similarly we obtain the Lee form  $\theta_J$  associated to  $(g, J)$ . We can now express the generalized Kähler-Ricci flow in the *I*-fixed gauge as

$$\frac{\partial}{\partial t} \omega_I = -\rho_I^{1,1}, \quad \frac{\partial}{\partial t} J = L_{\frac{1}{2}(\theta_J^\# - \theta_I^\#)} J. \tag{1.1}$$

The evolution of the complex structure  $J$  is derived in [18], arising from delicate gauge manipulations and curvature identities. On the other hand it has been shown in several special cases (cf. §4 below) that the generalized Kähler-Ricci is driven entirely by  $\rho_I$ . Using our description of canonical deformations, and a further subtle curvature identity for generalized Kähler manifolds (Proposition 4.2), we confirm that this is true in full generality, and give a very simple description of generalized Kähler-Ricci flow in terms of the associated generalized complex structures.

**Theorem 1.4.** *Let  $(M^{2n}, g_t, I, J_t)$  be a solution of generalized Kähler-Ricci flow in the *I*-fixed gauge. The one-parameter family of pairs of associated generalized complex structures  $(\mathbf{J}_1^t, \mathbf{J}_2^t)$  evolve by*

$$\frac{\partial}{\partial t} \mathbf{J}_1 = [\mathbf{J}_1, e^{-\rho_I} \mathbf{J}_1], \quad \frac{\partial}{\partial t} \mathbf{J}_2 = [\mathbf{J}_2, e^{-\rho_I} \mathbf{J}_2]. \tag{1.2}$$

*In other words, the generalized Kähler-Ricci flow is the canonical deformation driven by the negative of the *I*-Bismut-Ricci tensor.*

Immediately following from Theorem 1.4 and Corollary 1.3 is the fact that generalized Kähler-Ricci flow preserves the underlying real Poisson tensor  $\sigma$ , and moreover preserves the generalized Kähler cone associated to the initial data.

**Corollary 1.5.** *Let  $(M^{2n}, g_t, I, J_t)$  be a solution of generalized Kähler-Ricci flow in the *I*-fixed gauge. The associated one-parameter families of generalized complex structures  $(\mathbf{J}_1, \mathbf{J}_2)$  lies in the generalized Kähler cone associated to the initial data. In particular, for all  $t$  such that the flow is defined,*

$$\sigma^t \equiv \sigma^0,$$

*in other words, the real Poisson tensor  $\sigma$  is fixed along the flow.*

## 2 Formal deformations of generalized complex structure

### 2.1 Background

Given  $M$  a smooth manifold, the generalized tangent bundle is given by  $T \oplus T^*$ . This bundle comes equipped with a family of natural brackets determined by a closed three-form  $H$ . In particular, given  $H \in \Lambda^3 T^*$ ,  $dH = 0$ , define the twisted Courant bracket  $[\cdot, \cdot]$  for sections of  $T \oplus T^*$  via

$$[X + \xi, Y + \eta] = [X, Y] + L_X \eta - L_Y \xi + \frac{1}{2} d(\xi(Y) - \eta(X)) + i_Y i_X H. \quad (2.1)$$

A generalized complex structure  $\mathbf{J}$  is then an almost complex structure on  $T \oplus T^*$ , whose  $\sqrt{-1}$ -eigenbundle, denoted by  $L$ , is integrable with respect to the twisted Courant bracket. This condition is naturally captured by a corresponding version of the Nijenhuis tensor, where for a given almost complex structure on  $T \oplus T^*$  we associate the natural projection maps  $\pi_{0,1}$ ,  $\pi_{1,0}$  and then for  $\vec{x}, \vec{y} \in T \oplus T^*$  we have

$$N_{\mathbf{J}}(\vec{x}, \vec{y}) = \pi_{0,1}[\pi_{1,0}(\vec{x}), \pi_{1,0}(\vec{y})]. \quad (2.2)$$

Direct computations show that this is tensorial, and vanishes if and only if the associated almost generalized complex structure is integrable. See [11] for further discussion.

### 2.2 Variations of generalized complex structure

To begin we define an action of  $B$ -fields on generalized complex structures.

**Definition 2.1.** Given a smooth manifold  $M$  and  $K \in \Lambda^2$ , define

$$\begin{aligned} \Phi_K &: \text{End}(T \oplus T^*) \rightarrow \text{End}(T \oplus T^*), \\ \Phi_K(\mathbf{J}) &= [\mathbf{J}, e^K \mathbf{J}], \end{aligned}$$

where

$$e^K = \begin{pmatrix} 1 & 0 \\ K & 1 \end{pmatrix} \in \text{End}(T \oplus T^*).$$

For a given  $\mathbf{J}$ , we intend to use  $\Phi_K(\mathbf{J})$  as a tangent vector to a one-parameter variation of  $\mathbf{J}$  through generalized complex structures. We first note that variations of this kind will indeed preserve the space of generalized almost complex structures.

**Lemma 2.2.** *Let  $\mathbf{J}_t$  be a one-parameter family of endomorphisms of  $T \oplus T^*$  such that  $\mathbf{J}_0$  is an almost generalized complex structure and*

$$\frac{\partial}{\partial t} \mathbf{J}_t = [\mathbf{J}_t, e^{K_t} \mathbf{J}_t].$$

*Then  $\mathbf{J}_t^2 = -1$  for each  $t$ , i.e.  $\mathbf{J}_t$  is a family of generalized almost complex structures.*

*Proof.* Differentiating the expression  $\mathbf{J}_t^2$  yields

$$\frac{\partial}{\partial t} \mathbf{J}_t^2 = \left( \frac{\partial}{\partial t} \mathbf{J} \right) \mathbf{J} + \mathbf{J} \left( \frac{\partial}{\partial t} \mathbf{J} \right) = \left( \mathbf{J} e^K \mathbf{J} + e^K \right) \mathbf{J} + \mathbf{J} \left( \mathbf{J} e^K \mathbf{J} + e^K \right) = 0.$$

Thus  $\frac{\partial}{\partial t} \mathbf{J}_t^2 = 0$ , and since  $\mathbf{J}_0^2 = -1$ , the lemma follows.  $\square$

Next we can characterize the condition for these deformations to preserve integrability of  $\mathbf{J}_t$ .

**Proposition 2.3.** *Let  $\mathbf{J}_t$  be a one-parameter family of generalized almost complex structures such that*

$$\left. \frac{\partial}{\partial t} \mathbf{J}_t \right|_{t=0} = \Phi_K(\mathbf{J}_0).$$

Then for  $\vec{x}, \vec{y} \in T \oplus T^*$ ,

$$\begin{aligned} \left. \frac{\partial}{\partial t} \right|_{t=0} N_{\mathbf{J}_t}(\vec{x}, \vec{y}) &= \sqrt{-1} \pi_{0,1} (dK(\pi_T \pi_{1,0}(\vec{y}), \pi_T \pi_{1,0}(\vec{x}), \cdot)) \\ &\quad - \sqrt{-1} N_{\mathbf{J}_0}(\vec{x}, \vec{y}) + \mathbf{J}_0 e^K N_{\mathbf{J}_0}(\vec{x}, \vec{y}) + N_{\mathbf{J}_0}(e^K \mathbf{J}_0(\vec{x}), \vec{y}) + N_{\mathbf{J}_0}(\vec{x}, e^K \mathbf{J}_0 \vec{y}). \end{aligned}$$

*Proof.* For notational simplicity we set  $\mathbf{J} = \mathbf{J}_0$ . We differentiate the formula (2.2) at  $t = 0$ , using the explicit formulae for the projection maps, to obtain

$$\begin{aligned} &\left. \frac{\partial}{\partial t} (\pi_{0,1}^t [\pi_{1,0}^t(\vec{x}), \pi_{1,0}^t(\vec{y})]) \right. \\ &= \left. \frac{\partial}{\partial t} (\pi_{0,1}^t) [\pi_{1,0}^t(\vec{x}), \pi_{1,0}^t(\vec{y})] + \pi_{0,1}^t \left[ \frac{\partial}{\partial t} \pi_{1,0}^t(\vec{x}), \pi_{1,0}^t(\vec{y}) \right] + \pi_{0,1}^t \left[ \pi_{1,0}^t(\vec{x}), \frac{\partial}{\partial t} \pi_{1,0}^t(\vec{y}) \right] \right. \\ &= \frac{\sqrt{-1}}{2} \{ \Phi_K(\mathbf{J})[\pi_{1,0}(\vec{x}), \pi_{1,0}(\vec{y})] - \pi_{0,1}[\Phi_K(\mathbf{J})(\vec{x}), \pi_{1,0}(\vec{y})] - \pi_{0,1}[\pi_{1,0}(\vec{x}), \Phi_K(\mathbf{J})(\vec{y})] \}. \end{aligned} \quad (2.3)$$

Note that for any  $\vec{z} \in T \oplus T^*$  we may write  $\vec{z} = 2\pi_{1,0}(\vec{z}) + \sqrt{-1}\mathbf{J}(\vec{z})$  which leads to

$$\begin{aligned} \Phi_K(\mathbf{J})(\vec{z}) &= (\mathbf{J}e^K\mathbf{J} + e^K)(\vec{z}) \\ &= \mathbf{J}e^K\mathbf{J}(\vec{z}) + e^K(2\pi_{1,0}(\vec{z}) + \sqrt{-1}\mathbf{J}(\vec{z})) \\ &= 2e^K\pi_{1,0}(\vec{z}) + \sqrt{-1}(e^K\mathbf{J}\vec{z} - \sqrt{-1}\mathbf{J}(e^K\mathbf{J}\vec{z})) \\ &= 2e^K\pi_{1,0}(\vec{z}) + 2\sqrt{-1}\pi_{1,0}(e^K\mathbf{J}(\vec{z})). \end{aligned}$$

This observation allows us to rewrite the final line of (2.3) as

$$\begin{aligned} \dots &= \sqrt{-1}e^K\pi_{1,0}[\pi_{1,0}(\vec{x}), \pi_{1,0}(\vec{y})] - \pi_{1,0}e^K\mathbf{J}[\pi_{1,0}(\vec{x}), \pi_{1,0}(\vec{y})] \\ &\quad - \sqrt{-1}\pi_{0,1}[e^K\pi_{1,0}(\vec{x}) + \sqrt{-1}\pi_{1,0}e^K\mathbf{J}(\vec{x}), \pi_{1,0}(\vec{y})] \\ &\quad - \sqrt{-1}[\pi_{1,0}(\vec{x}), e^K\pi_{1,0}(\vec{y}) + \sqrt{-1}\pi_{1,0}e^K\mathbf{J}(\vec{y})] \\ &= \sqrt{-1}e^K\pi_{1,0}[\pi_{1,0}(\vec{x}), \pi_{1,0}(\vec{y})] - \pi_{1,0}e^K\mathbf{J}[\pi_{1,0}(\vec{x}), \pi_{1,0}(\vec{y})] \\ &\quad - \sqrt{-1}\pi_{0,1}[e^K\pi_{1,0}(\vec{x}), \pi_{1,0}(\vec{y})] - \sqrt{-1}\pi_{0,1}[\pi_{1,0}(\vec{x}), e^K\pi_{1,0}(\vec{y})] \\ &\quad + \pi_{0,1}([\pi_{1,0}e^K\mathbf{J}(\vec{x}), \pi_{1,0}(\vec{y})] + [\pi_{1,0}(\vec{x}), \pi_{1,0}(e^K\mathbf{J}\vec{y})]). \end{aligned}$$

Focusing on the first two terms, letting  $\vec{z} = [\pi_{1,0}(\vec{x}), \pi_{1,0}(\vec{y})]$ ,

$$\begin{aligned} &\sqrt{-1}e^K\pi_{1,0}\vec{z} - \pi_{1,0}e^K\mathbf{J}\vec{z} \\ &= \frac{\sqrt{-1}}{2}e^K(\vec{z} - \sqrt{-1}\mathbf{J}\vec{z}) - \frac{1}{2}(e^K\mathbf{J}\vec{z} - \sqrt{-1}\mathbf{J}e^K\mathbf{J}\vec{z}) \\ &= \frac{\sqrt{-1}}{2}(e^K\vec{z} + \mathbf{J}e^K\mathbf{J}\vec{z}) \\ &= \sqrt{-1}\pi_{0,1}e^K\vec{z} + \frac{1}{2}\mathbf{J}e^K\vec{z} + \frac{\sqrt{-1}}{2}\mathbf{J}e^K\mathbf{J}\vec{z} \\ &= \sqrt{-1}\pi_{0,1}(e^K\vec{z}) + \frac{1}{2}\mathbf{J}e^K(\vec{z} + \sqrt{-1}\mathbf{J}\vec{z}) \\ &= \sqrt{-1}\pi_{0,1}(e^K\vec{z}) + \mathbf{J}e^K\pi_{0,1}(\vec{z}). \end{aligned}$$

Equation (2.3) now simplifies further to

$$\begin{aligned} \dots &= \sqrt{-1}\pi_{0,1} \left( e^K[\pi_{1,0}(\vec{x}), \pi_{1,0}(\vec{y})] - [e^K\pi_{1,0}(\vec{x}), \pi_{1,0}(\vec{y})] - [\pi_{1,0}(\vec{x}), e^K\pi_{1,0}(\vec{y})] \right) \\ &\quad + \mathbf{J}e^K\pi_{0,1}[\pi_{1,0}(\vec{x}), \pi_{1,0}(\vec{y})] + \pi_{0,1} \left( [\pi_{1,0}e^K\mathbf{J}(\vec{x}), \pi_{1,0}(\vec{y})] + [\pi_{1,0}(\vec{x}), \pi_{1,0}(e^K\mathbf{J}\vec{y})] \right) \\ &= \sqrt{-1}\pi_{0,1} \left( e^K[\pi_{1,0}(\vec{x}), \pi_{1,0}(\vec{y})] - [e^K\pi_{1,0}(\vec{x}), \pi_{1,0}(\vec{y})] - [\pi_{1,0}(\vec{x}), e^K\pi_{1,0}(\vec{y})] \right) \\ &\quad + \mathbf{J}e^KN(\vec{x}, \vec{y}) + N(e^K\mathbf{J}(\vec{x}), \vec{y}) + N(\vec{x}, e^K\mathbf{J}\vec{y}). \end{aligned} \quad (2.4)$$

The Courant bracket satisfies  $e^K[X + \xi, Y + \eta] = [e^K(X + \xi), e^K(Y + \eta)] + \iota_{X+\xi} \iota_Y dK$ . Using this together with the fact that the Courant bracket involving a section with no tangent vector component vanishes we see that the first term of (2.4) becomes

$$\begin{aligned} & \sqrt{-1}\pi_{0,1} \left( [e^K\pi_{1,0}(\vec{x}), e^K\pi_{1,0}(\vec{y})] - [e^K\pi_{1,0}(\vec{x}), \pi_{1,0}(\vec{y})] - [\pi_{1,0}(\vec{x}), e^K\pi_{1,0}(\vec{y})] + dK(\pi_T\pi_{1,0}(\vec{y}), \pi_T\pi_{1,0}(\vec{x}), \cdot) \right) \\ &= \sqrt{-1}\pi_{0,1} \left( -[\pi_{1,0}(\vec{x}), \pi_{1,0}(\vec{y})] + dK(\pi_T\pi_{1,0}(\vec{y}), \pi_T\pi_{1,0}(\vec{x}), \cdot) \right) \\ &= -\sqrt{-1}N(\vec{x}, \vec{y}) + \sqrt{-1}\pi_{0,1}(dK(\pi_T\pi_{1,0}(\vec{y}), \pi_T\pi_{1,0}(\vec{x}), \cdot)). \end{aligned}$$

Collecting these computations gives the result.  $\square$

**Corollary 2.4.** *Let  $\mathbf{J}_t$  be a one-parameter family of generalized almost complex structures such that  $\mathbf{J}_0$  is integrable and for all  $t$  one has*

$$\frac{\partial}{\partial t} \mathbf{J} = \Phi_{K_t}(\mathbf{J}),$$

where furthermore for all  $\vec{x}, \vec{y} \in T \oplus T^*$  one has

$$\pi_{0,1}^t dK_t \left( \pi_T\pi_{1,0}^t(\vec{y}), \pi_T\pi_{1,0}^t(\vec{x}), \cdot \right) = 0. \quad (2.5)$$

Then  $\mathbf{J}_t$  are integrable for each  $t$ .

*Proof.* Choosing any Hermitian metric on  $(T \oplus T^*) \otimes \mathbb{C}$ , using Proposition 2.3 and the hypothesis (2.5) one directly derives for all  $t$

$$\frac{\partial}{\partial t} |N_{\mathbf{J}_t}|^2 \leq C(K, \mathbf{J}) |N_{\mathbf{J}_t}|^2.$$

Since  $N_{\mathbf{J}_0} = 0$  the result follows from Gronwall's inequality.  $\square$

**Remark 2.5.** We may also formulate integrability of generalized complex structures in terms of its  $-\sqrt{-1}$ -eigenbundle,  $\bar{L}_t$ , being Courant integrable. Replicating the above arguments with the roles of  $\pi_{1,0}$  and  $\pi_{0,1}$  reversed we obtain that the relevant condition on  $K_t$  is

$$0 = \pi_{1,0}^t dK_t \left( \pi_T\pi_{0,1}^t(\vec{y}), \pi_T\pi_{0,1}^t(\vec{x}), \cdot \right).$$

### 3 Variations of generalized Kähler structure

Having defined certain variations of generalized complex structure, we now extend this to defining variations of generalized Kähler structure. A naive guess would be that we should simply take a variation of one of the underlying generalized complex structures and seek the further integrability conditions. However, for reasons to be illuminated by the examples below, it is much more natural to vary *both* generalized complex structures by a single  $B$ -field as described in §2.2.

#### 3.1 Background

A generalized Kähler structure is a pair of commuting generalized complex structures  $\mathbf{J}_1, \mathbf{J}_2$  such that  $\mathbf{G} = -\mathbf{J}_1\mathbf{J}_2$  defines a generalized metric, i.e.  $\langle \mathbf{G}\cdot, \cdot \rangle$  is a positive definite inner product on  $T \oplus T^*$ , where  $\langle \cdot, \cdot \rangle$  denotes the symmetric neutral inner product on  $T \oplus T^*$ , i.e.

$$\langle X + \xi, Y + \eta \rangle = \frac{1}{2} (\xi(Y) + \eta(X)).$$

A fundamental theorem of Gualtieri ([11] Chapter 6) says that a generalized Kähler structure  $(\mathbf{J}_1, \mathbf{J}_2)$  as defined here corresponds to a bihermitian structure  $(g, I, J, b)$ , with Kähler forms  $\omega_I, \omega_J$ , as described in the introduction. The explicit relationship is given by

$$\mathbf{J}_{1/2} = \frac{1}{2} e^b \begin{pmatrix} I \pm J & -(\omega_I^{-1} \mp \omega_J^{-1}) \\ \omega_I \mp \omega_J & -(I^* \pm J^*) \end{pmatrix} e^{-b}. \quad (3.1)$$

We recall that a generalized Kähler structure induces a fourfold decomposition of the complexified generalized tangent bundle. Specifically, letting  $L_i$  and  $\bar{L}_i$  denote the  $\pm\sqrt{-1}$ -eigenbundles of  $\mathbf{J}_i$  respectively, we have the following decomposition:

$$(T \oplus T^*) \otimes \mathbb{C} = L_1^+ \oplus L_1^- \oplus \bar{L}_1^- \oplus \bar{L}_1^+ := (L_1 \cap L_2) \oplus (L_1 \cap \bar{L}_2) \oplus (\bar{L}_1 \cap L_2) \oplus (\bar{L}_1 \cap \bar{L}_2).$$

### 3.2 Definitions

**Definition 3.1.** A one-parameter family of generalized Kähler structures  $(\mathbf{J}_1^t, \mathbf{J}_2^t)$  is a *canonical family* if for all  $t$ , there exists  $K_t \in \Lambda^2$  such that

$$\frac{\partial}{\partial t} \mathbf{J}_1^t = \Phi_{K_t}(\mathbf{J}_1^t), \quad \frac{\partial}{\partial t} \mathbf{J}_2^t = \Phi_{K_t}(\mathbf{J}_2^t).$$

Given  $\tilde{\mathbf{J}}_1, \tilde{\mathbf{J}}_2$  another generalized Kähler structure, we define an equivalence relation where

$$(\tilde{\mathbf{J}}_1, \tilde{\mathbf{J}}_2) \sim (\mathbf{J}_1, \mathbf{J}_2)$$

if and only if there exists a canonical family  $(\mathbf{J}_1^t, \mathbf{J}_2^t)$ ,  $t \in [0, 1]$ , such that  $(\mathbf{J}_1^0, \mathbf{J}_2^0) = (\mathbf{J}_1, \mathbf{J}_2)$ ,  $(\mathbf{J}_1^1, \mathbf{J}_2^1) = (\tilde{\mathbf{J}}_1, \tilde{\mathbf{J}}_2)$ . Furthermore, the *generalized Kähler cone associated to*  $(\mathbf{J}_1, \mathbf{J}_2)$  is

$$\mathcal{GK}(\mathbf{J}_1, \mathbf{J}_2) = \{(\tilde{\mathbf{J}}_1, \tilde{\mathbf{J}}_2) \text{ generalized Kähler} \mid (\tilde{\mathbf{J}}_1, \tilde{\mathbf{J}}_2) \sim (\mathbf{J}_1, \mathbf{J}_2)\}.$$

Below we will establish that  $K_t$  must be closed to preserve the integrability of the generalized Kähler structure. Given this, it follows that these deformations can be described using the Hamiltonian deformations of [4, 12], as we will make explicit below.

### 3.3 Compatibility Condition

Here we address the condition required for a canonical deformation to preserve the algebraic compatibility condition of generalized Kähler structures. We first prove a formal lemma reducing this to an algebraic condition on  $K$ , then analyze this explicitly using the Gualtieri map.

**Lemma 3.2.** Let  $\mathbf{J}_1^t, \mathbf{J}_2^t$  be one-parameter families of generalized almost complex structures, with  $[\mathbf{J}_1^0, \mathbf{J}_2^0] = 0$ , which satisfy

$$\frac{\partial}{\partial t} \mathbf{J}_1^t = \Phi_{K_t}(\mathbf{J}_1^t), \quad \frac{\partial}{\partial t} \mathbf{J}_2^t = \Phi_{K_t}(\mathbf{J}_2^t).$$

Then  $[\mathbf{J}_1^t, \mathbf{J}_2^t] = 0$  for all  $t$  if and only if

$$[\Phi_{K_t}(\mathbf{J}_1^t), \mathbf{J}_2^t] = [\Phi_{K_t}(\mathbf{J}_2^t), \mathbf{J}_1^t]$$

for all  $t$ .

*Proof.* Differentiating  $[\mathbf{J}_1^t, \mathbf{J}_2^t]$  at any time  $t$  shows

$$\begin{aligned} \frac{\partial}{\partial t} [\mathbf{J}_1^t, \mathbf{J}_2^t] &= \Phi_K(\mathbf{J}_1) \mathbf{J}_2 + \mathbf{J}_1 \Phi_K(\mathbf{J}_2) - \Phi_K(\mathbf{J}_2) \mathbf{J}_1 - \mathbf{J}_2 \Phi_K(\mathbf{J}_1) \\ &= [\Phi_K(\mathbf{J}_1), \mathbf{J}_2] - [\Phi_K(\mathbf{J}_2), \mathbf{J}_1]. \end{aligned}$$

Since  $[\mathbf{J}_1^0, \mathbf{J}_2^0] = 0$ , the result follows.  $\square$



Here we reformulate the compatibility condition of Lemma 3.2 by expanding the necessary equation in terms of the Gualtieri map and analyzing the result, which simplifies dramatically.

**Proposition 3.3.** *Given  $M$  a smooth manifold and  $(\mathbf{J}_1, \mathbf{J}_2)$  a generalized Kähler structure, for  $K \in \Lambda^2$  one has*

$$[\Phi_K(\mathbf{J}_1), \mathbf{J}_2] = [\Phi_K(\mathbf{J}_2), \mathbf{J}_1]$$

if and only if

$$K \in \Lambda_J^{1,1}.$$

*Proof.* Let  $Y_{1/2} = \frac{1}{2} \begin{pmatrix} (I \pm J) & -(\omega_I^{-1} \mp \omega_J^{-1}) \\ (\omega_I \mp \omega_J) & -(I^* \pm J^*) \end{pmatrix}$ , so that (3.1) can be expressed as  $\mathbf{J}_{1/2} = e^b Y_{1/2} e^{-b}$ . Using this notation and the fact that  $e^K$  and  $e^b$  commute, it follows easily that

$$\begin{aligned} [\Phi_K(\mathbf{J}_1), \mathbf{J}_2] &= e^b [\Phi_K(Y_1), Y_2] e^{-b}, \\ [\Phi_K(\mathbf{J}_2), \mathbf{J}_1] &= e^b [\Phi_K(Y_2), Y_1] e^{-b}. \end{aligned}$$

Hence  $[\Phi_K(\mathbf{J}_1), \mathbf{J}_2] = [\Phi_K(\mathbf{J}_2), \mathbf{J}_1]$  reduces to the condition  $[\Phi_K(Y_1), Y_2] = [\Phi_K(Y_2), Y_1]$ . As a first step, we record the simplified forms of  $\Phi_K(Y_{1/2})$  obtained through a direct computation:

$$\Phi_K(Y_{1/2}) = \frac{1}{4} \begin{pmatrix} -(\omega_I^{-1} \mp \omega_J^{-1})K(I \pm J) & (\omega_I^{-1} \mp \omega_J^{-1})K(\omega_I^{-1} \mp \omega_J^{-1}) \\ 4K - (I^* \pm J^*)K(I \pm J) & (I^* \pm J^*)K(\omega_I^{-1} \mp \omega_J^{-1}) \end{pmatrix}.$$

Further tedious computation yields

$$\begin{aligned} [\Phi_K(\mathbf{J}_1), \mathbf{J}_2] &= \frac{1}{2} \begin{pmatrix} (\omega_I^{-1} + \omega_J^{-1})K + g^{-1}K(I + J) & (\omega_I^{-1} - \omega_J^{-1})Kg^{-1} - g^{-1}K(\omega_I^{-1} - \omega_J^{-1}) \\ K(I - J) + (I^* - J^*)K & (I^* + J^*)Kg^{-1} - K(\omega_I^{-1} + \omega_J^{-1}) \end{pmatrix}, \\ [\Phi_K(\mathbf{J}_2), \mathbf{J}_1] &= \frac{1}{2} \begin{pmatrix} (\omega_I^{-1} - \omega_J^{-1})K + g^{-1}K(I - J) & (\omega_I^{-1} + \omega_J^{-1})Kg^{-1} - g^{-1}K(\omega_I^{-1} + \omega_J^{-1}) \\ K(I + J) + (I^* + J^*)K & (I^* - J^*)Kg^{-1} - K(\omega_I^{-1} - \omega_J^{-1}) \end{pmatrix}. \end{aligned}$$

By comparing each entry of the matrices above, we see equality holds if and only if  $KJ = -J^*K$ , as required.  $\square$

### 3.4 Integrability Condition

We next address the integrability condition. Since our deformations should preserve integrability of each generalized complex structure  $\mathbf{J}_i$ , Proposition 2.3 yields two partial integrability conditions which  $K$  must satisfy. We again emphasize that neither of these conditions alone will force  $dK = 0$ , while somewhat surprisingly the combination of the two conditions does.

**Proposition 3.4.** *Given  $M$  a smooth manifold and  $(\mathbf{J}_1, \mathbf{J}_2)$  a generalized Kähler structure, for  $K \in \Lambda^2$  one has*

$$\pi_{0,1}^{\mathbf{J}_i}(t_X t_Y dK) = 0 \quad \text{for all } X, Y \in \pi_T(L_i), \quad i = 1, 2.$$

if and only if

$$dK = 0.$$

*Proof.* The sufficiency of  $dK = 0$  is obvious, we prove it is necessary. Note that for a pure covector  $\xi$ , one has

$$\mathbf{J}_{1/2}\xi = e^b Y_{1/2}\xi = -\frac{1}{2} \begin{pmatrix} (\omega_I^{-1} \mp \omega_J^{-1})\xi \\ b(\omega_I^{-1} \mp \omega_J^{-1})\xi + (I^* \pm J^*)\xi \end{pmatrix} \text{ and so}$$

$$\pi_{0,1}^{1/2}(\xi) = \left( \begin{array}{c} \frac{-\sqrt{-1}}{2}(\omega_I^{-1} \mp \omega_J^{-1})\xi \\ \xi - \frac{\sqrt{-1}}{2} (b(\omega_I^{-1} \mp \omega_J^{-1})\xi + (I^* \pm J^*)\xi) \end{array} \right). \quad (3.2)$$

Fix vectors  $X, Y \in T_I^{1,0}$ , and then choose lifts  $X_+, Y_+$  to  $C_+$ , the  $+1$ -eigenspace of  $\mathbf{G}$ . Using the representation of  $\mathbf{J}_i$  with respect to the  $\pm 1$ -eigenspace decomposition induced by  $\mathbf{G}$  ([11] Proposition 6.12), it follows that  $X_+, Y_+ \in L_1 \cap L_2 = L_1^+$ . Now let  $\xi = i_Y i_X dK$ , and note that Proposition 2.3 applied to both  $\mathbf{J}_1$  and  $\mathbf{J}_2$  implies that

$$\pi_{0,1}^{1/2}(\xi) = 0.$$

Comparing against equation (3.2) we obtain  $(\omega_I^{-1} \mp \omega_J^{-1})(\xi) = 0$ . Therefore  $i_Y i_X dK = 0$ , for all  $X, Y \in T_I^{1,0}$ . Since  $K$  is real it follows that  $dK = 0$ .  $\square$

*Proof of Theorem 1.1.* Fix  $(\mathbf{J}_1, \mathbf{J}_2)$  generalized Kähler and fix  $(\mathbf{J}_1^t, \mathbf{J}_2^t)$  a one-parameter family as in the statement. First let us assume conditions (1), (2), and (3) hold for this family. Since  $dK_t = 0$  for all  $t$ , it follows from Corollary 2.4 that  $(\mathbf{J}_1^t, \mathbf{J}_2^t)$  are integrable generalized complex structures. Furthermore, using that  $K_t \in \Lambda_{J_t}^{1,1}$  for all  $t$ , it follows from Lemma 2.2 and Proposition 3.3 that  $[\mathbf{J}_1^t, \mathbf{J}_2^t] = 0$  for all  $t$ . Since we have assumed the positivity of  $\langle -\mathbf{J}_1 \mathbf{J}_2 \cdot, \cdot \rangle$  in condition (3), it follows that  $(\mathbf{J}_1^t, \mathbf{J}_2^t)$  is generalized Kähler for all  $t$ .

Conversely, suppose  $(\mathbf{J}_1^t, \mathbf{J}_2^t)$  defines a generalized Kähler structure for all  $t$ . Condition (3) then holds by definition, and condition (1) holds by Lemma 2.2 and Proposition 3.3. As the structures  $(\mathbf{J}_1^t, \mathbf{J}_2^t)$  are assumed integrable for all times  $t$ , their Nijenhuis tensors vanish for all  $t$ , and thus it follows from Proposition 2.3 that

$$0 = \pi_{0,1}^{J_i} dK \left( \pi_T \pi_{1,0}^{J_i}(\vec{x}), \pi_T \pi_{1,0}^{J_i}(\vec{y}), \cdot \right)$$

for all  $\vec{x}, \vec{y} \in T \oplus T^*$ , all  $t$ , and  $i = 1, 2$ . It then follows from Proposition 3.4 that  $dK = 0$ , as required.  $\square$

With this characterization of canonical deformations in hand, we can now give the definition of generalized Kähler *classes* which emerges naturally from Theorem 1.1. We again note that these definitions are closely related to notions of equivalence of generalized Kähler structure arising in [4, 12]. Specifically, our canonical families can be described in terms of the deformations in ([12] Proposition 4.6) with  $\beta_1 = \beta_2 = \sqrt{-1} J_0^t K$ . We refer to [4, 5, 12] for more detailed results and the geometric nature of these deformations, and related notions of generalized Kähler cone.

**Definition 3.5.** Let  $M$  be a smooth manifold. An *exact canonical deformation* is a one-parameter family of generalized Kähler structures  $(\mathbf{J}_1^t, \mathbf{J}_2^t)$  such that, for all  $t$ ,

$$\frac{\partial}{\partial t} \mathbf{J}_i^t = \Phi_{da_t} \mathbf{J}_i^t,$$

for some  $a_t \in \Lambda^1$ . Given  $\tilde{\mathbf{J}}_1, \tilde{\mathbf{J}}_2$  another generalized Kähler structure, we define an equivalence relation where

$$(\tilde{\mathbf{J}}_1, \tilde{\mathbf{J}}_2) \sim_{\text{exact}} (\mathbf{J}_1, \mathbf{J}_2)$$

if and only if there exists an exact canonical deformation  $(\mathbf{J}_1^t, \mathbf{J}_2^t)$ ,  $t \in [0, 1]$ , such that  $(\mathbf{J}_1^0, \mathbf{J}_2^0) = (\mathbf{J}_1, \mathbf{J}_2)$ ,  $(\mathbf{J}_1^1, \mathbf{J}_2^1) = (\tilde{\mathbf{J}}_1, \tilde{\mathbf{J}}_2)$ . Furthermore, the *generalized Kähler class* of  $(\mathbf{J}_1, \mathbf{J}_2)$  is

$$[(\mathbf{J}_1, \mathbf{J}_2)] = \{(\tilde{\mathbf{J}}_1, \tilde{\mathbf{J}}_2) \text{ generalized Kähler} \mid (\tilde{\mathbf{J}}_1, \tilde{\mathbf{J}}_2) \sim_{\text{exact}} (\mathbf{J}_1, \mathbf{J}_2)\}.$$

### 3.5 Induced variations

In this section we derive the variation on the associated bihermitian data induced by a canonical deformation through an analysis of the Gualtieri map.

**Proposition 3.6.** *Let  $(\mathbf{J}_1^t, \mathbf{J}_2^t)$  be a canonical family, and let  $(g_t, b_t, I_t, J_t)$  denote the corresponding one-parameter family of bihermitian data. Then*

$$\begin{aligned}\dot{g} &= -\frac{1}{2}[K, I], & \dot{b} &= -\frac{1}{2}\{K, I\}, \\ \dot{\omega}_I &= -\frac{1}{2}[K, I]I, & \dot{\omega}_J &= -\frac{1}{2}\{K, IJ\}, \\ \dot{I} &= 0, & \dot{J} &= \frac{1}{2}[I, J]g^{-1}K.\end{aligned}$$

*Proof.* We use the notation and computations of Proposition 3.3. In particular, we recall that

$$e^{-b}\mathbf{J}_{1/2}e^b = \frac{1}{4} \begin{pmatrix} -(\omega_I^{-1} \mp \omega_J^{-1})K(I \pm J) & (\omega_I^{-1} \mp \omega_J^{-1})K(\omega_I^{-1} \mp \omega_J^{-1}) \\ 4K - (I^* \pm J^*)K(I \pm J) & (I^* \pm J^*)K(\omega_I^{-1} \mp \omega_J^{-1}) \end{pmatrix}.$$

On the other hand, writing expression (3.1) as  $\mathbf{J}_{1/2} = \frac{1}{2}e^b Y_{1/2} e^{-b}$  and differentiating, using the fact that  $e^b \dot{e}^b = \dot{e}^b = \dot{e}^b e^b$ , yields

$$\begin{aligned}e^{-b}\dot{\mathbf{J}}_{1/2}e^b &= \frac{1}{2}e^{-b} \left( \dot{e}^b Y_{1/2} e^{-b} + e^b \dot{Y}_{1/2} e^{-b} - e^b Y_{1/2} \dot{e}^b \right) \\ &= \frac{1}{2}[\dot{e}^b, Y_{1/2}] + \frac{1}{2}\dot{Y}_{1/2} \\ &= \frac{1}{2} \begin{pmatrix} (\omega_I^{-1} \mp \omega_J^{-1})\dot{b} + (\dot{I} \pm \dot{J}) & -(\dot{\omega}_I^{-1} \mp \dot{\omega}_J^{-1}) \\ \{\dot{b}, I \pm J\} + (\dot{\omega}_I \mp \dot{\omega}_J) & -\dot{b}(\omega_I^{-1} \mp \omega_J^{-1}) - (I^* \pm J^*) \end{pmatrix}.\end{aligned}\tag{3.3}$$

Then equating the appropriate expressions coming from above shows

$$\begin{aligned}e^{-b}(\dot{\mathbf{J}}_1 + \dot{\mathbf{J}}_2)e^b &= \begin{pmatrix} \dot{I} + \omega_I^{-1}\dot{b} & \omega_I^{-1}\dot{\omega}_I\omega_I^{-1} \\ \dot{\omega}_I + \{\dot{b}, I\} & -(I^* + \dot{b}\omega_I^{-1}) \end{pmatrix} \\ &= \frac{1}{2} \begin{pmatrix} \omega_J^{-1}KJ - \omega_I^{-1}KI & \omega_I^{-1}K\omega_I^{-1} + \omega_J^{-1}K\omega_J^{-1} \\ 4K - (I^*KI + J^*KJ) & I^*K\omega_I^{-1} - J^*K\omega_J^{-1} \end{pmatrix}, \\ e^{-b}(\dot{\mathbf{J}}_1 - \dot{\mathbf{J}}_2)e^b &= \begin{pmatrix} \dot{J} - \omega_J^{-1}\dot{b} & -\omega_J^{-1}\dot{\omega}_J\omega_J^{-1} \\ \{\dot{b}, J\} - \dot{\omega}_J & \dot{b}\omega_J^{-1} - \dot{J}^* \end{pmatrix} \\ &= \frac{1}{2} \begin{pmatrix} \omega_J^{-1}KI - \omega_I^{-1}KJ & -(\omega_I^{-1}K\omega_J^{-1} + \omega_J^{-1}K\omega_I^{-1}) \\ -(I^*KJ + J^*KI) & J^*K\omega_I^{-1} - I^*K\omega_J^{-1} \end{pmatrix}.\end{aligned}$$

Turning to the top right entries of each expression shows

$$\begin{aligned}\dot{\omega}_I &= \frac{1}{2}(K + \omega_I\omega_J^{-1}K\omega_J^{-1}\omega_I) = \frac{1}{2}(K + I^*J^*KJI) = \frac{1}{2}(K + I^*KI) = -\frac{1}{2}[K, I]I, \\ \dot{\omega}_J &= \frac{1}{2}(\omega_J\omega_I^{-1}K + K\omega_I^{-1}\omega_J) = -\frac{1}{2}(J^*I^*K + KIJ) = -\frac{1}{2}\{K, IJ\}.\end{aligned}$$

For the remaining data we differentiate the generalized metric  $\mathbf{G}_t = -(\mathbf{J}_1)_t(\mathbf{J}_2)_t = \begin{pmatrix} A_t & g_t^{-1} \\ \delta_t & A_t^* \end{pmatrix}$ , where  $g_t$  is the associated metric and  $b_t = -g_t A_t$ . It follows that

$$e^{-b}\dot{\mathbf{G}}e^b = \begin{pmatrix} \dot{A} - g^{-1}\dot{g}g^{-1}b & -g^{-1}\dot{g}g^{-1} \\ -b\dot{A} + \dot{A}^*b + \dot{\delta} - bg^{-1}\dot{g}g^{-1} & \dot{A}^* - bg^{-1}\dot{g}g^{-1} \end{pmatrix}.$$

Focusing on the top row, we furthermore compute

$$\begin{aligned}
e^{-b} \dot{\mathbf{G}} e^b &= -(e^{-b} \mathbf{j}_1 e^b)(e^{-b} \mathbf{J}_2 e^b) - (e^{-b} \mathbf{J}_1 e^b)(e^{-b} \mathbf{j}_2 e^b) \\
&= -\frac{1}{8} \begin{pmatrix} -(\omega_I^{-1} - \omega_J^{-1})K(I+J) & (\omega_I^{-1} - \omega_J^{-1})K(\omega_I^{-1} - \omega_J^{-1}) \\ 4K - (I^* + J^*)K(I+J) & (I^* + J^*)K(\omega_I^{-1} - \omega_J^{-1}) \end{pmatrix} \begin{pmatrix} I - J & -(\omega_I^{-1} + \omega_J^{-1}) \\ \omega_I + \omega_J & -(I^* - J^*) \end{pmatrix} \\
&\quad - \frac{1}{8} \begin{pmatrix} I + J & -(\omega_I^{-1} - \omega_J^{-1}) \\ \omega_I - \omega_J & -(I^* + J^*) \end{pmatrix} \begin{pmatrix} -(\omega_I^{-1} + \omega_J^{-1})K(I - J) & (\omega_I^{-1} + \omega_J^{-1})K(\omega_I^{-1} + \omega_J^{-1}) \\ 4K - (I^* - J^*)K(I - J) & (I^* - J^*)K(\omega_I^{-1} + \omega_J^{-1}) \end{pmatrix} \\
&= \frac{1}{2} \begin{pmatrix} g^{-1}K(I - J) + (\omega_I^{-1} - \omega_J^{-1})K & -(\omega_I^{-1} - \omega_J^{-1})Kg^{-1} - g^{-1}K(\omega_I^{-1} + \omega_J^{-1}) \\ * & * \end{pmatrix} \\
&= \frac{1}{2} \begin{pmatrix} g^{-1}\{K, I\} & -(\omega_I^{-1} - \omega_J^{-1})Kg^{-1} - g^{-1}K(\omega_I^{-1} + \omega_J^{-1}) \\ * & * \end{pmatrix}.
\end{aligned}$$

Thus

$$\begin{aligned}
\dot{g} &= \frac{1}{2}g(\omega_I^{-1} - \omega_J^{-1})K + \frac{1}{2}K(\omega_I^{-1} + \omega_J^{-1})g \\
&= \frac{1}{2}(I^*K - KI) - \frac{1}{2}(KJ + J^*K) \\
&= -\frac{1}{2}[K, I],
\end{aligned}$$

where we have used that  $K \in \Lambda_J^{1,1}$ . Then differentiating the formulas  $I/J = -\omega_{IJ}^{-1}g$  gives

$$\begin{aligned}
\dot{I} &= \omega_I^{-1} \dot{\omega}_I \omega_I^{-1} g - \omega_I^{-1} \dot{g} \\
&= -\frac{1}{2} \omega_I^{-1} [K, I] I \omega_I^{-1} g + \frac{1}{2} \omega_I^{-1} [K, I] = 0, \\
\dot{J} &= \omega_J^{-1} \dot{\omega}_J \omega_J^{-1} g - \omega_J^{-1} \dot{g} \\
&= -\frac{1}{2} \omega_J^{-1} \{K, IJ\} \omega_J^{-1} + \frac{1}{2} \omega_J^{-1} [K, I] \\
&= -\frac{1}{2} \omega_J^{-1} KI - \frac{1}{2} g^{-1} I^* KJ + \frac{1}{2} \omega_J^{-1} KI - \frac{1}{2} \omega_J^{-1} I^* K \\
&= \frac{1}{2} g^{-1} I^* J^* K - \frac{1}{2} g^{-1} J^* I^* K = \frac{1}{2} [I, J] g^{-1} K.
\end{aligned}$$

Lastly, differentiating the formula  $b = -gA$  yields

$$\begin{aligned}
\dot{b} &= -\dot{g}A - g\dot{A} \\
&= -\dot{g}A - g(g^{-1}\dot{g}g^{-1}b + \frac{1}{2}g^{-1}\{K, I\}) \\
&= -\frac{1}{2}\{K, I\}.
\end{aligned}$$

□

Using Proposition 3.6 we give the proof of Corollary 1.3.

*Proof of Corollary 1.3.* By Proposition 3.6 we know that  $\dot{I} = 0$ , and moreover it follows that

$$\begin{aligned}
\dot{\sigma} &= \frac{1}{2} \left( [I, J] g^{-1} K - [I, J] g^{-1} KI \right) g^{-1} + \frac{1}{2} [I, J] g^{-1} [K, I] g^{-1} \\
&= \frac{1}{2} [I, J] g^{-1} K g^{-1} - \frac{1}{2} [I, J] g^{-1} I^* K g^{-1} \\
&= \frac{1}{2} ([I, J] + [I, J] I) g^{-1} K g^{-1} \\
&= 0.
\end{aligned}$$

□

### 3.6 Examples

**Example 3.7.** Given  $(M^{2n}, g, J)$  a Kähler manifold, we interpret this as a generalized Kähler structure by setting  $I = J$ , and  $b = 0$ . Suppose  $(g_t, b_t, I_t, J_t)$  is a canonical family with this initial condition. Initially we have  $\sigma = 0$ , thus  $\sigma_t \equiv 0$  from Corollary 1.3. It follows that  $[I, J] \equiv 0$  for all times, so from the equations of Proposition 3.6 it follows that  $\dot{J} \equiv 0$  for all times, and so  $J_t \equiv J$ . In turn it follows easily that

$$\dot{\omega}_I = \dot{\omega}_J = K,$$

in other words, the complex structures stay fixed and the Kähler forms change by  $K$ . By the  $\sqrt{-1}\partial\bar{\partial}$ -Lemma, an exact canonical deformation satisfies  $K = da = dJdu$  for some  $u \in C^\infty(M)$ . Using the construction above and the normalization that  $\sqrt{-1}\partial\bar{\partial} = dId$  we verify item (1) of Proposition 1.2, noting that positivity of the Kähler forms  $(\omega_I)_t = \omega_I + tJdu$  is equivalent to the positivity condition (3) in Theorem 1.1.

**Example 3.8.** Suppose  $(M^{2n}, g, b, I, J)$  is a generalized Kähler structure and  $[I, J] \equiv 0$ . These manifolds are characterized by a holomorphic splitting of the tangent bundle and simple examples are given by quotients of products, for instance on Hopf surfaces. For further background on these structures see ([2]). Setting  $Q = -IJ$  we see that  $Q^2 = 1$  and so we can split  $T = T_+ \oplus T_-$  in terms of the  $\pm 1$ -eigenbundles of  $Q$ . Let  $J_+ = I|_{T_+}$  and  $J_- = I|_{T_-}$ . Since on  $T_\pm$  we have  $I = \pm J$ , it follows  $J|_{T_+} = J_+$  and  $J|_{T_-} = -J_-$ . Similarly, we define  $\omega_\pm = \omega_I|_{T_\pm}$ . Note that the Kähler form indeed block diagonalizes along  $T_\pm$  since

$$\omega_I(X_+, IY_-) = g(IX_+, IY_-) = g(JX_+, JY_-) = g(-IX_+, IY_-) = -\omega_I(X_+, IY_-) = 0.$$

Now suppose  $(g_t, b_t, I_t, J_t)$  is a canonical family with this initial condition. Arguing as above, since  $\sigma = 0$  for the given structure, it follows that  $[I, J] \equiv 0$  for all times  $t$ , and hence  $\dot{J} = 0$ . Thus along the variation we preserve the splitting induced by  $Q$ , and we can decompose  $K = K_{++} + K_{+-} + K_{-+} + K_{--}$ . Tracing through the formulas in Proposition 3.6 yields

$$\begin{aligned} \dot{g} &= \begin{pmatrix} J_+^* K_{++} & 0 \\ 0 & J_-^* K_{--} \end{pmatrix}, & \dot{b} &= \begin{pmatrix} 0 & -J_+^* K_{-+} \\ -J_-^* K_{+-} & 0 \end{pmatrix}, \\ \dot{\omega}_I &= \begin{pmatrix} K_{++} & 0 \\ 0 & K_{--} \end{pmatrix}, & \dot{\omega}_J &= \begin{pmatrix} -K_{++} & 0 \\ 0 & K_{--} \end{pmatrix}. \end{aligned}$$

A special case of this occurs when  $K = dJdu$ , yielding

$$\dot{\omega}_I = \sqrt{-1} (\partial_+ \bar{\partial}_+ - \partial_- \bar{\partial}_-) u,$$

where  $d = \partial_+ + \partial_- + \bar{\partial}_+ + \bar{\partial}_-$  is the fourfold splitting of  $d$  induced by  $Q$ . This construction verifies item (2) of Proposition 1.2, again noting that positivity of

$$(\omega_I)_t = \omega_I + t\sqrt{-1} (\partial_+ \bar{\partial}_+ - \partial_- \bar{\partial}_-) u$$

is equivalent to the positivity condition of item (3) in Theorem 1.1.

**Example 3.9.** Suppose  $(M^{2n}, g, b, I, J)$  is a generalized Kähler structure where the associated Poisson tensor  $\sigma$  is nondegenerate. In this setting the endomorphism  $[I, J]$  is invertible, and this in turn implies that  $I \pm J$  are invertible. We define the 2-forms  $F_\pm = -2g(I \pm J)^{-1}$ , which it turns out are closed, and furthermore let  $\Omega = \sigma^{-1}$ . It turns out that the three symplectic forms  $F_\pm, \Omega$  completely determine the generalized Kähler structure in this case. Direct computations show that the generalized complex structures can be expressed as

$$J_1 = e^{-4\Omega} \begin{pmatrix} 0 & -F_-^{-1} \\ F_- & 0 \end{pmatrix} e^{4\Omega}, \quad J_2 = \begin{pmatrix} 0 & -F_+^{-1} \\ F_+ & 0 \end{pmatrix},$$

Let  $K$  be an infinitesimal deformation of the generalized Kahler pair  $(J_1, J_2)$ . Then with respect to the data  $(F_+, F_-, \Omega)$ , direct computations show that

$$\dot{F}_+ = \dot{F}_- = K, \quad \dot{\Omega} = 0.$$

As in the above examples we can choose  $K_t = dJ_t du_t$ . We claim that for such a variation,

$$\dot{J} = L_{\sigma du} J.$$

To show this we compute, using that  $\sigma = \Omega^{-1}$  and  $d\Omega = 0$ ,

$$(L_{\sigma du} J) \Omega = L_{\sigma du} (J \Omega) - J L_{\sigma du} \Omega = dJdu = K.$$

Comparing against Proposition 3.6 we see that

$$\dot{J} = \sigma K = L_{\sigma du} J,$$

as required. It follows that  $J_t = \phi_t^* J_0$ , where  $\phi$  is the one-parameter family of  $\Omega$ -Hamiltonian diffeomorphisms driven by  $u_t$ . As discussed in [3], the positivity of  $-\text{Im} \pi_{\Lambda_I^{1,1}} \Omega_J$  is equivalent to the positivity condition (3) of Theorem 1.1. This verifies item (3) of Proposition 1.2.

*Proof of Proposition 1.2.* The three deformations claimed in the proposition are described in the examples above.  $\square$

## 4 Generalized Kähler-Ricci flow as canonical deformation

In this section we establish Theorem 1.4, namely that solutions to the generalized Kähler-Ricci flow are canonical deformations, driven by the Bismut Ricci curvature. This generalizes and unifies various instances of this phenomenon which have previously been observed. In particular, it is well known that Kähler-Ricci flow moves within the Kähler cone against a fixed complex structure, and so is a canonical deformation as in Example 3.7. Next, comparing Example 3.8 against the curvature identities of [17], we see that the generalized Kähler-Ricci flow in the case  $[I, J] = 0$  is a canonical deformation driven by  $\rho_I$ . Also, we can compare the discussion in Example 3.9 with [3] to see the phenomenon holds in the nondegenerate case. The key point in establishing the general case is to show that the evolution of  $J$  is indeed always determined by  $K = -\rho_I$ . This requires a delicate curvature identity we build up below.

**Lemma 4.1.** (cf. [15]) *Let  $(M^{2n}, g, I)$  be a pluriclosed structure. Then*

$$\begin{aligned} \rho_I(X, Y) &= -\text{Rc}^B(X, IY) - \nabla_X^B \theta(IY) \\ \rho_I^{1,1}(\cdot, I\cdot) &= \text{Rc}^g - \frac{1}{4} H^2 + \frac{1}{2} L_{\theta^\sharp} g, \\ \rho_I^{2,0+0,2}(X, IY) &= \frac{1}{2} \left( -d^* H(X, Y) + d^\nabla \theta(X, Y) \right) \\ &= (dI\theta)^{2,0+0,2}(X, Y). \end{aligned}$$

**Proposition 4.2.** *Let  $(M^{2n}, g, I, J)$  be a generalized Kähler structure. Then*

$$g \left( \left( L_{\theta^\sharp - \theta^\sharp} J \right) X, Y \right) = \rho_I(X, [I, J]Y).$$

*Proof.* We will use that for a Hermitian manifold  $(M^{2n}, g, J)$  one has

$$g((L_X J)Y, Z) = g((D_X J)Y - D_{JY}X + J D_Y X, Z). \tag{4.1}$$

First note, using that  $\rho_I \in \Lambda_J^{1,1}$ ,

$$\begin{aligned} \rho_I([I, J]X, JY) + \rho_I([I, J]JY, X) &= \rho_I(IJX - JIX, JY) - \rho_I(X, IJJY - JIJY) \\ &= \rho_I(IJX, JY) - \rho_I(IX, Y) + \rho_I(X, IY) - \rho_I(JX, IJY) \\ &= -\rho_I(JY, IJX) + \rho_I(Y, IX) + \rho_I(X, IY) - \rho_I(JX, IJY). \end{aligned}$$

Applying Lemma 4.1 we further compute

$$\begin{aligned}
& \rho_I(JY, IJX) - \rho_I(Y, IX) - \rho_I(X, IY) + \rho_I(JX, IJY) \\
&= \operatorname{Rc}^I(JY, JX) + \nabla_{JY}^I \theta^I(JX) - \operatorname{Rc}^I(Y, X) - \nabla_Y^I \theta^I(X) \\
&\quad - \operatorname{Rc}^I(X, Y) - \nabla_X^I \theta^I(Y) + \operatorname{Rc}^I(JX, JY) + \nabla_{JX}^I \theta^I(JY) \\
&= \operatorname{Rc}^J(JY, JX) + \operatorname{Rc}^J(JX, JY) - \operatorname{Rc}^J(X, Y) - \operatorname{Rc}^J(Y, X) \\
&\quad + \nabla_{JY}^I \theta^I(JX) - \nabla_Y^I \theta^I(X) - \nabla_X^I \theta^I(Y) + \nabla_{JX}^I \theta^I(JY) \\
&= \rho_J(JY, X) + \rho_J(JX, Y) + \rho_J(X, JY) + \rho_J(Y, JX) \\
&\quad - \nabla_{JY}^J \theta^J(JX) - \nabla_{JX}^J \theta^J(JY) + \nabla_X^J \theta^J(Y) + \nabla_Y^J \theta^J(X) \\
&\quad + \nabla_{JY}^J \theta^J(JX) - \nabla_Y^J \theta^J(X) - \nabla_X^J \theta^J(Y) + \nabla_{JX}^J \theta^J(JY) \\
&= D_X(\theta^J - \theta^I)(Y) + D_Y(\theta^J - \theta^I)(X) - D_{JX}(\theta^J - \theta^I)(JY) - D_{JY}(\theta^J - \theta^I)(JX) \\
&= g\left(\left(L_{\theta_j^\# - \theta_i^\#} J\right) X, JY\right) + g\left(\left(L_{\theta_j^\# - \theta_i^\#} J\right) JY, X\right).
\end{aligned}$$

where the last line follows by comparing (4.1).

To address the skew symmetric piece we first compute, again using that  $\rho_I \in \Lambda_J^{1,1}$ ,

$$\begin{aligned}
\rho_I([I, J]X, Y) - \rho_I([I, J]Y, X) &= \rho_I((IJ - JI)X, Y) + \rho_I(X, (IJ - JI)Y) \\
&= \rho_I(IJX, Y) + \rho_I(IX, JY) + \rho_I(X, IJY) + \rho_I(JX, IY) \\
&= 2\left((\rho_I)^{2,0}(JX, IY) + (\rho_I)^{2,0}(IX, JY)\right).
\end{aligned}$$

Now we compute using Lemma 4.1 and the last piece of equation (4.1),

$$\begin{aligned}
2\left((\rho_I)^{2,0}(JX, IY) + (\rho_I)^{2,0}(IX, JY)\right) &= -d^* H_I(JX, Y) + d^* H_I(JY, X) + d^{\nabla^I} \theta^I(JX, Y) - d^{\nabla^I} \theta^I(JY, X) \\
&= d^* H_J(JX, Y) - d^* H_J(JY, X) + d^{\nabla^I} \theta^I(JX, Y) - d^{\nabla^I} \theta^I(JY, X) \\
&= -2(\rho_J)^{2,0}(X, Y) + d^{\nabla^I} \theta^I(JX, Y) + 2(\rho_J)^{2,0}(Y, X) - d^{\nabla^I} \theta^I(JY, X) \\
&\quad + d^{\nabla^I} \theta^I(JX, Y) - d^{\nabla^I} \theta^I(JY, X) \\
&= -d\theta^J(JX, Y) + d\theta^J(JY, X) + d\theta^I(JX, Y) - d\theta^I(JY, X) \\
&= g\left(\left(L_{\theta_j^\# - \theta_i^\#} J\right) X, Y\right) - g\left(\left(L_{\theta_j^\# - \theta_i^\#} J\right) Y, X\right).
\end{aligned}$$

Combining the above equations yields the result.  $\square$

*Proof of Theorem 1.4.* We first assume that equations (1.2) hold. By Proposition 3.6 with  $K = -\rho_I$  one notes that  $I$  is fixed and one easily derives the evolution equation  $\dot{\omega}_I = -(\rho_I)^{1,1}$ . By Proposition 4.2 it follows that

$$\dot{J} = -\frac{1}{2}[I, J]g^{-1}\rho_I = L_{\frac{1}{2}(\theta_j^\# - \theta_i^\#)}J.$$

as required. Thus we have verified the evolution equations of (1.1). Assuming equations (1.1) and imposing  $\dot{b} = -\{\rho_B^I, I\}$ , we can apply the computations in Proposition 3.6 and use Proposition 4.2 to establish equations (1.2).  $\square$

## References

- [1] Apostolov, V., Gauduchon, P., and Grantcharov, G. Bi-Hermitian structures on complex surfaces. *Proc. London Math. Soc.* (3) 79, 2 (1999), 414–428.
- [2] Apostolov, V., and Gualtieri, M. Generalized Kähler manifolds, commuting complex structures, and split tangent bundles. *Comm. Math. Phys.* 271, 2 (2007), 561–575.

- [3] Apostolov, V., and Streets, J. The nondegenerate generalized Kähler Calabi-Yau problem. *ArXiv e-prints* (Mar. 2017).
- [4] Bischoff, F. Morita equivalence and Generalized Kähler geometry Diss. Univ. Toronto
- [5] Bischoff, F., Gualtieri, M., and Zabzine, M. *Morita equivalence and the generalized Kahler potential* to appear in Journal of Differential Geometry
- [6] Brieskorn, E. Über holomorphe  $P_n$ -Bündel über  $P_1$ . *Math. Ann.* 157 (1965), 343–357.
- [7] Demailly, J.-P., and Paun, M. Numerical characterization of the Kähler cone of a compact Kähler manifold. *Ann. of Math. (2)* 159, 3 (2004), 1247–1274.
- [8] Gates, Jr., S. J., Hull, C. M., and Roček, M. Twisted multiplets and new supersymmetric nonlinear  $\sigma$ -models. *Nuclear Phys. B* 248, 1 (1984), 157–186.
- [9] Goto, R. Deformations of generalized complex and generalized Kähler structures. *J. Differential Geom.* 84, 3 (2010), 525–560.
- [10] Gualtieri, M. Branes on Poisson varieties *The many facets of geometry*, Oxford Univ. Press, 2010
- [11] Gualtieri, M. *Generalized complex geometry*. PhD thesis, St. John’s College, University of Oxford, 11 2003.
- [12] Gualtieri, M. Generalized Kähler metrics from Hamiltonian deformations *Geometry and Physics: A Festschrift in Honour of Nigel Hitchin, Vol. 1*, Oxford University Press, 2018
- [13] Hitchin, N. Generalized Calabi-Yau manifolds. *Q. J. Math.* 54, 3 (2003), 281–308.
- [14] Hitchin, N. Bihermitian metrics on del Pezzo surfaces. *J. Symplectic Geom.* 5, 1 (2007), 1–8.
- [15] Ivanov, S., and Papadopoulos, G. Vanishing theorems and string backgrounds. *Classical Quantum Gravity* 18, 6 (2001), 1089–1110.
- [16] Streets, J., Pluriclosed flow, Born-Infeld geometry, and rigidity results for generalized Kähler manifolds, *Comm. Partial Differential Equations* 41 (2016), no. 2, 318–374.
- [17] Streets, J. Pluriclosed flow on generalized Kähler manifolds with split tangent bundle. *J. Reine Angew. Math.* 739 (2018), 241–276.
- [18] Streets, J., and Tian, G. Generalized Kähler geometry and the pluriclosed flow. *Nuclear Phys. B* 858, 2 (2012), 366–376.