

Stefan Richter* and Brett D. Wick

A remark on the multipliers on spaces of Weak Products of functions

DOI 10.1515/conop-2016-0004

Received October 30, 2015; accepted January 27, 2016.

Abstract: If \mathcal{H} denotes a Hilbert space of analytic functions on a region $\Omega \subseteq \mathbb{C}^d$, then the weak product is defined by

$$\mathcal{H} \odot \mathcal{H} = \left\{ h = \sum_{n=1}^{\infty} f_n g_n : \sum_{n=1}^{\infty} \|f_n\|_{\mathcal{H}} \|g_n\|_{\mathcal{H}} < \infty \right\}.$$

We prove that if \mathcal{H} is a first order holomorphic Besov Hilbert space on the unit ball of \mathbb{C}^d , then the multiplier algebras of \mathcal{H} and of $\mathcal{H} \odot \mathcal{H}$ coincide.

Keywords: Dirichlet space, Drury-Arveson space, Weak product, Multiplier

MSC: 47B37

1 Introduction

Let d be a positive integer and let $R = \sum_{i=1}^d z_i \frac{\partial}{\partial z_i}$ denote the radial derivative operator. For $s \in \mathbb{R}$ the holomorphic Besov space B_s is defined to be the space of holomorphic functions f on the unit ball \mathbb{B}_d of \mathbb{C}^d such that for some nonnegative integer $k > s$

$$\|f\|_{k,s}^2 = \int_{\mathbb{B}_d} |(I + R)^k f(z)|^2 (1 - |z|^2)^{2(k-s)-1} dV(z) < \infty.$$

Here dV denotes Lebesgue measure on \mathbb{B}_d . It is well-known that for any $f \in \text{Hol}(\mathbb{B}_d)$ and any $s \in \mathbb{R}$ the quantity $\|f\|_{k,s}$ is finite for some nonnegative integer $k > s$ if and only if it is finite for all nonnegative integers $k > s$, and that for each $k > s$ $\|\cdot\|_{k,s}$ defines a norm on B_s , and that all these norms are equivalent to one another, see [2]. For $s < 0$ one can take $k = 0$ and these spaces are weighted Bergman spaces. In particular, $B_{-1/2} = L_a^2(\mathbb{B}_d)$ is the unweighted Bergman space. For $s = 0$ one obtains the Hardy space of \mathbb{B}_d and one has that for each $k \geq 1$ $\|f\|_{k,0}^2$ is equivalent to $\int_{\partial\mathbb{B}_d} |f|^2 d\sigma$, where σ is the rotationally invariant probability measure on $\partial\mathbb{B}_d$. We also note that for $s = (d-1)/2$ we have $B_s = H_d^2$, the Drury-Arveson space. If $d = 1$ and $s = 1/2$, then $B_s = D$, the classical Dirichlet space of the unit disc.

Let $\mathcal{H} \subseteq \text{Hol}(\mathbb{B}_d)$ be a reproducing kernel Hilbert space such that $1 \in \mathcal{H}$. The weak product of \mathcal{H} is denoted by $\mathcal{H} \odot \mathcal{H}$ and it is defined to be the collection of all functions $h \in \text{Hol}(\mathbb{B}_d)$ such that there are sequences $\{f_i\}_{i \geq 1}, \{g_i\}_{i \geq 1} \subseteq \mathcal{H}$ with $\sum_{i=1}^{\infty} \|f_i\|_{\mathcal{H}} \|g_i\|_{\mathcal{H}} < \infty$ and for all $z \in \mathbb{B}_d$, $h(z) = \sum_{i=1}^{\infty} f_i(z)g_i(z)$.

*Corresponding Author: Stefan Richter: Department of Mathematics, University of Tennessee, Knoxville, TN 37996, USA,

E-mail: richter@math.utk.edu

Brett D. Wick: Department of Mathematics, Washington University – St. Louis, St. Louis, MO 63130, USA and School of Mathematics, Georgia Institute of Technology, Atlanta GA 30332-0160, USA, E-mail: wick@math.wustl.edu

We define a norm on $\mathcal{H} \odot \mathcal{H}$ by

$$\|h\|_* = \inf \left\{ \sum_{i=1}^{\infty} \|f_i\|_{\mathcal{H}} \|g_i\|_{\mathcal{H}} : h(z) = \sum_{i=1}^{\infty} f_i(z)g_i(z) \text{ for all } z \in \mathbb{B}_d \right\}.$$

In what appears below we will frequently take $\mathcal{H} = B_s$, and will use the same notation for this weak product.

Weak products have their origin in the work of Coifman, Rochberg, and Weiss [5]. In the frame work of the Hilbert space \mathcal{H} one may consider the weak product to be an analogue of the Hardy H^1 -space. For example, one has $H^2(\partial\mathbb{B}_d) \odot H^2(\partial\mathbb{B}_d) = H^1(\partial\mathbb{B}_d)$ and $L_a^2(\mathbb{B}_d) \odot L_a^2(\mathbb{B}_d) = L_a^1(\mathbb{B}_d)$, see [5]. For the Dirichlet space D the weak product $D \odot D$ has recently been considered in [1, 3, 6, 7, 9]. The space $H_d^2 \odot H_d^2$ was used in [10]. For further motivation and general background on weak products we refer the reader to [1] and [9].

Let \mathcal{B} be a Banach space of analytic functions on \mathbb{B}_d such that point evaluations are continuous and such that $1 \in \mathcal{B}$. We use $M(\mathcal{B})$ to denote the multiplier algebra of \mathcal{B} ,

$$M(\mathcal{B}) = \{\varphi : \varphi f \in \mathcal{B} \text{ for all } f \in \mathcal{B}\}.$$

The multiplier norm $\|\varphi\|_M$ is defined to be the norm of the associated multiplication operator $M_\varphi : \mathcal{B} \rightarrow \mathcal{B}$. It is easy to check and is well-known that $M(\mathcal{B}) \subseteq H^\infty(\mathbb{B}_d)$, and that for $s \leq 0$ we have $M(B_s) = H^\infty(\mathbb{B}_d)$. For $s > d/2$ the space B_s is an algebra [2], hence $B_s = M(B_s)$, but for $0 < s \leq d/2$ one has $M(B_s) \subsetneq B_s \cap H^\infty(\partial\mathbb{B}_d)$. For those cases $M(B_s)$ has been described by a certain Carleson measure condition, see [4, 8].

It is easy to see that $M(\mathcal{H}) \subseteq M(\mathcal{H} \odot \mathcal{H}) \subseteq H^\infty$ (see Proposition 3.1). Thus, if $s \leq 0$, then $M(B_s) = M(B_s \odot B_s) = H^\infty$. Furthermore, if $s > d/2$, then $B_s = B_s \odot B_s = M(B_s)$ since B_s is an algebra. This raises the question whether $M(B_s)$ and $M(B_s \odot B_s)$ always agree. We prove the following:

Theorem 1.1. *Let $s \in \mathbb{R}$ and $d \in \mathbb{N}$. If $s \leq 1$ or $d \leq 2$, then $M(B_s) = M(B_s \odot B_s)$.*

Note that when $d \leq 2$, then B_s is an algebra for all $s > 1$. Thus for each $d \in \mathbb{N}$ the nontrivial range of the Theorem is $0 < s \leq 1$. If $d = 1$ then the theorem applies to the classical Dirichlet space of the unit disc and for $d \leq 3$ it applies to the Drury-Arveson space.

2 Preliminaries

For $z = (z_1, \dots, z_d) \in \mathbb{C}^d$ and $t \in \mathbb{R}$ we write $e^{it}z = (e^{it}z_1, \dots, e^{it}z_d)$ and we write $\langle z, w \rangle$ for the inner product in \mathbb{C}^d . Furthermore, if h is a function on \mathbb{B}_d , then we define $T_t f$ by $(T_t f)(z) = f(e^{it}z)$. We say that a space $\mathcal{H} \subseteq \text{Hol}(\mathbb{B}_d)$ is radially symmetric, if each T_t acts isometrically on \mathcal{H} and if for all $t_0 \in \mathbb{R}$, $T_t \rightarrow T_{t_0}$ in the strong operator topology as $t \rightarrow t_0$, i.e. if $\|T_t f\|_{\mathcal{H}} = \|f\|_{\mathcal{H}}$ and $\|T_t f - T_{t_0} f\|_{\mathcal{H}} \rightarrow 0$ for all $f \in \mathcal{H}$. For example, for each $s \in \mathbb{R}$ the holomorphic Besov space B_s is radially symmetric when equipped with any of the norms $\|\cdot\|_{k,s}$, $k > s$.

It is elementary to verify the following lemma.

Lemma 2.1. *If $\mathcal{H} \subseteq \text{Hol}(\mathbb{B}_d)$ is radially symmetric, then so is $\mathcal{H} \odot \mathcal{H}$.*

Note that if h and φ are functions on \mathbb{B}_d , then for every $t \in \mathbb{R}$ we have $(T_t \varphi)h = T_t(\varphi T_{-t}h)$, hence if a space is radially symmetric, then T_t acts isometrically on the multiplier algebra. For $0 < r < 1$ we write $f_r(z) = f(rz)$.

Lemma 2.2. *If $\mathcal{H} \subseteq \text{Hol}(\mathbb{B}_d)$ is radially symmetric, and if $\varphi \in M(\mathcal{H} \odot \mathcal{H})$, then for all $0 < r < 1$ we have $\|\varphi_r\|_{M(\mathcal{H} \odot \mathcal{H})} \leq \|\varphi\|_{M(\mathcal{H} \odot \mathcal{H})}$.*

Proof. Let $\varphi \in M(\mathcal{H} \odot \mathcal{H})$ and $h \in \mathcal{H} \odot \mathcal{H}$, then for $0 < r < 1$ we have

$$\varphi_r h = \int_{-\pi}^{\pi} \frac{1-r^2}{|1-re^{it}|^2} (T_t \varphi)h \frac{dt}{2\pi}.$$

This implies

$$\|\varphi_r h\|_* \leq \int_{-\pi}^{\pi} \frac{1-r^2}{|1-re^{it}|^2} \|(T_t \varphi)h\|_* \frac{dt}{2\pi} \leq \|\varphi\|_{M(\mathcal{H} \odot \mathcal{H})} \|h\|_*.$$

Thus, $\|\varphi_r\|_{M(\mathcal{H} \odot \mathcal{H})} \leq \|\varphi\|_{M(\mathcal{H} \odot \mathcal{H})}$. □

3 Multipliers

The following Proposition is elementary.

Proposition 3.1. *We have $M(\mathcal{H}) \subseteq M(\mathcal{H} \odot \mathcal{H}) \subseteq H^\infty$ and if $\varphi \in M(\mathcal{H})$, $\|\varphi\|_{M(\mathcal{H} \odot \mathcal{H})} \leq \|\varphi\|_{M(\mathcal{H})}$.*

As explained in the Introduction, the following will establish Theorem 1.1.

Theorem 3.2. *Let $0 < s \leq 1$. Then $M(B_s) = M(B_s \odot B_s)$ and there is a $C_s > 0$ such that*

$$\|\varphi\|_{M(B_s \odot B_s)} \leq \|\varphi\|_{M(B_s)} \leq C_s \|\varphi\|_{M(B_s \odot B_s)}$$

for all $\varphi \in M(B_s)$.

Here for each s we have the norm on B_s to be $\|\cdot\|_{k,s}$, where k is the smallest natural number $> s$.

Proof. We first do the case $0 < s < 1$. Then $k = 1$, and $\|f\|_{B_s}^2 = \int_{\mathbb{B}_d} |(I+R)f(z)|^2 dV_s(z)$, where $dV_s(z) = (1-|z|^2)^{1-2s} dV(z)$. For later reference we note that a short calculation shows that $\int_{\mathbb{B}_d} |Rf|^2 dV_s \leq \|f\|_{B_s}^2$.

We write $\|R\varphi\|_{C\alpha(B_s)}$ for the Carleson measure norm of $|R\varphi|^2$, i.e.

$$\|R\varphi\|_{C\alpha(B_s)}^2 = \inf \left\{ C > 0 : \int_{\mathbb{B}_d} |f|^2 |R\varphi|^2 dV_s \leq C \|f\|_{B_s}^2 \text{ for all } f \in B_s \right\}.$$

Since $\|\varphi f\|_{B_s}^2 = \int_{\mathbb{B}_d} |\varphi(z)(I+R)f(z) + f(z)R\varphi(z)|^2 dV_s(z)$ it is clear that $\|\varphi\|_{M(B_s)}$ is equivalent to $\|\varphi\|_\infty + \|R\varphi\|_{C\alpha(B_s)}$. Thus, it suffices to show that there is a $c > 0$ such that $\|R\varphi\|_{C\alpha(B_s)} \leq c \|\varphi\|_{M(B_s \odot B_s)}$ for all $\varphi \in M(B_s \odot B_s)$.

First we note that if b is holomorphic in a neighborhood of $\overline{\mathbb{B}_d}$ and $h = \sum_{i=1}^{\infty} f_i g_i \in B_s \odot B_s$, then

$$\begin{aligned} \int_{\mathbb{B}_d} |(Rh)Rb| dV_s &\leq \sum_{i=1}^{\infty} \int_{\mathbb{B}_d} |(Rf_i)g_i Rb| dV_s + \int_{\mathbb{B}_d} |(Rg_i)f_i Rb| dV_s \\ &\leq \sum_{i=1}^{\infty} \|f_i\|_{B_s} \left(\int_{\mathbb{B}_d} |g_i Rb|^2 dV_s \right)^{1/2} + \|g_i\|_{B_s} \left(\int_{\mathbb{B}_d} |f_i Rb|^2 dV_s \right)^{1/2} \\ &\leq 2 \sum_{i=1}^{\infty} \|f_i\|_{B_s} \|g_i\|_{B_s} \|Rb\|_{C\alpha(B_s)}. \end{aligned}$$

Hence

$$\int_{\mathbb{B}_d} |(Rh)Rb| dV_s \leq 2 \|h\|_* \|Rb\|_{C\alpha(B_s)},$$

where we have continued to write $\|\cdot\|_*$ for $\|\cdot\|_{B_s \odot B_s}$.

Let $\varphi \in M(B_s \odot B_s)$ and let $0 < r < 1$. Then for all $f \in B_s$ we have $f^2, \varphi_r f^2 \in B_s \odot B_s$, hence

$$\int_{\mathbb{B}_d} |f|^2 |R\varphi_r|^2 dV_s = \int_{\mathbb{B}_d} |R(\varphi_r f^2) - \varphi_r R(f^2)| |R\varphi_r| dV_s$$

$$\begin{aligned}
&\leq 2(\|\varphi_r f^2\|_* + \|\varphi\|_\infty \|f^2\|_*) \|R\varphi_r\|_{Ca(B_s)} \\
&\leq 2(\|\varphi\|_{M(B_s \odot B_s)} \|f^2\|_* + \|\varphi\|_\infty \|f^2\|_*) \|R\varphi_r\|_{Ca(B_s)} \\
&\leq 4\|\varphi\|_{M(B_s \odot B_s)} \|f\|_{B_s}^2 \|R\varphi_r\|_{Ca(B_s)}.
\end{aligned}$$

Next we take the sup of the left hand side of this expression over all f with $\|f\|_{B_s} = 1$ and we obtain $\|R\varphi_r\|_{Ca(B_s)}^2 \leq 4\|\varphi\|_{M(B_s \odot B_s)} \|R\varphi_r\|_{Ca(B_s)}$ which implies that $\|R\varphi_r\|_{Ca(B_s)} \leq 4\|\varphi\|_{M(B_s \odot B_s)}$ holds for all $0 < r < 1$. Thus, for $0 < s < 1$ the result follows from Fatou's lemma as $r \rightarrow 1$.

If $s = 1$, then $\|f\|_{2,1}^2 \sim \int_{\partial\mathbb{B}_d} |(I + R)f(z)|^2 d\sigma(z)$ and the argument proceeds as above. \square

Acknowledgement: B.D. Wick's research supported in part by National Science Foundation DMS grant #1603246 and #1560955.

This work was supported by the LABEX MILYON (ANR-10-LABX-0070) of Université de Lyon, within the program "Investissements d'Avenir" (ANR-11-IDEX-0007) operated by the French National Research Agency (ANR).

References

- [1] Arcozzi, Nicola and Rochberg, Richard and Sawyer, Eric and Wick, Brett D., Bilinear forms on the Dirichlet space, *Anal. PDE*, **3**, 2010, no. 1, 21–47.
- [2] Beatrous, Frank and Burbea, Jacob, Holomorphic Sobolev spaces on the ball, *Dissertationes Math. (Rozprawy Mat.)*, Polska Akademia Nauk. Instytut Matematyczny. Dissertationes Mathematicae. Rozprawy Matematyczne, **276**, 1989.
- [3] Cascante, Carme and Ortega, Joaquín M., On a characterization of bilinear forms on the Dirichlet space, *Proc. Amer. Math. Soc.*, **140**, 2012, no. 7, 2429–2440.
- [4] Cascante, Carme and Fàbrega, Joan and Ortega, Joaquín M., On weighted Toeplitz, big Hankel operators and Carleson measures, *Integral Equations Operator Theory*, **66**, 2010, no.4, 495–528.
- [5] Coifman, R. R. and Rochberg, R. and Weiss, Guido, Factorization theorems for Hardy spaces in several variables, *Ann. of Math. (2)*, **103**, 1976, no. 3, 611–635.
- [6] Luo, Shuaibing and Richter, Stefan, Hankel operators and invariant subspaces of the Dirichlet space, *J. Lond. Math. Soc. (2)*, **91**, 2015, no. 2, 423–438.
- [7] Luo, Shuaibing, On the Index of Invariant Subspaces in the Space of Weak Products of Dirichlet Functions, *Complex Anal. Oper. Theory*, **9**, 2015, no. 6, 1311–1323.
- [8] Ortega, Joaquín and Fàbrega, Joan, Multipliers in Hardy-Sobolev spaces, *Integral Equations Operator Theory*, **55**, 2006, no. 4, 535–560.
- [9] Richter, Stefan and Sundberg, Carl, Weak products of Dirichlet functions, *J. Funct. Anal.*, **266**, 2014, no. 8, 5270–5299.
- [10] Richter, Stefan and Sunkes, James, Hankel operators, Invariant subspaces, and cyclic vectors in the Drury-Arveson space, *Proc. Amer. Math. Soc.*, *Proc. Amer. Math. Soc.*, to appear.