Clinical Image

John Carter*, DO, Jessica Ball, DO, MPH, George McKean, DO and Samir Patel, MD, DPM

Epiglottic squamous cell carcinoma

<https://doi.org/10.1515/jom-2021-0071>

Received February 26, 2021; accepted July 1, 2021;

published online July 15, 2021

A 56 year old woman with history of epiglottic squamous cell cancer (SCC), chronic obstructive pulmonary disease (COPD), requiring 4 L O₂ supplementation, asthma, coronary artery disease, myocardial infarction, gastroesophageal reflux disease, dysphagia, and tobacco abuse presented for laparoscopic jejunostomy tube placement to supplement nutritional intake during radiation therapy. Less than one month prior to presentation, she underwent awake intubation utilizing video laryngoscopy with a full view of the glottis. A computed tomography scan of the neck demonstrated a 3.1 × 1.8 cm epiglottic mass (Image A) and nasolaryngoscopy revealed a mass on the epiglottis to the mid-laryngeal surface without obstruction of the glottis. Given the patient's reassuring nasolaryngoscopy examination and prior successful intubation, intubation for the jejunostomy tube placement was performed utilizing video laryngoscopy. After induction, easy bag-mask ventilation was confirmed before succinylcholine, a short-acting neuromuscular blocker, was given to optimize intubating conditions. Using a Glidescope® LoPro video laryngoscope (Verathon Inc.), the mass was visualized to be partially obstructing the laryngeal inlet (Image B); however, intubation went without complication. On postoperative day 4, the patient stated that she no longer wanted to move forward with radiation therapy as she felt that she was experiencing significant weakness. At that time, the patient changed her code status to Do-Not-Resuscitate Comfort Care and elected hospice care. Unfortunately, the patient expired on postoperative day 5.

SCC is the most common malignancy in the head and neck [1, 2]. Alcohol and tobacco abuse are the most prevalent risk factors, contributing to approximately 75–85% of cases [3]. Human papilloma virus (HPV) infection, primarily HPV-16, is a growing cause of head and neck SCC,

specifically oropharyngeal cancer with the prevalence of HPV-positive oropharyngeal SCC in Europe increasing to 50% recently and North America plateauing at approximately 65% in the last decade [3, 4]. Presenting symptoms for laryngeal SCC include dysphagia, dysphonia, referred otalgia, and neck mass [5]. More advanced tumors may present with airway compromise due to obstruction [5].

Image A: CT neck (axial view) demonstrates the presence of an epiglottic mass (denoted by the red arrow).

Image B: Video laryngoscopy of the patient's airway demonstrated an epiglottic mass partially obstructing the view of the glottic opening.

***Corresponding author: John Carter, DO,** Department of Anesthesiology, Doctors Hospital, 5100 West Broad Street, Columbus, OH, 43228, USA, E-mail: john.carter2@ohiohealth.com

Jessica Ball, DO, MPH, Department of Otolaryngology, Doctors Hospital, Columbus, OH, USA

George McKean, DO and Samir Patel, MD, DPM, Department of Anesthesiology, Doctors Hospital, Columbus, OH, USA

Prevalent comorbidities at diagnosis often include hypertension, hyperlipidemia, COPD, and diabetes [6]. Treatment options for laryngeal cancer are determined by subsite, stage, and pathologic findings [7]. Early stage laryngeal cancers may be treated with surgery or radiation while advanced stage laryngeal cancers require combined modality therapy with either chemoradiation or surgery and adjuvant radiation [7, 8]. As demonstrated in this case, pathologies that can distort the airway must be considered to ensure proper airway management.

Research funding: None reported.

Author contributions: All authors provided substantial contributions to conception and design, acquisition of data, or analysis and interpretation of data; all authors drafted the article or revised it critically for important intellectual content; all authors gave final approval of the version of the article to be published; and all authors agree to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Competing interests: None reported.

Informed consent: The patient in this report provided written informed consent.

References

1. Landry D, Glastonbury C. Squamous cell carcinoma of the upper aerodigestive tract: a review. *Radiol Clin* 2015;53:81–97.
2. Farsi NJ, Rousseau MC, Schlecht N, Castonguay G, Allison P, Nguyen-Tan PF, et al. Aetiological heterogeneity of head and neck squamous cell carcinomas: the role of human papillomavirus infections, smoking and alcohol. *Carcinogenesis* 2017;38:1188–95.
3. Machiels JP, Leemans CR, Golusinski W, Grau C, Licitra L, Gregoire V, et al. Squamous cell carcinoma of the oral cavity, larynx, oropharynx and hypopharynx: EHNS-ESMO-ESTRO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol* 2020;31:1462–75.
4. Stein A, Saha S, Kraninger JL, Swick AD, Yu M, Lambert PF, et al. Prevalence of human papillomavirus in oropharyngeal cancer: a systematic review. *Canc J* 2015;21:138–46.
5. Obid R, Redlich M, Tomeh C. The treatment of laryngeal cancer. *Oral Maxillofac Surg Clin* 2019;31:1–11.
6. Eytan D, Blackford A, Eisele D, Fakhry C. Prevalence of comorbidities among older head and neck cancer survivors in the United States. *Otolaryngol Head Neck Surg* 2019;160:85–92.
7. Pfister D, Spencer S, Adelstein D, Adkins D, Anzai Y, Brizel DM, et al. Head and neck cancers, version 2.2020, NCCN clinical practice guidelines in oncology. *J Natl Compr Canc Netw* 2020;18:873–98.
8. Patel S, Qureshi M, Dyer MA, Jalisi S, Grillone G, Truong MT. Comparing surgical and nonsurgical larynx-preserving treatments with total laryngectomy for locally advanced laryngeal cancer. *Cancer* 2019;125:3367–77.