

Clinical Image

Austin B. Ambur*, DO, Timothy A. Nyckowski, DO and Aashni Bhukhan, BS

Annular bullous lesions in a child from Uganda: chronic bullous disease of childhood

<https://doi.org/10.1515/jom-2022-0168>

Received August 13, 2022; accepted September 22, 2022;
published online November 2, 2022

A 10-year-old male presented to our global outreach clinic in Uganda in May 2018 with a recurrent pruritic blistering rash on the lower leg of 1 year in duration. His medical history was unremarkable for autoimmune conditions or exposure to medications. Cutaneous examination showed urticarial plaques and numerous tense bullae in an annular arrangement on the left lower leg (Figure 1). Direct immunofluorescence (DIF) demonstrated a linear band of IgA at the dermoepidermal junction, consistent with a diagnosis of chronic bullous disease of childhood (CBDC). Oral dapsone was initiated with full resolution over the following week.

CBDC is a subepidermal blistering condition driven by autoimmune production of IgA. The disease is characterized by the abrupt onset of edematous plaques and large tense bullae, typically in the first decade of life [1]. Lesions are distributed in an annular, or “crown of jewels,” arrangement, and they commonly affect the trunk and limbs [1]. Although children present with a cutaneous distribution, neonates may have more mucocutaneous involvement [2]. DIF classically demonstrates linear IgA deposition along the basement membrane zone (BMZ) [3]. Although the disease is often idiopathic, genetic susceptibility, infections, drugs, and vaccinations have been implicated in the pathogenesis [1, 3]. Bullous impetigo, epidermolysis bullosa acquisita, and other autoimmune bullous disorders can be considered as differential



Figure 1: Urticarial plaques and numerous tense bullae in an annular arrangement on the lower left leg and foot.

diagnoses [1]. Although the condition can follow a relapsing and remitting course for several years, spontaneous resolution is common before puberty [4]. Treatment options may include dapsone and topical steroids.

Research funding: None reported.

Author contributions: All authors provided substantial contributions to conception and design, acquisition of data, or analysis and interpretation of data; all authors drafted the article or revised it critically for important intellectual content; all authors gave final approval of the version of the article to be published; and all authors agree to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Competing interests: None reported.

*Corresponding author: Austin B. Ambur, DO, KCU-GME, Advanced Dermatology and Cosmetic Surgery, 1410 W Broadway Street, Suite 205, Oviedo, FL 32765, USA, E-mail: austin.b.ambur@gmail.com.
<https://orcid.org/0000-0003-2191-7224>

Timothy A. Nyckowski, DO, Kansas City University at Advanced Dermatology and Cosmetic Surgery, Orlando, FL, USA

Aashni Bhukhan, BS, Nova Southeastern University Dr. Kiran C. Patel College of Osteopathic Medicine, Fort Lauderdale, FL, USA

Informed consent: A parent of the patient described in this report provided written informed consent.

References

1. Mintz EM, Morel KD. Clinical features, diagnosis, and pathogenesis of chronic bullous disease of childhood. *Dermatol Clin* 2011;29:459–ix.
2. Wilson A, Murrell DF. Clinical features of chronic bullous dermatosis of childhood. *Clin Exp Dermatol* 2022;47:873–81.
3. Mori F, Saretta F, Liotti L, Giovannini M, Castagnoli R, Arasi S, et al. Linear immunoglobulin a bullous dermatosis in children. *Front Pediatr* 2022;10:937528.
4. Tate C, Christian W, Newell L. Chronic bullous dermatosis of childhood and the string of pearls sign. *J Pediatr* 2018;202:325–e1.