Research Article

Ahmed Alsaedi, Bashir Ahmad*, Badrah Alghamdi, and Sotiris K. Ntouyas

On a nonlinear system of Riemann-Liouville fractional differential equations with semi-coupled integro-multipoint boundary conditions

<https://doi.org/10.1515/math-2021-0069>

received February 16, 2021; accepted June 29, 2021

Abstract: We study a nonlinear system of Riemann-Liouville fractional differential equations equipped with nonseparated semi-coupled integro-multipoint boundary conditions. We make use of the tools of the fixed-point theory to obtain the desired results, which are well-supported with numerical examples.

Keywords: Riemann-Liouville fractional derivative, integro-differential inclusions, nonlocal multi-point boundary conditions, existence, fixed point theorems

MSC 2020: 34A08, 34B15

1 Introduction

Nonlinear boundary value problems involving Riemann-Liouville fractional derivatives have been studied by many researchers, for example, see [1–6]. In a recent article [7], the authors discussed the existence and Ulam-type stability for nonlinear Riemann-Liouville fractional differential equations with constant delay.

The nonlocal nature of fractional derivative operators significantly contributed to the popularity of fractional calculus. Nowadays, one can find extensive application of fractional-order operators in the mathematical models of several real-world phenomena occurring in physical and applied sciences, such as continuum mechanics [8], bioengineering [9], financial economics [10], fractals [11], etc.

The topic of fractional differential systems also received considerable attention in view of their applications in diverse fields such as anomalous diffusion [12], disease models [13] biological models [14], hybrid systems [15], rheological models [16], diffusion systems [17], ecological models [18], etc. For theoretical details of such systems, see [19–29].

In this paper, we investigate the existence of solutions for a system of nonlinear Riemann-Liouville fractional differential equations

$$\begin{cases} D^\alpha u(t) = F(t, u(t), v(t)), & 1 < \alpha \leq 2, \quad t \in [0, T], \\ D^\beta v(t) = G(t, u(t), v(t)), & 1 < \beta \leq 2, \quad t \in [0, T], \end{cases} \quad (1)$$

* **Corresponding author: Bashir Ahmad**, Department of Mathematics, Faculty of Science, Nonlinear Analysis and Applied Mathematics (NAAM)-Research Group, King Abdulaziz University, P.O. Box 80203, Jeddah 21589, Saudi Arabia, e-mail: bashirahmad-qau@yahoo.com

Ahmed Alsaedi: Department of Mathematics, Faculty of Science, Nonlinear Analysis and Applied Mathematics (NAAM)-Research Group, King Abdulaziz University, P.O. Box 80203, Jeddah 21589, Saudi Arabia, e-mail: aalsaedi@hotmail.com

Badrah Alghamdi: Department of Mathematics, Faculty of Science, Nonlinear Analysis and Applied Mathematics (NAAM)-Research Group, King Abdulaziz University, P.O. Box 80203, Jeddah 21589, Saudi Arabia, e-mail: bsalghamdi@kau.edu.sa

Sotiris K. Ntouyas: Department of Mathematics, University of Ioannina, 451 10, Ioannina, Greece, e-mail: sntouyas@uoi.gr

equipped with nonlocal semi-coupled fractional integro-multipoint boundary conditions of the form:

$$\begin{cases} D^{\alpha-2}u(0^+) + a_0 D^{\alpha-2}u(T^-) = \lambda_1, \\ D^{\alpha-1}u(0^+) + a_1 D^{\alpha-1}u(T^-) = \nu I^{\alpha-1}v(\eta_1) + \sum_{i=1}^m \mu_i v(\xi_i), \\ D^{\beta-2}v(0^+) + b_0 D^{\beta-2}v(T^-) = \lambda_2, \\ D^{\beta-1}v(0^+) + b_1 D^{\beta-1}v(T^-) = \mu I^{\beta-1}u(\eta_2) + \sum_{j=1}^n \sigma_j u(\zeta_j), \end{cases} \quad (2)$$

where D^χ is the Riemann-Liouville fractional derivative of order $\chi \in \{\alpha, \beta\}$, $0 < \eta_1 < \eta_2 < \xi_1 < \xi_2 < \dots < \xi_m < \zeta_1 < \zeta_2 < \dots < \zeta_n < T$, $a_0, a_1, b_0, b_1, \nu, \mu, \lambda_1, \lambda_2, \mu_i$ ($i = 1, 2, \dots, m$), σ_j ($j = 1, 2, \dots, n$) are real constants and $a_0 \neq -1, a_1 \neq -1, b_0 \neq -1$, and $F, G : [0, T] \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ are continuous functions.

Here we emphasize that our results are new with respect to the semi-coupled boundary conditions (2) and enrich the related literature on the topic.

In Section 2, we prove an auxiliary lemma dealing with the linear variant of systems (1)–(2), which plays a fundamental role in establishing the existence and uniqueness results for the given nonlinear problem, presented in Section 3. Illustrative examples demonstrating the application of the obtained results are given in Section 4.

2 Preliminaries

Let us begin this section with some related definitions [30,31].

Definition 2.1. The (left) Riemann-Liouville fractional integral of order $\chi > 0$ for $\varrho \in L_1[a, b]$, $-\infty < a \leq t \leq b < +\infty$, existing almost everywhere on $[a, b]$, is defined as

$$(I_{a+}^\chi \varrho)(t) = \frac{1}{\Gamma(\chi)} \int_a^t (t-s)^{\chi-1} \varrho(s) ds, \quad \chi > 0,$$

where $\Gamma(\cdot)$ is the gamma function and $(I_{a+}^0 \varrho)(x) = \varrho(x)$.

Definition 2.2. For $\varrho, \varrho^{(m)} \in L^1[a, b]$, the (left) Riemann-Liouville fractional derivative $D_{a+}^\chi \varrho$ of order $\chi \in (m-1, m]$, $m \in \mathbb{N}$ is defined as

$$D_{a+}^\chi \varrho(t) = \frac{1}{\Gamma(m-\chi)} \frac{d^m}{dt^m} \int_a^t (t-s)^{m-1-\chi} \varrho(s) ds, \quad -\infty \leq a < t < b \leq +\infty.$$

In the sequel, we will write Riemann-Liouville fractional integral and derivative operators as I^χ and D^χ instead of I_{a+}^χ and D_{a+}^χ , respectively.

Lemma 2.1. Let $\psi_1, \psi_2 \in C[0, T] \cap L[0, T]$ and $\Lambda \neq 0$. Then the unique solution of the following linear system

$$\begin{cases} D^\alpha u(t) = \psi_1(t), & 1 < \alpha \leq 2, \quad t \in [0, T], \\ D^\beta v(t) = \psi_2(t), & 1 < \beta \leq 2, \quad t \in [0, T], \end{cases} \quad (3)$$

subject to the boundary conditions (2) is given by

$$\begin{aligned} u(t) = & \lambda_1 \rho_1(t) + \lambda_2 \omega_1 \rho_2(t) - a_1 \rho_2(t) I^1 \psi_1(T) - b_1 \rho_3(t) I^1 \psi_2(T) - a_0 \rho_1(t) I^2 \psi_1(T) - b_0 \omega_1 \rho_2(t) I^2 \psi_2(T) \\ & + \mu \rho_3(t) I^{\alpha+\beta-1} \psi_1(\eta_2) + \nu \rho_2(t) I^{\alpha+\beta-1} \psi_2(\eta_1) + \rho_3(t) \sum_{j=1}^n \sigma_j I^\alpha \psi_1(\zeta_j) + \rho_2(t) \sum_{i=1}^m \mu_i I^\beta \psi_2(\xi_i) + I^\alpha \psi_1(t), \end{aligned} \quad (4)$$

$$\begin{aligned}
v(t) = & \lambda_2 \rho_1^*(t) + \lambda_1 v_1 \rho_2^*(t) - a_1 \rho_3^*(t) I^1 \psi_1(T) - b_1 \rho_2^*(t) I^1 \psi_2(T) - a_0 v_1 \rho_2^*(t) I^2 \psi_1(T) - b_0 \rho_1^*(t) I^2 \psi_2(T) \\
& + \mu \rho_2^*(t) I^{\alpha+\beta-1} \psi_1(\eta_2) + v \rho_3^*(t) I^{\alpha+\beta-1} \psi_2(\eta_1) + \rho_2^*(t) \sum_{j=1}^n \sigma_j I^\alpha \psi_1(\zeta_j) + \rho_3^*(t) \sum_{i=1}^m \mu_i I^\beta \psi_2(\xi_i) + I^\beta \psi_2(t),
\end{aligned} \quad (5)$$

where

$$\begin{aligned}
\rho_1(t) &= t^{\alpha-2} \left[\frac{1}{(1+a_0)\Gamma(\alpha-1)} + \frac{v_1}{\Lambda} \frac{a_0 T \Gamma(\alpha)}{(1+a_0)\Gamma(\alpha-1)} \left(\omega_2 - \frac{b_0 T \Gamma(\beta)}{(1+a_1)\Gamma(\alpha)} \omega_1 \right) \right] - t^{\alpha-1} \frac{v_1}{\Lambda} \left[\omega_2 - \frac{b_0 T \Gamma(\beta)}{(1+a_1)\Gamma(\alpha)} \omega_1 \right], \\
\rho_2(t) &= \frac{1}{\Lambda} \left[t^{\alpha-2} \frac{a_0 T (b_1+1) \Gamma(\beta)}{(1+a_1)(1+a_0)\Gamma(\alpha-1)} - t^{\alpha-1} \frac{(1+b_1)\Gamma(\beta)}{(1+a_1)\Gamma(\alpha)} \right], \\
\rho_3(t) &= \frac{1}{\Lambda} \left[\omega_2 - \frac{b_0 T \Gamma(\beta)}{(1+a_1)\Gamma(\alpha)} \omega_1 \right] \left[t^{\alpha-2} \frac{a_0 T \Gamma(\alpha)}{(1+a_0)\Gamma(\alpha-1)} - t^{\alpha-1} \right], \\
\rho_1^*(t) &= t^{\beta-2} \left[\frac{1}{(1+b_0)\Gamma(\beta-1)} + \frac{\omega_1}{\Lambda} \frac{b_0 T \Gamma(\beta)}{(1+b_0)\Gamma(\beta-1)} \left(v_2 - \frac{a_0 T}{(1+a_1)} v_1 \right) \right] - t^{\beta-1} \frac{\omega_1}{\Lambda} \left[v_2 - \frac{a_0 T}{(1+a_1)} v_1 \right], \\
\rho_2^*(t) &= \frac{1}{\Lambda} \left[t^{\beta-2} \frac{b_0 T \Gamma(\beta)}{(1+b_0)\Gamma(\beta-1)} - t^{\beta-1} \right], \\
\rho_3^*(t) &= \frac{1}{\Lambda} \left[v_2 - \frac{a_0 T}{(1+a_1)} v_1 \right] \left[t^{\beta-2} \frac{b_0 T \Gamma(\beta)}{(1+b_0)\Gamma(\beta-1)} - t^{\beta-1} \right], \\
\Lambda &= -\Gamma(\beta)(1+b_1) + \left[v_2 - \frac{a_0 T}{(1+a_1)} v_1 \right] [\Gamma(\alpha)(1+a_1)\omega_2 - b_0 T \Gamma(\beta)\omega_1], \\
v_1 &= \frac{1}{(1+a_0)\Gamma(\alpha-1)} \left[\mu \eta_2^{\alpha+\beta-3} \frac{\Gamma(\alpha-1)}{\Gamma(\alpha+\beta-2)} + \sum_{j=1}^n \sigma_j \zeta_j^{\alpha-2} \right], \\
v_2 &= \frac{1}{(1+a_1)\Gamma(\alpha)} \left[\mu \eta_2^{\alpha+\beta-2} \frac{\Gamma(\alpha)}{\Gamma(\alpha+\beta-1)} + \sum_{j=1}^n \sigma_j \zeta_j^{\alpha-1} \right], \\
\omega_1 &= \frac{1}{(1+b_0)\Gamma(\beta-1)} \left[v \eta_1^{\alpha+\beta-3} \frac{\Gamma(\beta-1)}{\Gamma(\alpha+\beta-2)} + \sum_{i=1}^m \mu_i \xi_i^{\beta-2} \right], \\
\omega_2 &= \frac{1}{(1+a_1)\Gamma(\alpha)} \left[v \eta_1^{\alpha+\beta-2} \frac{\Gamma(\beta)}{\Gamma(\alpha+\beta-1)} + \sum_{i=1}^m \mu_i \xi_i^{\beta-1} \right].
\end{aligned} \quad (6)$$

Proof. It is well known that the solutions of fractional differential equations in (3) can be written as

$$u(t) = I^\alpha \psi_1(t) + c_0 t^{\alpha-2} + c_1 t^{\alpha-1}, \quad (7)$$

$$v(t) = I^\beta \psi_2(t) + c_2 t^{\beta-2} + c_3 t^{\beta-1}, \quad (8)$$

where $c_i \in \mathbb{R}$, $i = 0, 1, 2, 3$ are unknown arbitrary constants. From (7) and (8), we have

$$D^{\alpha-2} u(t) = c_0 \Gamma(\alpha-1) + c_1 \Gamma(\alpha) t + I^2 \psi_1(t), \quad (9)$$

$$D^{\beta-2} v(t) = c_2 \Gamma(\beta-1) + c_3 \Gamma(\beta) t + I^2 \psi_2(t), \quad (10)$$

$$D^{\alpha-1} u(t) = c_1 \Gamma(\alpha) + I^1 \psi_1(t), \quad (11)$$

$$D^{\beta-1} v(t) = c_3 \Gamma(\beta) + I^1 \psi_2(t). \quad (12)$$

Using (9) and (10) in (2), we get

$$(1+a_0)\Gamma(\alpha-1)c_0 + a_0 T \Gamma(\alpha)c_1 = \lambda_1 - a_0 I^2 \psi_1(T), \quad (13)$$

$$(1+b_0)\Gamma(\beta-1)c_2 + b_0 T \Gamma(\beta)c_3 = \lambda_2 - b_0 I^2 \psi_2(T). \quad (14)$$

Combining (11) and (12) with (2), we obtain

$$\begin{aligned} & \Gamma(\alpha)(1+a_1)c_1 - \left[\frac{v\eta_1^{\alpha+\beta-2}\Gamma(\beta)}{\Gamma(\alpha+\beta-1)} + \sum_{i=1}^m \mu_i \xi_i^{\beta-1} - \frac{b_0 T \Gamma(\beta)}{(1+b_0)\Gamma(\beta-1)} \left(v\eta_1^{\alpha+\beta-3} \frac{\Gamma(\beta-1)}{\Gamma(\alpha+\beta-2)} + \sum_{i=1}^m \mu_i \xi_i^{\beta-2} \right) \right] c_3 \\ &= - \frac{b_0}{(1+b_0)\Gamma(\beta-1)} \left[v\eta_1^{\alpha+\beta-3} \frac{\Gamma(\beta-1)}{\Gamma(\alpha+\beta-2)} + \sum_{i=1}^m \mu_i \xi_i^{\beta-2} \right] I^2 \psi_2(t) + \sum_{i=1}^m \mu_i I^\beta \psi_2(\xi_i) - a_1 I^1 \psi_1(T) \\ &+ \frac{\lambda_2}{(1+b_0)\Gamma(\beta-1)} \left[v\eta_1^{\alpha+\beta-3} \frac{\Gamma(\beta-1)}{\Gamma(\alpha+\beta-2)} + \sum_{i=1}^m \mu_i \xi_i^{\beta-2} \right] + v I^{\alpha+\beta-1} \psi_2(\eta_1), \end{aligned} \quad (15)$$

$$\begin{aligned} & - \left[\mu \eta_2^{\alpha+\beta-2} \frac{\Gamma(\alpha)}{\Gamma(\alpha+\beta-1)} + \sum_{j=1}^n \sigma_j \zeta_j^{\alpha-1} \right] c_1 + \Gamma(\beta)(1+b_1)c_3 \\ &= \sum_{j=1}^n \sigma_j I^\alpha \psi_1(\zeta_j) - b_1 I^1 \psi_2(T) + \frac{1}{(1+a_0)\Gamma(\alpha-1)} \left[\mu \eta_2^{\alpha+\beta-3} \frac{\Gamma(\alpha-1)}{\Gamma(\alpha+\beta-2)} + \sum_{j=1}^n \sigma_j \zeta_j^{\alpha-2} \right] \\ & \times [\lambda_1 - a_0 c_1 T \Gamma(\alpha) - a_0 I^2 \psi_1(t)] + \mu I^{\alpha+\beta-1} \psi_1(\eta_2). \end{aligned} \quad (16)$$

Solving (15) and (16) for c_1 and c_3 , we find that

$$\begin{aligned} c_1 &= \left[\frac{1}{\Lambda} \frac{(b_1+1)\Gamma(\beta)}{(1+a_1)\Gamma(\alpha)} \right] \left[-\omega_1 \lambda_2 + b_0 \omega_1 I^2 \psi_2(T) - v I^{\alpha+\beta-1} \psi_2(\eta_1) - \sum_{i=1}^m \mu_i I^\beta \psi_2(\xi_i) + a_1 I^1 \psi_1(T) \right] \\ &+ \frac{1}{\Lambda} \left[\omega_2 - \frac{b_0 T \Gamma(\beta)}{(1+a_1)\Gamma(\alpha)} \omega_1 \right] \left[-\lambda_1 v_1 - \mu I^{\alpha+\beta-1} \psi_1(\eta_2) - \sum_{j=1}^n \sigma_j I^\alpha \psi_1(\zeta_j) + b_1 I^1 \psi_2(T) + a_0 v_1 I^2 \psi_1(T) \right], \end{aligned} \quad (17)$$

$$\begin{aligned} c_3 &= \frac{1}{\Lambda} \left[-\lambda_1 v_1 - \lambda_2 \omega_1 \left(v_2 - \frac{a_0 T}{(1+a_1)} v_1 \right) + a_0 v_1 I^2 \psi_1(t) + b_0 \omega_1 \left(v_2 - \frac{a_0 T}{(1+a_1)} v_1 \right) I^2 \psi_2(T) \right. \\ &- v \left(v_2 - \frac{a_0 T}{(1+a_1)} v_1 \right) I^{\alpha+\beta-1} \psi_2(\eta_1) - \mu I^{\alpha+\beta-1} \psi_1(\eta_2) - \left(v_2 - \frac{a_0 T}{(1+a_1)} v_1 \right) \sum_{i=1}^m \mu_i I^\beta \psi_2(\xi_i) \\ &\left. - \sum_{j=1}^n \sigma_j I^\alpha \psi_1(\zeta_j) + a_1 \left(v_2 - \frac{a_0 T}{(1+a_1)} v_1 \right) I^1 \psi_1(T) + b_1 I^1 \psi_2(T) \right]. \end{aligned} \quad (18)$$

Substituting the values of c_1 and c_3 in (13) and (14), respectively, we get

$$\begin{aligned} c_0 &= [\lambda_1 - a_0 I^2 \psi_1(T)] \left[\frac{1}{(1+a_0)\Gamma(\alpha-1)} + \frac{v_1}{\Lambda} \frac{a_0 T \Gamma(\alpha)}{(1+a_0)\Gamma(\alpha-1)} \left(\omega_2 - \frac{b_0 T \Gamma(\beta)}{(1+a_1)\Gamma(\alpha)} \omega_1 \right) \right] \\ &+ \lambda_2 \left[\frac{\omega_1}{\Lambda} \frac{a_0(1+b_1)T \Gamma(\beta)}{(1+a_0)(1+a_1)\Gamma(\alpha-1)} \right] - I^2 \psi_2(T) \left[\frac{\omega_1}{\Lambda} \frac{a_0 b_0(1+b_1)T \Gamma(\beta)}{(1+a_0)(1+a_1)\Gamma(\alpha-1)} \right] \\ &+ I^{\alpha+\beta-1} \psi_2(\eta_1) \left[\frac{v}{\Lambda} \frac{a_0(1+b_1)T \Gamma(\beta)}{(1+a_0)(1+a_1)\Gamma(\alpha-1)} \right] + \sum_{i=1}^m \mu_i I^\beta \psi_2(\xi_i) \left[\frac{1}{\Lambda} \frac{a_0(1+b_1)T \Gamma(\beta)}{(1+a_0)(1+a_1)\Gamma(\alpha-1)} \right] \\ &- I^1 \psi_1(T) \left[\frac{1}{\Lambda} \frac{a_0 a_1(1+b_1)T \Gamma(\beta)}{(1+a_0)(1+a_1)\Gamma(\alpha-1)} \right] + I^{\alpha+\beta-1} \psi_1(\eta_2) \left[\frac{\mu}{\Lambda} \frac{a_0 T \Gamma(\alpha)}{(1+a_0)\Gamma(\alpha-1)} \left(\omega_2 - \frac{b_0 T \Gamma(\beta)}{(1+a_1)\Gamma(\alpha)} \omega_1 \right) \right] \\ &+ \sum_{j=1}^n \sigma_j I^\alpha \psi_1(\zeta_j) \left[\frac{1}{\Lambda} \frac{a_0 T \Gamma(\alpha)}{(1+a_0)\Gamma(\alpha-1)} \left(\omega_2 - \frac{b_0 T \Gamma(\beta)}{(1+a_1)\Gamma(\alpha)} \omega_1 \right) \right] \\ &+ I^1 \psi_2(T) \left[\frac{-b_1}{\Lambda} \frac{a_0 T \Gamma(\alpha)}{(1+a_0)\Gamma(\alpha-1)} \left(\omega_2 - \frac{b_0 T \Gamma(\beta)}{(1+a_1)\Gamma(\alpha)} \omega_1 \right) \right], \end{aligned}$$

$$\begin{aligned}
c_2 = & [\lambda_2 - b_0 I^2 \psi_2(T)] \left[\frac{1}{(1+b_0)\Gamma(\beta-1)} + \frac{\omega_1}{\Lambda} \frac{b_0 T \Gamma(\beta)}{(1+b_0)\Gamma(\beta-1)} \left(v_2 - \frac{a_0 T}{(1+b_1)} v_1 \right) \right] \\
& + \lambda_1 \left[\frac{v_1}{\Lambda} \frac{b_0 T \Gamma(\beta)}{(1+b_0)\Gamma(\beta-1)} \right] - I^2 \psi_1(T) \left[\frac{v_1}{\Lambda} \frac{a_0 b_0 T \Gamma(\beta)}{(1+b_0)\Gamma(\beta-1)} \right] \\
& + I^{\alpha+\beta-1} \psi_2(\eta_1) \left[\frac{v}{\Lambda} \frac{b_0 T \Gamma(\beta)}{(1+b_0)\Gamma(\beta-1)} \left(v_2 - \frac{a_0 T}{(1+a_1)} v_1 \right) \right] \\
& + I^{\alpha+\beta-1} \psi_1(\eta_2) \left[\frac{\mu}{\Lambda} \frac{b_0 T \Gamma(\beta)}{(1+b_0)\Gamma(\beta-1)} \right] + \sum_{i=1}^m \mu_i I^\beta \psi_2(\xi_i) \left[\frac{1}{\Lambda} \frac{b_0 T \Gamma(\beta)}{(1+b_0)\Gamma(\beta-1)} \left(v_2 - \frac{a_0 T}{(1+a_1)} v_1 \right) \right] \\
& - I^1 \psi_1(T) \left[\frac{1}{\Lambda} \frac{b_0 a_1 T \Gamma(\beta)}{(1+b_0)\Gamma(\beta-1)} \left(v_2 - \frac{a_0 T}{(1+a_1)} v_1 \right) \right] \\
& + \sum_{j=1}^n \sigma_j I^\alpha \psi_1(\zeta_j) \left[\frac{1}{\Lambda} \frac{b_0 T \Gamma(\beta)}{(1+b_0)\Gamma(\beta-1)} \right] - I^1 \psi_2(T) \left[\frac{1}{\Lambda} \frac{b_0 b_1 T \Gamma(\beta)}{(1+b_0)\Gamma(\beta-1)} \right].
\end{aligned}$$

Inserting the value of c_0 , c_1 , c_2 , and c_3 in (7) and (8) together with the notations (6) leads to the solutions (4) and (5). The converse of this lemma follows by direct computation. The proof is completed. $\square$

3 Main results

Let $C([0, T], \mathbb{R})$ denote the Banach space of all continuous real-valued functions defined on $[0, T]$ with norm $\|u\| = \sup_{t \in [0, T]} |u(t)|$. For $t \in [0, T]$, let $C_r([0, T], \mathbb{R})$ denote the space of all functions u_r such that $u_r \in C([0, T], \mathbb{R})$, which is a Banach space endowed with norm $\|u\|_r = \sup_{t \in [0, T]} \{t^r |u(t)|\}$. Let $X = \{u : u \in C_{2-\alpha}([0, T], \mathbb{R})\}$ and $Y = \{v : v \in C_{2-\beta}([0, T], \mathbb{R})\}$ be equipped with the norm $\|u\|_X = \sup_{t \in [0, T]} \{t^{2-\alpha} |u(t)|\}$ and $\|v\|_Y = \sup_{t \in [0, T]} \{t^{2-\beta} |v(t)|\}$, respectively. Then the product space $(X \times Y, \|\cdot\|_{X \times Y})$ is a Banach space with the norm

$$\|(u, v)\|_{X \times Y} = \|u\|_X + \|v\|_Y.$$

Next we introduce an operator $\mathcal{P} : X \times Y \rightarrow X \times Y$ by

$$\mathcal{P}(u, v)(t) = (\mathcal{P}_1(u, v)(t), \mathcal{P}_2(u, v)(t)),$$

where

$$\begin{aligned}
\mathcal{P}_1(u, v)(t) = & \lambda_1 \rho_1(t) + \lambda_2 \omega_1 \rho_2(t) - a_1 \rho_2(t) I^1 F(T, u(T), v(T)) - b_1 \rho_3(t) I^1 G(T, u(T), v(T)) \\
& - a_0 \rho_1(t) I^2 F(T, u(T), v(T)) - b_0 \omega_1 \rho_2(t) I^2 G(T, u(T), v(T)) \\
& + \mu \rho_3(t) I^{\alpha+\beta-1} F(\eta_2, u(\eta_2), v(\eta_2)) + \nu \rho_2(t) I^{\alpha+\beta-1} G(\eta_1, u(\eta_1), v(\eta_1)) \\
& + \rho_3(t) \sum_{j=1}^n \sigma_j I^\alpha F(\zeta_j, u(\zeta_j), v(\zeta_j)) + \rho_2(t) \sum_{i=1}^m \mu_i I^\beta G(\xi_i, u(\xi_i), v(\xi_i)) \\
& + I^\alpha F(t, u(t), v(t)), \quad t \in [0, T], \\
\mathcal{P}_2(u, v)(t) = & \lambda_2 \rho_1^*(t) + \lambda_1 \nu \rho_2^*(t) - a_1 \rho_3^*(t) I^1 F(T, u(T), v(T)) - b_1 \rho_2^*(t) I^1 G(T, u(T), v(T)) \\
& - a_0 \nu \rho_2^*(t) I^2 F(T, u(T), v(T)) - b_0 \rho_1^*(t) I^2 G(T, u(T), v(T)) \\
& + \mu \rho_2^*(t) I^{\alpha+\beta-1} F(\eta_2, u(\eta_2), v(\eta_2)) + \nu \rho_3^*(t) I^{\alpha+\beta-1} G(\eta_1, u(\eta_1), v(\eta_1)) \\
& + \rho_2^*(t) \sum_{j=1}^n \sigma_j I^\alpha F(\zeta_j, u(\zeta_j), v(\zeta_j)) + \rho_3^*(t) \sum_{i=1}^m \mu_i I^\beta G(\xi_i, u(\xi_i), v(\xi_i)) \\
& + I^\beta G(t, u(t), v(t)), \quad t \in [0, T].
\end{aligned} \tag{19}$$

For the sake of brevity, we set

$$\begin{aligned} N_1 = & |\mu|\delta_3 \left[\frac{\eta_2^{2\alpha+\beta-3}\Gamma(\alpha-1)}{\Gamma(2\alpha+\beta-2)} + \frac{\eta_2^{\alpha+2\beta-3}\Gamma(\beta-1)}{\Gamma(\alpha+2\beta-2)} \right] + |a_0|\delta_1 \left[\frac{T^\alpha\Gamma(\alpha-1)}{\Gamma(\alpha+1)} + \frac{T^\beta\Gamma(\beta-1)}{\Gamma(\beta+1)} \right] \\ & + \delta_3 \sum_{j=1}^n |\sigma_j| \left[\frac{\zeta_j^{2\alpha-2}\Gamma(\alpha-1)}{\Gamma(2\alpha-1)} + \frac{\zeta_j^{\alpha+\beta-2}\Gamma(\beta-1)}{\Gamma(\alpha+\beta-1)} \right] + |a_1|\delta_2 \left[\frac{T^{\alpha-1}}{(\alpha-1)} + \frac{T^{\beta-1}}{(\beta-1)} \right] \\ & + \frac{T^\alpha\Gamma(\alpha-1)}{\Gamma(2\alpha-1)} + \frac{T^\beta\Gamma(\beta-1)}{\Gamma(\alpha+\beta-1)}, \end{aligned} \quad (20)$$

$$\begin{aligned} N_2 = & |\nu|\delta_2 \left[\frac{\eta_1^{2\alpha+\beta-3}\Gamma(\alpha-1)}{\Gamma(2\alpha+\beta-2)} + \frac{\eta_1^{\alpha+2\beta-3}\Gamma(\beta-1)}{\Gamma(\alpha+2\beta-2)} \right] + |b_0|\omega_1|\delta_2 \left[\frac{T^\alpha\Gamma(\alpha-1)}{\Gamma(\alpha+1)} + \frac{T^\beta\Gamma(\beta-1)}{\Gamma(\beta+1)} \right] \\ & + \delta_2 \sum_{i=1}^m |\mu_i| \left[\frac{\xi_i^{\alpha+\beta-2}\Gamma(\alpha-1)}{\Gamma(\alpha+\beta-1)} + \frac{\xi_i^{2\beta-2}\Gamma(\beta-1)}{\Gamma(2\beta-1)} \right] + |b_1|\delta_3 \left[\frac{T^{\alpha-1}}{\alpha-1} + \frac{T^{\beta-1}}{\beta-1} \right], \end{aligned} \quad (21)$$

$$\begin{aligned} N_1^* = & |\mu|\delta_2^* \left[\frac{\eta_2^{2\alpha+\beta-3}\Gamma(\alpha-1)}{\Gamma(2\alpha+\beta-2)} + \frac{\eta_2^{\alpha+2\beta-3}\Gamma(\beta-1)}{\Gamma(\alpha+2\beta-2)} \right] + |a_0\nu_1|\delta_2^* \left[\frac{T^\alpha\Gamma(\alpha-1)}{\Gamma(\alpha+1)} + \frac{T^\beta\Gamma(\beta-1)}{\Gamma(\beta+1)} \right] \\ & + \delta_2^* \sum_{j=1}^n |\sigma_j| \left[\frac{\zeta_j^{2\alpha-2}\Gamma(\alpha-1)}{\Gamma(2\alpha-1)} + \frac{\zeta_j^{\alpha+\beta-2}\Gamma(\beta-1)}{\Gamma(\alpha+\beta-1)} \right] + |a_1|\delta_3^* \left[\frac{T^{\alpha-1}}{(\alpha-1)} + \frac{T^{\beta-1}}{(\beta-1)} \right], \end{aligned} \quad (22)$$

$$\begin{aligned} N_2^* = & |\nu|\delta_3^* \left[\frac{\eta_1^{2\alpha+\beta-3}\Gamma(\alpha-1)}{\Gamma(2\alpha+\beta-2)} + \frac{\eta_1^{\alpha+2\beta-3}\Gamma(\beta-1)}{\Gamma(\alpha+2\beta-2)} \right] + |b_0|\delta_1^* \left[\frac{T^\alpha\Gamma(\alpha-1)}{\Gamma(\alpha+1)} + \frac{T^\beta\Gamma(\beta-1)}{\Gamma(\beta+1)} \right] \\ & + \delta_3^* \sum_{i=1}^m |\mu_i| \left[\frac{\xi_i^{\alpha+\beta-2}\Gamma(\alpha-1)}{\Gamma(\alpha+\beta-1)} + \frac{\xi_i^{2\beta-2}\Gamma(\beta-1)}{\Gamma(2\beta-1)} \right] + |b_1|\delta_2^* \left[\frac{T^{\alpha-1}}{\alpha-1} + \frac{T^{\beta-1}}{\beta-1} \right] \\ & + \frac{T^\alpha\Gamma(\alpha-1)}{\Gamma(\alpha+\beta-1)} + \frac{T^\beta\Gamma(\beta-1)}{\Gamma(2\beta-1)}. \end{aligned} \quad (23)$$

Theorem 3.1. Assume that:

(A1) $F, G : [0, T] \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ are continuous functions and there exist positive constants L_1 and L_2 such that, for all $t \in [0, T]$ and $u_i, v_i \in \mathbb{R}, i = 1, 2$,

$$|F(t, u_1, v_1) - F(t, u_2, v_2)| \leq L_1(|u_1 - u_2| + |v_1 - v_2|),$$

$$|G(t, u_1, v_1) - G(t, u_2, v_2)| \leq L_2(|u_1 - u_2| + |v_1 - v_2|).$$

Then the system (1)–(2) has a unique solution on $[0, T]$, provided that

$$L_1(N_1 + N_1^*) + L_2(N_2 + N_2^*) < 1, \quad (24)$$

where, N_1, N_2, N_1^* , and N_2^* are, respectively, given by (20), (21), (22), and (23).

Proof. Define $\sup_{t \in [0, T]} F(t, 0, 0) = M_1$, $\sup_{t \in [0, T]} G(t, 0, 0) = M_2$, and choose $r > 0$ such that

$$r \geq \frac{|\lambda_1|(\delta_1 + \delta_2^*|\nu_1|) + |\lambda_2|(|\omega_1|\delta_2 + \delta_1^*) + M_1(e_1 + e_1^*) + M_2(e_2 + e_2^*)}{1 - [L_1(N_1 + N_1^*) + L_2(N_2 + N_2^*)]}, \quad (25)$$

where

$$\begin{aligned} e_1 = & |a_0|\delta_1 \frac{T^2}{2} + |a_1|\delta_2 T + |\mu|\delta_3 \frac{\eta_2^{\alpha+\beta-1}}{\Gamma(\alpha+\beta)} + \delta_3 \sum_{j=1}^n |\sigma_j| \frac{\zeta_j^\alpha}{\Gamma(\alpha+1)} + \frac{T^2}{\Gamma(\alpha+1)}, \\ e_2 = & |b_1|\delta_3 T + |\nu|\delta_2 \frac{\eta_1^{\alpha+\beta-1}}{\Gamma(\alpha+\beta)} + \delta_2 \sum_{i=1}^m |\mu_i| \frac{\xi_i^\beta}{\Gamma(\beta+1)} + |b_0\omega_1|\delta_2 \frac{T^2}{2}, \end{aligned} \quad (26)$$

$$\begin{aligned}
e_1^* &= |a_0 v_1| \delta_2^* \frac{T^2}{2} + |a_1| \delta_3^* T + |\mu| \delta_2^* \frac{\eta_2^{\alpha+\beta-1}}{\Gamma(\alpha+\beta)} + \delta_2^* \sum_{j=1}^n |\sigma_j| \frac{\zeta_j^\alpha}{\Gamma(\alpha+1)}, \\
e_2^* &= |b_1| \delta_2^* T + |v| \delta_3^* \frac{\eta_1^{\alpha+\beta-1}}{\Gamma(\alpha+\beta)} + \delta_3^* \sum_{i=1}^m |\mu_i| \frac{\xi_i^\beta}{\Gamma(\beta+1)} + |b_0| \delta_1^* \frac{T^2}{2} + \frac{T^2}{\Gamma(\beta+1)}, \\
\delta_m &= \sup_{t \in [0, T]} \{t^{2-\alpha} |\rho_m(t)|\}, \quad \delta_m^* = \sup_{t \in [0, T]} \{t^{2-\beta} |\rho_m^*(t)|\}, \quad m = 1, 2, 3.
\end{aligned} \tag{27}$$

In the first step, it will be shown that $\mathcal{P}\mathbb{B}_r \subset \mathbb{B}_r$, where $\mathbb{B}_r = \{(u, v) \in X \times Y : \|(u, v)\|_{X \times Y} \leq r\}$. By the assumption (A_1) , for $(u, v) \in \mathbb{B}_r$, $t \in [0, T]$, we have

$$|F(t, u(t), v(t))| \leq |F(t, u(t), v(t)) - F(t, 0, 0)| + |F(t, 0, 0)| \leq L_1(|u| + |v|) + M_1$$

and

$$|G(t, u(t), v(t))| \leq |G(t, u(t), v(t)) - G(t, 0, 0)| + |G(t, 0, 0)| \leq L_2(|u| + |v|) + M_2.$$

In consequence, by using the relation for beta function $B(\cdot, \cdot)$:

$$B(a, b) = \int_0^1 u^{a-1} (1-u)^{b-1} du = \frac{\Gamma(a)\Gamma(b)}{\Gamma(a+b)},$$

we obtain

$$\begin{aligned}
\|\mathcal{P}_1(u, v)\|_X &= \sup_{t \in [0, T]} \{t^{2-\alpha} |\mathcal{P}_1(u, v)(t)|\} \\
&\leq \sup_{t \in [0, T]} \left\{ t^{2-\alpha} \left[|\lambda_1| |\rho_1(t)| + |\lambda_2 \omega_1| |\rho_2(t)| + |a_0| |\rho_1(t)| \int_0^T \frac{(T-s)}{\Gamma(2)} (L_1(|u| + |v|) + M_1) ds \right. \right. \\
&\quad + |a_1| |\rho_2(t)| \int_0^T (L_1(|u| + |v|) + M_1) ds + |b_1| |\rho_3(t)| \int_0^T (L_2(|u| + |v|) + M_2) ds \\
&\quad + |v| |\rho_2(t)| \int_0^{\eta_1} \frac{(\eta_1 - s)^{\alpha+\beta-2}}{\Gamma(\alpha+\beta-1)} (L_2(|u| + |v|) + M_2) ds + |\mu| |\rho_3(t)| \int_0^{\eta_2} \frac{(\eta_2 - s)^{\alpha+\beta-2}}{\Gamma(\alpha+\beta-1)} (L_1(|u| + |v|) + M_1) ds \\
&\quad + |\rho_2(t)| \sum_{i=1}^m |\mu_i| \int_0^{\xi_i} \frac{(\xi_i - s)^{\beta-1}}{\Gamma(\beta)} (L_2(|u| + |v|) + M_2) ds + |\rho_3(t)| \sum_{j=1}^n |\sigma_j| \int_0^{\zeta_j} \frac{(\zeta_j - s)^{\alpha-1}}{\Gamma(\alpha)} (L_1(|u| + |v|) + M_1) ds \\
&\quad \left. \left. + |b_0 \omega_1| |\rho_2(t)| \int_0^T \frac{(T-s)}{\Gamma(2)} (L_2(|u| + |v|) + M_2) ds + \int_0^t \frac{(t-s)^{\alpha-1}}{\Gamma(\alpha)} (L_1(|u| + |v|) + M_1) ds \right] \right\} \\
&\leq |\lambda_1| \delta_1 + |\lambda_2 \omega_1| \delta_2 + |a_0| \delta_1 L_1 \int_0^T (T-s) [s^{\alpha-2} \|u\|_X + s^{\beta-2} \|v\|_Y] ds + M_1 |a_0| \delta_1 \frac{T^2}{2} \\
&\quad + |a_1| \delta_2 L_1 \int_0^T [s^{\alpha-2} \|u\|_X + s^{\beta-2} \|v\|_Y] ds + |a_1| \delta_2 M_1 T + |b_1| \delta_3 L_2 \int_0^T [s^{\alpha-2} \|u\|_X + s^{\beta-2} \|v\|_Y] ds \\
&\quad + M_2 |b_1| \delta_3 T + |v| \delta_2 L_2 \int_0^{\eta_1} \frac{(\eta_1 - s)^{\alpha+\beta-2}}{\Gamma(\alpha+\beta-1)} [s^{\alpha-2} \|u\|_X + s^{\beta-2} \|v\|_Y] ds \\
&\quad + M_2 |v| \delta_2 \frac{(\eta_1)^{\alpha+\beta-1}}{\Gamma(\alpha+\beta)} + |\mu| \delta_3 L_1 \int_0^{\eta_2} \frac{(\eta_2 - s)^{\alpha+\beta-2}}{\Gamma(\alpha+\beta-1)} [s^{\alpha-2} \|u\|_X + s^{\beta-2} \|v\|_Y] ds \\
&\quad + M_1 |\mu| \delta_3 \frac{(\eta_2)^{\alpha+\beta-1}}{\Gamma(\alpha+\beta)} + \delta_2 \sum_{i=1}^m |\mu_i| L_2 \int_0^{\xi_i} \frac{(\xi_i - s)^{\beta-1}}{\Gamma(\beta)} [s^{\alpha-2} \|u\|_X + s^{\beta-2} \|v\|_Y] ds
\end{aligned}$$

$$\begin{aligned}
& + M_2 \delta_2 \sum_{i=1}^m \frac{|\mu_i| \zeta_i^\beta}{\Gamma(\beta+1)} + \delta_3 \sum_{j=1}^n |\sigma_j| L_1 \int_0^{\zeta_j} \frac{(\zeta_j - s)^{\alpha-1}}{\Gamma(\alpha)} [s^{\alpha-2} \|u\|_X + s^{\beta-2} \|v\|_Y] ds \\
& + M_1 \delta_3 \sum_{j=1}^n \frac{|\sigma_j| \zeta_j^\alpha}{\Gamma(\alpha+1)} + |b_0 \omega_1| \delta_2 L_2 \int_0^T (T-s) [s^{\alpha-2} \|u\|_X + s^{\beta-2} \|v\|_Y] ds \\
& + M_1 |b_0 \omega_1| \delta_2 \frac{T^2}{2} + L_1 t^{2-\alpha} \int_0^t \frac{(t-s)^{\alpha-1}}{\Gamma(\alpha)} [s^{\alpha-2} \|u\|_X + s^{\beta-2} \|v\|_Y] ds + M_1 \frac{T^2}{\Gamma(\alpha+1)} \\
& \leq |\lambda_1| \delta_1 + |\lambda_2 \omega_1| \delta_2 + [L_1 N_1 + L_2 N_2] r + M_1 e_1 + M_2 e_2.
\end{aligned} \tag{28}$$

Similarly, one can get

$$\|\mathcal{P}_2(u, v)\|_Y \leq |\lambda_2| \delta_1^* + |\lambda_1 v_1| \delta_2^* + [L_1 N_1^* + L_2 N_2^*] r + M_1 e_1^* + M_2 e_2^*. \tag{29}$$

In view of (28) and (29) together with (25), we have

$$\|\mathcal{P}(u, v)\|_{X \times Y} \leq |\lambda_1| [\delta_1 + |v_1| \delta_2^*] + |\lambda_2| [\delta_2 | \omega_1| + \delta_1^*] + [L_1 (N_1 + N_1^*) + L_2 (N_2 + N_2^*)] r + M_1 (e_1 + e_1^*) + M_2 (e_2 + e_2^*) \leq r.$$

Now, for $(u_1, v_1), (u_2, v_2) \in X \times Y$, and for any $t \in [0, T]$, we get

$$\begin{aligned}
& t^{2-\alpha} |\mathcal{P}_1(u_2, v_2)(t) - \mathcal{P}_1(u_1, v_1)(t)| \\
& \leq \sup_{t \in [0, T]} \left\{ t^{2-\alpha} \left[|a_0 \rho_1(t)| \int_0^T (T-s) |F(s, u_2, v_2) - F(s, u_1, v_1)| ds \right. \right. \\
& \quad + |a_1 \rho_2(t)| \int_0^T |F(s, u_2, v_2) - F(s, u_1, v_1)| ds + |b_1 \rho_3(t)| \int_0^T |G(s, u_2, v_2) - G(s, u_1, v_1)| ds \\
& \quad + |v \rho_2(t)| \int_0^{\eta_1} \frac{(\eta_1 - s)^{\alpha+\beta-2}}{\Gamma(\alpha+\beta-1)} |G(s, u_2, v_2) - G(s, u_1, v_1)| ds \\
& \quad + |\mu \rho_3(t)| \int_0^{\eta_2} \frac{(\eta_2 - s)^{\alpha+\beta-2}}{\Gamma(\alpha+\beta-1)} |F(s, u_2, v_2) - F(s, u_1, v_1)| ds \\
& \quad + |\rho_2(t)| \sum_{i=1}^m |\mu_i| \int_0^{\xi_i} \frac{(\xi_i - s)^{\beta-1}}{\Gamma(\beta)} |G(s, u_2, v_2) - G(s, u_1, v_1)| ds \\
& \quad + |\rho_3(t)| \sum_{j=1}^n |\sigma_j| \int_0^{\zeta_j} \frac{(\zeta_j - s)^{\alpha-1}}{\Gamma(\alpha)} |F(s, u_2, v_2) - F(s, u_1, v_1)| ds \\
& \quad \left. + |b_0 \omega_1 \rho_2(t)| \int_0^T (T-s) |G(s, u_2, v_2) - G(s, u_1, v_1)| ds + \int_0^t \frac{(t-s)^{\alpha-1}}{\Gamma(\alpha)} |F(s, u_2, v_2) - F(s, u_1, v_1)| ds \right] \Bigg\}.
\end{aligned} \tag{30}$$

Using (A_1) and the relation $|u_2 - u_1| + |v_2 - v_1| \leq t^{\alpha-2} \|u_2 - u_1\|_X + t^{\beta-2} \|v_2 - v_1\|_Y$ in (30) yields

$$\|\mathcal{P}_1(u_2, v_2) - \mathcal{P}_1(u_1, v_1)\|_X \leq (L_1 N_1 + L_2 N_2) [\|u_2 - u_1\|_X + \|v_2 - v_1\|_Y]. \tag{31}$$

In a similar manner, one can get

$$\|\mathcal{P}_2(u_2, v_2) - \mathcal{P}_2(u_1, v_1)\|_Y \leq (L_1 N_1^* + L_2 N_2^*) [\|u_2 - u_1\|_X + \|v_2 - v_1\|_Y]. \tag{32}$$

Thus, it follows from (31) and (32) that

$$\|\mathcal{P}(u_2, v_2) - \mathcal{P}(u_1, v_1)\|_{X \times Y} \leq [L_1 (N_1 + N_1^*) + L_2 (N_2 + N_2^*)] [\|u_2 - u_1\|_X + \|v_2 - v_1\|_Y],$$

which, in view of condition (24), implies that $\mathcal{P}$ is a contraction. Hence, by Banach's fixed point theorem, the operator $\mathcal{P}$ has a unique fixed point, which is indeed a unique solution of the problem (1)–(2) on $[0, T]$. This completes the proof. $\square$

In the following result, we present the sufficient conditions ensuring the existence of solutions for the problem (1)–(2). We apply Leray-Schauder alternative [32] to prove this result.

Theorem 3.2. Assume that

(A1) $F, G : [0, T] \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ are continuous functions and there exist real constants $k_i, \gamma_i \geq 0$, ($i = 1, 2$) and $k_0 > 0$ and $\gamma_0 > 0$ such that, for all $t \in [0, T]$ and $u, v \in \mathbb{R}$,

$$|F(t, u, v)| \leq k_0 + k_1|u| + k_2|v|, \quad |G(t, u, v)| \leq \gamma_0 + \gamma_1|u| + \gamma_2|v|.$$

(A2) $k(N_1 + N_1^*) + \gamma(N_2 + N_2^*) < 1$, where $k = \max\{k_1, k_2\}$, $\gamma = \max\{\gamma_1, \gamma_2\}$.

Then the system (1)–(2) has at least one solution on $[0, T]$.

Proof. Let us first note that continuity of the operator $\mathcal{P}$ follows from that of the functions F and G . Let $\mathbb{B} \subset X \times Y$ be bounded such that $|F(t, u, v)| \leq K_F$, $|G(t, u, v)| \leq K_G$, $\forall (u, v) \in \mathbb{B}$, for positive constants K_F and K_G . Then, for any $(u, v) \in \mathbb{B}$, we have

$$\begin{aligned} & t^{2-\alpha}|\mathcal{P}_1(u, v)(t)| \\ & \leq |\lambda_1|\delta_1 + |\lambda_2\omega_1|\delta_2 + K_F \left[|a_0|\delta_1 \frac{T^2}{2} + |a_1|T\delta_2 + |\mu|\delta_3 \frac{\eta_2^{\alpha+\beta-1}}{\Gamma(\alpha+\beta)} + \delta_3 \sum_{j=1}^n |\sigma_j| \frac{\zeta_j^\alpha}{\Gamma(\alpha+1)} + \frac{T^2}{\Gamma(\alpha+1)} \right] \\ & \quad + K_G \left[|v|\delta_2 \frac{\eta_1^{\alpha+\beta-1}}{\Gamma(\alpha+\beta)} + |b_1|\delta_3 T + \delta_2 \sum_{i=1}^m |\mu_i| \frac{\xi_i^\beta}{\Gamma(\beta+1)} + |b_0\omega_1|\delta_2 \frac{T^2}{2} \right] \\ & = |\lambda_1|\delta_1 + |\lambda_2\omega_1|\delta_2 + K_F e_1 + K_G e_2, \end{aligned}$$

which implies that

$$\|\mathcal{P}_1(u, v)\|_X \leq |\lambda_1|\delta_1 + |\lambda_2\omega_1|\delta_2 + K_F e_1 + K_G e_2.$$

Similarly, one can show that

$$\|\mathcal{P}_2(u, v)\|_Y \leq |\lambda_2|\delta_1^* + |\lambda_1\nu_1|\delta_2^* + K_F e_1^* + K_G e_2^*.$$

In consequence, we get

$$\|\mathcal{P}(u, v)\|_{X \times Y} \leq |\lambda_1|\delta_1 + |\lambda_2|\delta_1^* + |\lambda_2\omega_1|\delta_2 + |\lambda_1\nu_1|\delta_2^* + K_F(e_1 + e_1^*) + K_G(e_2 + e_2^*) < \infty,$$

which shows that the operator $\mathcal{P}$ is uniformly bounded.

Next, we show that $\mathcal{P}$ is equicontinuous. Let $t_1, t_2 \in [0, T]$ with $t_1 > t_2$. Then we have

$$\begin{aligned} & |t_1^{2-\alpha}\mathcal{P}_1(u, v)(t_1) - t_2^{2-\alpha}\mathcal{P}_1(u, v)(t_2)| \\ & \leq |\lambda_1||t_1^{2-\alpha}\rho_1(t_1) - t_2^{2-\alpha}\rho_1(t_2)| + |\lambda_2\omega_1||t_1^{2-\alpha}\rho_2(t_1) - t_2^{2-\alpha}\rho_2(t_2)| \\ & \quad + K_F \left[|a_0||t_1^{2-\alpha}\rho_1(t_1) - t_2^{2-\alpha}\rho_1(t_2)| \frac{T^2}{2} + |a_1|T|t_1^{2-\alpha}\rho_2(t_1) - t_2^{2-\alpha}\rho_2(t_2)| \right. \\ & \quad \left. + |\mu||t_1^{2-\alpha}\rho_3(t_1) - t_2^{2-\alpha}\rho_3(t_2)| \frac{|\eta_2|^{\alpha+\beta-1}}{\Gamma(\alpha+\beta)} + |t_1^{2-\alpha}\rho_3(t_1) - t_2^{2-\alpha}\rho_3(t_2)| \sum_{j=1}^n |\sigma_j| \frac{\zeta_j^\alpha}{\Gamma(\alpha+1)} \right. \\ & \quad \left. + \left| t_1^{2-\alpha} \int_0^{t_1} \frac{(t_1-s)^{\alpha-1}}{\Gamma(\alpha)} ds - t_2^{2-\alpha} \int_0^{t_2} \frac{(t_2-s)^{\alpha-1}}{\Gamma(\alpha)} ds \right| \right] \\ & \quad + K_G \left[|v| \frac{|\eta_1|^{\alpha+\beta-1}}{\Gamma(\alpha+\beta)} |t_1^{2-\alpha}\rho_2(t_1) - t_2^{2-\alpha}\rho_2(t_2)| + |b_1|T|t_1^{2-\alpha}\rho_3(t_1) - t_2^{2-\alpha}\rho_3(t_2)| \right] \end{aligned}$$

$$\begin{aligned}
& + \sum_{i=1}^m |\mu_i| \frac{\xi_i^\beta}{\Gamma(\beta+1)} |t_1^{2-\alpha} \rho_2(t_1) - t_2^{2-\alpha} \rho_2(t_2)| + |b_0 \omega_1| \frac{T^2}{2} |t_1^{2-\alpha} \rho_2(t_1) - t_2^{2-\alpha} \rho_2(t_2)| \Bigg] \\
& \leq |t_1 - t_2| \left\{ \left[|\lambda_1| + |a_0| \frac{T^2}{2} K_F \right] \left[\left| \frac{v_1}{\Lambda} \right| \left| \omega_2 - \frac{b_0 T \Gamma(\beta)}{\Gamma(\alpha)(1+a_1)} \omega_1 \right| \right] \right. \\
& + \left[|\lambda_2 \omega_1| + |a_1| T K_F + |v| K_G \frac{\eta_1^{\alpha+\beta-1}}{\Gamma(\alpha+\beta)} + K_G \sum_{i=1}^m |\mu_i| \frac{\xi_i^\beta}{\Gamma(\beta+1)} + K_G |b_0 \omega_1| \frac{T^2}{2} \right] \left| \frac{\Gamma(\beta)(b_1+1)}{\Lambda \Gamma(\alpha)(a_1+1)} \right| \\
& + \left[\sum_{j=1}^n |\sigma_j| \frac{\zeta_j^\alpha}{\Gamma(\alpha+1)} K_G + |\mu| \frac{|\eta_2|^{\alpha+\beta-1}}{\Gamma(\alpha+\beta)} K_F + |b_1| T K_G \right] \left[\frac{1}{|\Lambda|} \left| \omega_2 - \frac{b_0 T \Gamma(\beta)}{\Gamma(\alpha)(1+a_1)} \omega_1 \right| \right] \Bigg\} \\
& + \frac{2K_F t_1^{2-\alpha}}{\Gamma(\alpha+1)} |t_1 - t_2|^\alpha + K_F \frac{|t_1^2 - t_2^2|}{\Gamma(\alpha+1)} \rightarrow 0 \quad \text{as } t_2 \rightarrow t_1,
\end{aligned}$$

independent of $(u, v) \in \mathbb{B}$. Also, we have

$$\begin{aligned}
& |t_1^{2-\beta} \mathcal{P}_2(u, v)(t_1) - t_2^{2-\beta} \mathcal{P}_2(u, v)(t_2)| \\
& \leq |t_1 - t_2| \left\{ \left[|\lambda_2| + |b_0| K_G \frac{T^2}{2} \right] \left[\left| \frac{\omega_1}{\Lambda} \right| \left| v_2 - \frac{a_0 T}{(1+a_1)} v_1 \right| \right] \right. \\
& + \left[|\lambda_1 v_1| + |a_0 v_1| K_F \frac{T^2}{2} + \mu K_F \frac{\eta_2^{\alpha+\beta-1}}{\Gamma(\alpha+\beta)} + |b_1| T K_G + K_F \sum_{j=1}^n |\sigma_j| \frac{\zeta_j^\alpha}{\Gamma(\alpha)} \right] \left[\frac{1}{|\Lambda|} \right] \\
& + \left[|a_1| T K_F + |v| K_G \frac{\eta_1^{\alpha+\beta-1}}{\Gamma(\alpha+\beta)} + K_G \sum_{i=1}^m |\mu_i| \frac{\xi_i^\beta}{\Gamma(\beta+1)} \right] \left[\frac{1}{|\Lambda|} \left| v_2 - \frac{a_0 T}{(1+a_1)} v_1 \right| \right] \Bigg\} \\
& + \frac{2K_G t_1^{2-\beta}}{\Gamma(\beta+1)} |t_1 - t_2|^\beta + K_G \frac{|t_1^2 - t_2^2|}{\Gamma(\beta+1)} \rightarrow 0 \quad \text{as } t_2 \rightarrow t_1,
\end{aligned}$$

independently of $(u, v) \in \mathbb{B}$. Thus, the operator $\mathcal{P}$ is equicontinuous. Thus, we deduce that the operator $\mathcal{P}$ is completely continuous.

Finally, we consider the set

$$V = \{(u, v) \in X \times Y \mid (u, v) = m\mathcal{P}(u, v), 0 \leq m \leq 1\}$$

and show that it is bounded. Let $(u, v) \in V$ with $(u, v) = m\mathcal{P}(u, v)$, $u = m\mathcal{P}_1(u, v)$, and $v = m\mathcal{P}_2(u, v)$. Then we have

$$\begin{aligned}
\|u\|_X & \leq |\lambda_1| \delta_1 + |\lambda_2 \omega_1| \delta_2 + (N_1 k_1 + N_2 \gamma_1) \|u\|_X + (N_1 k_2 + N_2 \gamma_2) \|v\|_Y + k_0 e_1 + \gamma_0 e_2 \\
& \leq |\lambda_1| \delta_1 + |\lambda_2 \omega_1| \delta_2 + (N_1 k + N_2 \gamma) \|u\|_X + (N_1 k + N_2 \gamma) \|v\|_Y + k_0 e_1 + \gamma_0 e_2, \\
\|v\|_Y & \leq |\lambda_2| \delta_1^* + |\lambda_1 v_1| \delta_2^* + (N_1^* k_1 + N_2^* \gamma_1) \|u\|_X + (k_2 N_1^* + N_2^* \gamma_2) \|v\|_Y + k_0 e_1^* + \gamma_0 e_2^* \\
& \leq |\lambda_2| \delta_1^* + |\lambda_1 v_1| \delta_2^* + (N_1^* k + N_2^* \gamma) \|u\|_X + (k N_1^* + N_2^* \gamma) \|v\|_Y + k_0 e_1^* + \gamma_0 e_2^*,
\end{aligned}$$

which imply that

$$\begin{aligned}
\|u\|_X + \|v\|_Y & \leq |\lambda_1| \delta_1 + |\lambda_2 \omega_1| \delta_2 + |\lambda_2| \delta_1^* + |\lambda_1 v_1| \delta_2^* + [k(N_1 + N_1^*) + \gamma(N_2 + N_2^*)] \|u\|_X \\
& + [k(N_1 + N_1^*) + \gamma(N_2 + N_2^*)] \|v\|_Y + k_0(e_1 + e_1^*) + \gamma_0(e_2 + e_2^*).
\end{aligned}$$

Thus,

$$\|(u, v)\|_{X \times Y} \leq \frac{|\lambda_1| \delta_1 + |\lambda_2 \omega_1| \delta_2 + |\lambda_2| \delta_1^* + |\lambda_1 v_1| \delta_2^* + k_0(e_1 + e_1^*) + \gamma_0(e_2 + e_2^*)}{1 - [k(N_1 + N_1^*) + \gamma(N_2 + N_2^*)]}.$$

Hence, the set V is bounded. Thus, by Leray-Schauder alternative, we deduce that the operator $\mathcal{P}$ has at least one fixed point, which corresponds to the fact that the problem (1)–(2) has at least one solution on $[0, T]$. The proof is completed. $\square$

4 Examples

This section is devoted to the illustration of the results derived in the previous section.

Example 4.1. Consider the system of fractional differential equations consisting of the equations given by

$$\begin{cases} D^{6/5}u(t) = \frac{1}{4\sqrt{1600+t}}(u(t) + \tan^{-1}v(t)) + \cos t, & t \in [0, 1], \\ D^{7/4}v(t) = \frac{1}{\sqrt{2500+t}}(\sin u(t) + v(t)) + t \sin t, & t \in [0, 1], \end{cases} \quad (33)$$

supplemented by the following boundary conditions

$$\begin{cases} D^{-4/5}u(0^+) + \frac{1}{4}D^{-4/5}u(1^-) = -4, \\ D^{1/5}u(0^+) - \frac{3}{2}D^{1/5}u(1^-) = -2\Gamma^{1/5}v\left(\frac{1}{4}\right) - 6v\left(\frac{1}{2}\right) - 4v\left(\frac{2}{3}\right), \\ D^{-1/4}v(0^+) - D^{-1/4}v(1^-) = 2, \\ D^{3/4}v(0^+) - \frac{3}{2}D^{3/4}v(1^-) = 3\Gamma^{3/4}u\left(\frac{1}{3}\right) + \frac{5}{3}u\left(\frac{11}{15}\right) + \frac{2}{5}u\left(\frac{3}{4}\right). \end{cases} \quad (34)$$

Here $\alpha = 6/5$, $\beta = 7/4$, $a_0 = 1/4$, $b_0 = -1/4$, $a_1 = -3/2$, $b_1 = -3/2$, $v = -2$, $\mu = 3$, $\lambda_1 = -4$, $\lambda_2 = 2$, $\mu_1 = -6$, $\mu_2 = -4$, $\sigma_1 = 5/3$, $\sigma_2 = 2/5$, $\eta_1 = 1/4$, $\xi_1 = 1/2$, $\xi_2 = 2/3$, $\eta_2 = 1/3$, $\zeta_1 = 11/15$, $\zeta_2 = 3/4$.

Using the given data, it is found that $\Lambda \approx 70.9531$, $v_1 \approx 2.9182$, $v_2 \approx -6.3910$, $\omega_1 \approx -15.3511$, $\omega_2 \approx 15.2944$, $\delta_1 \approx 0.7330$, $\delta_2 \approx 0.0135$, $\delta_3 \approx 0.3100$, $\delta_1^* \approx 0.8213$, $\delta_2^* \approx 0.0176$, $\delta_3^* \approx 0.0869$, $N_1 \approx 12.2414$, $N_2 \approx 3.6569$, $N_1^* \approx 1.1599$, $N_2^* \approx 9.4324$, $L_1 = 1/160$, $L_2 = 1/50$, and

$$L_1(N_1 + N_1^*) + L_2(N_2 + N_2^*) \approx 0.3455 < 1.$$

Thus, all the conditions of Theorem 3.1 are satisfied and hence the problem (33)–(34) has a unique solution on $[0, 1]$.

Example 4.2. Let us consider the problem (33)–(34) with

$$\begin{aligned} F(t, u(t), v(t)) &= \frac{\sin^2 t}{\sqrt{2+t^3}} + \frac{\tan^{-1}u(t)}{2\sqrt{400+t}} + \frac{v(t)|u(t)|}{50(1+|u(t)|)}, & t \in [0, 1], \\ G(t, u(t), v(t)) &= e^{-t} + \frac{|u(t)|\cos u(t)}{150+t} + \frac{v(t)}{640\sqrt{1+\sin^2 t}}, & t \in [0, 1]. \end{aligned} \quad (35)$$

Clearly, $|F(t, u(t), v(t))| < \frac{1}{2} + \frac{1}{40}|u(t)| + \frac{1}{50}|v(t)|$, $G(t, u(t), v(t)) < 1 + \frac{1}{150}|u(t)| + \frac{1}{640}|v(t)|$, and so $k_0 = 1/2$, $k_1 = 1/40$, $k_2 = 1/50$, $\gamma_0 = 1$, $\gamma_1 = 1/150$, $\gamma_2 = 1/640$, $k = \max\{k_1, k_2\} = 1/40$, $\gamma = \max\{\gamma_1, \gamma_2\} = 1/150$. Moreover,

$$k(N_1 + N_1^*) + \gamma(N_2 + N_2^*) \approx 0.4223 < 1.$$

Therefore, by Theorem 3.2, the problem (33)–(34) with F and G given by (35) has at least one solution on $[0, 1]$.

5 Conclusion

We have investigated the existence and uniqueness of solutions for a nonlinear system of Riemann-Liouville fractional differential equations, equipped with nonseparated semi-coupled integro-multipoint boundary conditions. We apply Banach contraction mapping principle to establish the existence of a

unique solution, while Leray-Schauder alternative is used to obtain the existence result for the problem at hand. We emphasize that the novelty of our results lies on the semi-coupled boundary conditions (2) and enrich the related literature on the topic. Our work also produces some special cases by fixing the parameters involved in the boundary conditions. For example, our results correspond to the ones for nonlocal semi-coupled fractional multipoint boundary conditions by fixing $\nu = 0 = \mu$ and the results for nonlocal semi-coupled fractional integral boundary conditions follow by taking all $\mu_i = 0$, $i = 1, \dots, m$ and $\sigma_j = 0$, $j = 1, \dots, n$.

Acknowledgment: This project was funded by the Deanship of Scientific Research (DSR), King Abdulaziz University, Jeddah, Saudi Arabia under grant no. (KEP-PhD-40-130-41). The authors, therefore, acknowledge with thanks DSR technical and financial support. The authors also thank the reviewers for their constructive remarks on our work.

Conflict of interest: Authors state no conflict of interest.

References

- [1] Z. Bai, *On positive solutions of a nonlocal fractional boundary value problem*, Nonlinear Anal. **72** (2010), 916–924.
- [2] B. Ahmad and J. J. Nieto, *Riemann-Liouville fractional integro-differential equations with fractional nonlocal integral boundary conditions*, Bound. Value Probl. **2011** (2011), 36.
- [3] B. Ahmad, A. Alsaedi, A. Assolami, and R. P. Agarwal, *A study of Riemann-Liouville fractional nonlocal integral boundary value problems*, Bound. Value Probl. **2013** (2013), 274.
- [4] R. Luca, *On a class of nonlinear singular Riemann-Liouville fractional differential equations*, Results Math. **73** (2018), no. 125, 1–15.
- [5] R. P. Agarwal and R. Luca, *Positive solutions for a semipositone singular Riemann-Liouville fractional differential problem*, Int. J. Nonlinear Sci. Numer. Simul. **20** (2019), 823–831.
- [6] R. P. Agarwal, S. Hristova, and D. O'Regan, *Exact solutions of linear Riemann-Liouville fractional differential equations with impulses*, Rocky Mountain J. Math. **50** (2020), 779–791.
- [7] R. Agarwal, S. Hristova, and D. O'Regan, *Existence and Ulam-type stability for nonlinear Riemann-Liouville fractional differential equations with constant delay*, Electron. J. Qual. Theory Differ. Equ. **2020** (2020), no. 67, 1–18.
- [8] F. Mainardi, *Some basic problems in continuum and statistical mechanics*, in: A. Carpinteri, F. Mainardi (eds), Fractals and Fractional Calculus in Continuum Mechanics, Springer, Berlin, 1997, pp. 291–348.
- [9] R. L. Magin, *Fractional Calculus in Bioengineering*, Begell House Publishers, Connecticut, 2006.
- [10] H. A. Fallahgoul, S. M. Focardi, and F. J. Fabozzi, *Fractional Calculus and Fractional Processes with Applications to Financial Economics, Theory and Application*, Elsevier/Academic Press, London, 2017.
- [11] A. K. Golmankhaneh and D. Baleanu, *Calculus on Fractals*, in: C. Cattani, H. M. Srivastava, X.-J. Yang (eds), Fractional Dynamics, Warsaw, Poland, De Gruyter Open, Poland, 2016, pp. 307–332.
- [12] I. M. Sokolov, J. Klafter, and A. Blumen, *Fractional kinetics*, Phys. Today **55** (2002), 48–54.
- [13] N. Nyamoradi, M. Javidi, and B. Ahmad, *Dynamics of SVEIS epidemic model with distinct incidence*, Int. J. Biomath. **8** (2015), no. 6, 1550076.
- [14] I. Petras and R. L. Magin, *Simulation of drug uptake in a two compartmental fractional model for a biological system*, Commun. Nonlinear. Sci. Numer. Simul. **16** (2011), 4588–4595.
- [15] W. Yukunthorn, B. Ahmad, S. K. Ntouyas, and J. Tariboon, *On Caputo-Hadamard type fractional impulsive hybrid systems with nonlinear fractional integral conditions*, Nonlinear Anal. Hybrid Syst. **19** (2016), 77–92.
- [16] N. Heymans and J. C. Bauwens, *Fractal rheological models and fractional differential equations for viscoelastic behavior*, Rheologica Acta **33** (1994), 210–219.
- [17] M. Kirane, B. Ahmad, A. Alsaedi, and M. Al-Yami, *Non-existence of global solutions to a system of fractional diffusion equations*, Acta Appl. Math. **133** (2014), 235–248.
- [18] S. Pati, J. R. Graef, and S. Padhi, *Positive periodic solutions to a system of nonlinear differential equations with applications to Lotka-Volterra-type ecological models with discrete and distributed delays*, J. Fixed Point Theory Appl. **21** (2019), 80.
- [19] J. Henderson and R. Luca, *Systems of Riemann-Liouville fractional equations with multi-point boundary conditions*, Appl. Math. Comput. **309** (2017), 303–323.
- [20] H. T. Tuan, A. Czornik, J. J. Nieto, and M. Niezabitowski, *Global attractivity for some classes of Riemann-Liouville fractional differential systems*, J. Integral Equ. Appl. **31** (2019), no. 2, 265–282.

- [21] A. Alsaedi, S. Aljoudi, and B. Ahmad, *Existence of solutions for Riemann-Liouville type coupled systems of fractional integro-differential equations and boundary conditions*, Electron. J. Differ. Equ. **2016** (2016), 211.
- [22] R. P. Agarwal, B. Ahmad, D. Garout, and A. Alsaedi, *Existence results for coupled nonlinear fractional differential equations equipped with nonlocal coupled flux and multi-point boundary conditions*, Chaos Soliton. Fract. **102** (2017), 149–161.
- [23] R. Luca, *Positive solutions for a system of fractional differential equations with p -Laplacian operator and multi-point boundary conditions*, Nonlinear Anal. Model. Control **23** (2018), 771–801.
- [24] B. Ahmad, A. Alsaedi, S. K. Ntouyas, and Y. Alruwaily, *On a fractional integro-differential system involving Riemann-Liouville and Caputo derivatives with coupled multi-point boundary conditions*, Int. J. Differ. Equ. **15** (2020), no. 2, 209–241.
- [25] A. Alsaedi, R. Luca, and B. Ahmad, *Existence of positive solutions for a system of singular fractional boundary value problems with p -Laplacian operators*, Mathematics **8** (2020), 1890.
- [26] S. Padhi, B. S. R. V. Prasad, and D. Mahendru, *Systems of Riemann-Liouville fractional differential equations with nonlocal boundary conditions – Existence, nonexistence, and multiplicity of solutions: Method of fixed point index*, Math. Meth. Appl. Sci. **44** (2021), no. 10, 8266–8285.
- [27] X. M. Zhang, *A new method for searching the integral solution of system of Riemann-Liouville fractional differential equations with non-instantaneous impulses*, J. Comput. Appl. Math. **388** (2021), 113307.
- [28] B. Ahmad, N. Alghamdi, A. Alsaedi, and S. K. Ntouyas, *Multi-term fractional differential equations with nonlocal boundary conditions*, Open Math. **16** (2018), 1519–1536.
- [29] B. Ahmad, M. Alghanmi, A. Alsaedi, and J. J. Nieto, *Existence and uniqueness results for a nonlinear coupled system involving Caputo fractional derivatives with a new kind of coupled boundary conditions*, Appl. Math. Lett. **116** (2021), 107018.
- [30] A. A. Kilbas, H. M. Srivastava, and J. J. Trujillo, *Theory and Applications of Fractional Differential Equations*, North-Holland Mathematics Studies, vol. 204, Elsevier, Amsterdam, 2006.
- [31] Y. Zhou, *Basic Theory of Fractional Differential Equations*, World Scientific, Singapore, 2014.
- [32] A. Granas and J. Dugundji, *Fixed Point Theory*, Springer-Verlag, New York, 2003.