Yee Seng Tan and Edward R.T. Tiekink*

Crystal structure of bis(μ -*N*-*i*-propyl-*N*-*n*-propyldithiocarbamato- κ^2 S:S') bis(*N*-*i*-propyl-*N*-*n*-propyldithiocarbamato- κ^2 S,S')dizinc(II), $C_{28}H_{56}N_4S_8Zn_2$

Table 1: Data collection and handling.

Crystal:	Prism, colourless
Size:	0.22 × 0.15 × 0.12 mm
Wavelength:	Mo K α radiation (0.71073 Å)
μ :	1.64 mm ⁻¹
Diffractometer, scan mode:	XtaLAB Synergy, φ and ω -scans
$\theta_{\max}$, completeness:	26.4°, >99%
$N(hkl)_{\text{measured}}$, $N(hkl)_{\text{unique}}$, R_{int} :	25064, 4077, 0.058
Criterion for I_{obs} , $N(hkl)_{\text{gt}}$:	$I_{\text{obs}} > 2 \sigma(I_{\text{obs}})$, 3556
$N(\text{param})_{\text{refined}}$:	196
Programs:	CrysAlis ^{PRO} [1], SHELX [2, 3], WinGX and ORTEP [4]

<https://doi.org/10.1515/ncrs-2017-0366>

Received November 26, 2017; accepted February 24, 2018; available online March 16, 2018

Abstract

$C_{28}H_{56}N_4S_8Zn_2$, monoclinic, $P2_1/n$ (no. 14), $a = 9.4123(2)$ Å, $b = 19.2708(4)$ Å, $c = 11.5228(3)$ Å, $\beta = 107.202(2)^\circ$, $V = 1996.54(8)$ Å³, $Z = 2$, $R_{\text{gt}}(F) = 0.0254$, $wR_{\text{ref}}(F^2) = 0.0572$, $T = 100(2)$ K.

CCDC no.: 1825598

The crystal structure is shown in the figure. Tables 1 and 2 contain details on crystal structure and measurement conditions and a list of the atoms including atomic coordinates and displacement parameters.

Source of materials

The compound was obtained from reacting a 1:2 mixture of $ZnCl_2$ (Acros Organic) and $Na[S_2CN(i\text{-Pr})n\text{-Pr}]$ [prepared

from the 1:1:1 reaction of CS_2 (Panreac), $HN(i\text{-Pr})n\text{-Pr}$ (Alfa Aesar) and $NaOH$ (Merck) in acetone solution] in water which resulted in an immediate white precipitate. This was extracted with CH_2Cl_2 and filtered. The filtrate was allowed to stand for slow evaporation under ambient conditions. Colourless crystals formed after a few days. **M.p.**: 384.2–384.4 K. **IR** (cm^{-1}): $\nu(C-S)$ 1189 (s, sh), 971 (s), $\nu(C-N)$ 1454 (s).

Experimental details

The C-bound H atoms were geometrically placed ($C-H = 0.98-1.00$ Å) and refined as riding with $U_{\text{iso}}(H) = 1.2-1.5U_{\text{eq}}(C)$.

Discussion

On-going interest in the structural chemistry of zinc dithiocarbamates ($\sim S_2CNRR'$) is reflected by the observation that there are approaching 250 “hits” for these derivatives in the Cambridge Structural Database (CSD) [5]. The motivations for continuing investigations broadly relate to crystal engineering [6–8] and biological considerations [9, 10]. It was in the former context that the centrosymmetric binuclear title compound, $\{Zn[S_2CN(i\text{-Pr})n\text{-Pr}]_2\}_2$, with disparate R substituents, was studied.

As seen from the figure (70% displacement ellipsoids), the molecular structure comprises equal numbers of bidentate bridging and chelating dithiocarbamate ligands. The Zn–S bond lengths span a relatively narrow range of 2.3335(5) Å [for Zn–S1] to 2.4332(5) Å [Zn–S4]. The coordination geometry at Zn(II) approximates a tetrahedron with angles ranging from a narrow 76.289(17)° (chelate angle) to

*Corresponding author: Edward R.T. Tiekink, Research Centre for Crystalline Materials, School of Science and Technology, Sunway University, 47500 Bandar Sunway, Selangor Darul Ehsan, Malaysia, e-mail: edwardt@sunway.edu.my

Yee Seng Tan: Research Centre for Crystalline Materials, School of Science and Technology, Sunway University, 47500 Bandar Sunway, Selangor Darul Ehsan, Malaysia

Table 2: Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²).

Atom	x	y	z	U _{iso} */U _{eq}
Zn	0.37991(2)	0.01105(2)	0.84818(2)	0.01427(7)
S1	0.23765(5)	−0.07186(2)	0.91107(4)	0.01680(11)
S2	0.37630(5)	0.03187(2)	1.10103(4)	0.01369(10)
S3	0.33473(5)	0.12972(2)	0.81372(4)	0.01565(10)
S4	0.27001(6)	0.01889(2)	0.62911(4)	0.01789(11)
N1	0.22070(17)	−0.07957(8)	1.13595(14)	0.0141(3)
N2	0.17867(17)	0.15084(8)	0.58165(14)	0.0152(3)
C1	0.27372(19)	−0.04443(9)	1.05904(17)	0.0126(4)
C2	0.2486(2)	−0.05890(10)	1.26581(17)	0.0177(4)
H2	0.3058	−0.0144	1.2785	0.021*
C3	0.3434(2)	−0.11270(12)	1.35064(19)	0.0265(5)
H3A	0.4324	−0.1224	1.3259	0.040*
H3B	0.2859	−0.1555	1.3467	0.040*
H3C	0.3729	−0.0949	1.4340	0.040*
C4	0.1023(2)	−0.04519(10)	1.29285(18)	0.0200(4)
H4A	0.0467	−0.0091	1.2380	0.030*
H4B	0.1225	−0.0297	1.3772	0.030*
H4C	0.0433	−0.0880	1.2808	0.030*
C5	0.1321(2)	−0.14292(9)	1.09520(18)	0.0164(4)
H5A	0.1332	−0.1710	1.1674	0.020*
H5B	0.1801	−0.1707	1.0449	0.020*
C6	−0.0290(2)	−0.12934(10)	1.02204(18)	0.0192(4)
H6A	−0.0320	−0.0958	0.9563	0.023*
H6B	−0.0829	−0.1087	1.0755	0.023*
C7	−0.1051(3)	−0.19655(11)	0.9673(2)	0.0280(5)
H7A	−0.2087	−0.1869	0.9212	0.042*
H7B	−0.1027	−0.2296	1.0325	0.042*
H7C	−0.0530	−0.2164	0.9129	0.042*
C8	0.2525(2)	0.10501(9)	0.66396(17)	0.0136(4)
C9	0.1123(2)	0.13041(10)	0.45210(18)	0.0230(5)
H9	0.1032	0.0787	0.4497	0.028*
C10	0.2184(3)	0.15005(13)	0.3801(2)	0.0370(6)
H10A	0.3164	0.1296	0.4187	0.056*
H10B	0.2276	0.2007	0.3785	0.056*
H10C	0.1797	0.1325	0.2969	0.056*
C11	−0.0426(3)	0.15985(12)	0.3985(2)	0.0336(6)
H11A	−0.1057	0.1456	0.4484	0.050*
H11B	−0.0850	0.1424	0.3156	0.050*
H11C	−0.0374	0.2106	0.3970	0.050*
C12	0.1667(2)	0.22389(9)	0.61543(18)	0.0165(4)
H12A	0.1462	0.2531	0.5416	0.020*
H12B	0.2630	0.2389	0.6719	0.020*
C13	0.0444(2)	0.23543(10)	0.67548(19)	0.0191(4)
H13A	−0.0532	0.2245	0.6164	0.023*
H13B	0.0596	0.2033	0.7451	0.023*
C14	0.0422(2)	0.30923(10)	0.7197(2)	0.0271(5)
H14A	−0.0393	0.3146	0.7558	0.041*
H14B	0.0275	0.3413	0.6511	0.041*
H14C	0.1370	0.3196	0.7809	0.041*

129.055(19)° for S1–Zn–S3. This wide angle may be traced to the close approach of the transannular S2 atom which is separated from the zinc atom by 2.9512(5) Å.

In the most recent comprehensive review of the structural chemistry of the binary zinc-triad 1,1-dithiolates, of which dithiocarbamate is an exemplar, two structural motifs were described for molecules of the general formula Zn(S₂CNRR')₂ [11]. One was based on the dimeric structure just described, as exemplified by [Zn(S₂CNMe₂)₂]₂ [12], while the other was monomeric, with Zn(S₂CNCy₂)₂ being archetypal [13]. The adoption of one motif over the other is ascribed to steric effects with bulky substituents precluding dimerisation. Of the approximately 60 structures of Zn(S₂CNRR')₂ included in the CSD [5], the ratio of dimer to monomer is almost exactly 2:1. The fine balance between the adoption of either motif is seen in the structure for R = i-Bu [14]. Here, the crystal comprises equal numbers of each motif. Finally, the structure of the cadmium analogue of the title compound is described in the following paper [15]. Essentially, the same structural motif is observed except that the transannular Cd–S is considered bonding and therefore, the cadmium five-coordinate. This difference arises owing to the larger size of the cadmium atom.

Acknowledgements: Sunway University is thanked for support of biological and crystal engineering studies of metal dithiocarbamates.

References

1. Rigaku/Oxford Diffraction: CrysAlis^{PRO}. Rigaku Corporation, The Woodlands, TX, USA (2017).
2. Sheldrick, G. M.: A short history of SHELX. *Acta Crystallogr. A* **64** (2008) 112–122.
3. Sheldrick, G. M.: Crystal structure refinement with SHELXL. *Acta Crystallogr. C* **71** (2015) 3–8.
4. Farrugia, L. J.: *WinGX* and *ORTEP* for Windows: an update. *J. Appl. Cryst.* **45** (2012) 849–854.
5. Groom, C. R.; Bruno, I. J.; Lightfoot, M. P.; Ward, S. C.: The Cambridge Structural Database. *Acta Crystallogr. B* **72** (2016) 171–179.
6. Benson, R. E.; Ellis, C. A.; Lewis, C. E.; Tiekink, E. R. T.: 3D-, 2D- and 1D-supramolecular structures of {Zn[S₂CN(CH₂CH₂OH)R]₂}₂ and their {Zn[S₂CN(CH₂CH₂OH)R]₂}₂(4,4'-bipyridine) adducts for R = CH₂CH₂OH, Me or Et: polymorphism and pseudo-polymorphism. *CrystEngComm* **9** (2007) 930–941.
7. Poplaukhin, P.; Tiekink, E. R. T.: Interwoven coordination polymers sustained by tautomeric forms of the bridging ligand. *CrystEngComm* **12** (2010) 1302–1306.
8. Manar, K. K.; Yadav, C. L.; Tiwari, N.; Singh, R. K.; Kumar, A.; Drew, M. G. B.; Singh, N.: Effect of functionalities on the crystal structures of new zinc(II) dithiocarbamates: a combined anti-leishmanial and thermal decomposition study. *CrystEngComm* **19** (2017) 2660–2672.
9. Nagy, E. M.; Sitran, S.; Montopoli, M.; Favaro, M.; Marchió, L.; Caparrotta, L.; Fregona, D.: Zinc(II) complexes with dithiocarbamate derivatives: Structural characterisation and biological

- assays on cancerous cell lines. *J. Inorg. Biochem.* **117** (2012) 131–139.
10. Tan, Y. S.; Ooi, K. K.; Ang, K. P.; Akim, A. M.; Cheah, Y.-K.; Halim, S. N. A.; Seng, H.-L.; Tiekink, E. R. T.: Molecular mechanisms of apoptosis and cell selectivity of zinc dithiocarbamates functionalized with hydroxyethyl substituents. *J. Inorg. Biochem.* **150** (2015) 48–62.
 11. Tiekink, E. R. T.: Molecular architecture and supramolecular association in the zinc-triad 1,1-dithiolates. Steric control as a design element in crystal engineering? *CrystEngComm* **5** (2003) 101–113.
 12. Klug, H. P.: The crystal structure of zinc dimethyldithiocarbamate. *Acta Crystallogr.* **21** (1966) 536–546.
 13. Cox, M. J.; Tiekink, E. R. T.: Structural features of zinc(II) bis(O-alkyldithiocarbonate) and zinc(II) bis(N,N-dialkyldithiocarbamate) compounds. *Z. Kristallogr.* **214** (1999) 184–190.
 14. Ivanov, A. V.; Korneeva, E. V.; Gerasimenko, A. V.; Forsling, W.: Structural organization of nickel(II), zinc(II), and copper(II) complexes with diisobutyldithiocarbamate: EPR, C-13 and N-15 CP/MAS NMR, and X-ray diffraction studies. *Russ. J. Coord. Chem.* **31** (2005) 695–707.
 15. Tan, Y. S.; Tiekink, E. R. T.: Crystal structure of bis(μ -*N*-*i*-propyl-*N*-*n*-propyldithiocarbamate- κ^3S,S')bis(*N*-*i*-propyl-*N*-*n*-propyldithiocarbamate- κ^2S,S')dicadmium(II), $C_{28}H_{56}Cd_2N_4S_8$. *Z. Kristallogr. –NCS* **233** (2018) 481–483.