Yee Seng Tan, Chien Ing Yeo and Edward R.T. Tiekink*

Crystal structure of *N*-(3-chlorophenyl)ethoxy-carbothioamide, C₉H₁₀ClNOS

Table 1: Data collection and handling.

Crystal:	Colourless prism
Size:	0.30 × 0.22 × 0.05 mm
Wavelength:	Mo K α radiation (0.71073 Å)
μ :	5.4 cm ⁻¹
Diffractometer, scan mode:	Bruker SMART APEX, ω scans
2 θ _{max} , completeness:	55°, >99%
$N(hkl)$ _{measured} , $N(hkl)$ _{unique} , R_{int} :	9015, 2327, 0.035
Criterion for I_{obs} , $N(hkl)$ _{gt} :	$I_{obs} > 2 \sigma(I_{obs})$, 2015
$N(param)$ _{refined} :	122
Programs:	Bruker programs [1, 2], SHELX [3, 4], ORTEP [5]

<https://doi.org/10.1515/ncrs-2017-0403>

Received December 12, 2017; accepted March 18, 2018; available online March 30, 2018

Abstract

C₉H₁₀ClNOS, monoclinic, C2/c (no. 15), $a = 18.8783(6)$ Å, $b = 4.5518(2)$ Å, $c = 23.6739(8)$ Å, $\beta = 91.501(4)^\circ$, $V = 2033.61(13)$ Å³, $Z = 8$, $R_{gt}(F) = 0.033$, $wR_{ref}(F^2) = 0.082$, $T = 100(2)$ K.

CCDC no.: 1830621

The asymmetric unit of the title crystal structure is shown in the figure. Tables 1 and 2 contain details of the measurement method and a list of the atoms including atomic coordinates and displacement parameters.

Source of materials

3-Chlorophenyl isothiocyanate (Sigma Aldrich; 2.5 mmol, 0.33 mL) was added to NaOH (Merck; 2.5 mmol, 0.10 g) in EtOH (Merck; 3 mL) and the mixture was stirred at room temperature for 2 h, followed by the addition of excess 5 M HCl solution. The resulting mixture was stirred for another 1.5 h. The final product was extracted with chloroform (Merck;

Table 2: Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²).

Atom	<i>x</i>	<i>y</i>	<i>z</i>	U_{iso}^*/U_{eq}
Cl1	0.39796(3)	0.55128(12)	0.24380(2)	0.03764(15)
S1	0.36135(2)	−0.36422(9)	−0.00973(2)	0.01921(12)
O1	0.42262(6)	−0.0977(2)	0.07864(5)	0.0194(2)
N1	0.30855(7)	0.0069(3)	0.06370(6)	0.0163(3)
H1N	0.2704(7)	−0.028(4)	0.0438(7)	0.020*
C1	0.36499(8)	−0.1440(3)	0.04656(6)	0.0157(3)
C2	0.30022(8)	0.2121(3)	0.10787(6)	0.0163(3)
C3	0.35058(8)	0.2705(4)	0.15030(7)	0.0199(3)
H3	0.3948	0.1711	0.1513	0.024*
C4	0.33465(9)	0.4769(4)	0.19103(7)	0.0218(4)
C5	0.27084(9)	0.6245(4)	0.19176(7)	0.0220(3)
H5	0.2614	0.7648	0.2203	0.026*
C6	0.22107(9)	0.5617(4)	0.14962(7)	0.0223(4)
H6	0.1765	0.6586	0.1494	0.027*
C7	0.23542(8)	0.3593(3)	0.10777(7)	0.0200(3)
H7	0.2009	0.3206	0.0788	0.024*
C8	0.48604(8)	−0.2722(4)	0.06945(7)	0.0223(4)
H2A	0.5083	−0.2145	0.0337	0.027*
H2B	0.4742	−0.4839	0.0676	0.027*
C9	0.53472(9)	−0.2092(5)	0.11901(8)	0.0339(5)
H3A	0.5780	−0.3261	0.1159	0.051*
H3B	0.5111	−0.2605	0.1540	0.051*

*Corresponding author: Edward R.T. Tiekink, Research Centre for Crystalline Materials, School of Science and Technology, Sunway University, 47500 Bandar Sunway, Selangor Darul Ehsan, Malaysia, e-mail: edwardt@sunway.edu.my

Yee Seng Tan and Chien Ing Yeo: Research Centre for Crystalline Materials, School of Science and Technology, Sunway University, 47500 Bandar Sunway, Selangor Darul Ehsan, Malaysia

15 mL) and left for evaporation at room temperature, yielding colourless crystals after 2 weeks. M.p. (Krüss KSP1N): 349–350 K. IR (Perkin Elmer Spectrum 400 F T Mid-IR/Far-IR; cm⁻¹): 3248 (s) ν (N–H), 1476 (s) ν (C–N), 1199 (s) ν (C=S), 1040 (s) ν (C–O). Elem. Anal. (Perkin Elmer PE 2400 CHN):

Calc. for C₉H₁₀ClNOS: C, 50.11; H, 4.67; N, 6.49%. Found: C, 50.21; H, 4.55; N, 6.69%.

Experimental details

The C-bound H atoms were geometrically placed (C—H = 0.95–0.99 Å) and refined as riding with $U_{\text{iso}}(\text{H}) = 1.2\text{--}1.5 U_{\text{eq}}(\text{C})$. The N-bound H-atoms were located in a difference Fourier map but were refined with a distance restraint of N—H = 0.88 ± 0.01 Å, and with $U_{\text{iso}}(\text{H})$ set to $1.2 U_{\text{eq}}(\text{N})$. Owing to poor agreement, two reflections, i.e. (2 0 0) and (0 0 2), were omitted from the final cycles of refinement, presumably owing to interference from the beam-stop.

Comment

The structural landscape of molecules of the general formula ROC(=S)N(H)R', for R, R' = alkyl/aryl, has been summarized recently [6]. The overwhelming majority of structures, including a more recently reported derivative [7], share common features such as a *syn*-disposition of the thioamide-N—H and thione-S atoms [6, 7] with the exceptions, i.e. with anti-dispositions of these atoms, arising owing to the dictates of thioamide-N—H...O, N hydrogen bonding, as the O— and/or N-bound substituents carry good oxygen or nitrogen hydrogen bond acceptors [6]. In continuation of these studies, the title compound EtOC(=S)N(H)(3-ClPh) was investigated by X-ray crystallography.

The molecular structure is shown in the Figure (50% displacement ellipsoids) and this features the anticipated *syn*-disposition of the thioamide-N—H and thione-S atoms. The central CNOS chromophore is planar and forms a dihedral angle of $7.94(9)^\circ$ with the N-bound 3-chlorophenyl ring. The whole molecule is practically planar with the r.m.s. for the 13 fitted non-hydrogen atoms being 0.0594 Å with maximum deviations of 0.1186(12) and 0.0973(12) Å for the O1 and N1 atoms, respectively. Overall, the molecule has a U-shape as

the ethoxy and phenyl-chloro groups lie to the same side of the molecule.

In the molecular packing, the *syn*-disposition of the thioamide-N—H and thione-S atoms facilitates the formation of thioamide-N—H...S(thione) hydrogen bonds between centrosymmetrically related molecules and therefore, eight-membered thioamide synthons $\{\cdots\text{HNCS}\}_2$; N1—H1n...S1ⁱ: 2.641(14) Å and $164.7(14)^\circ$ for symmetry operation *i*: $1/2 - x, -1/2 - y, -z$. Globally, in the crystal, dimeric aggregates stack in columns parallel to the *b* axis.

Acknowledgements: The University of Malaya's X-ray laboratory is thanked for the data collection. Sunway University is thanked for support of biological and crystal engineering studies of metal alkoxycarbothioamides.

References

1. Krause, L.; Herbst-Irmer, R.; Sheldrick, G. M.; Stalke, D.: Comparison of silver and molybdenum microfocus X-ray sources for single-crystal structure determination. *J. Appl. Cryst.* **48** (2015) 3–10.
2. Bruker. APEX2 and SAINT. Bruker AXS Inc., Madison, WI, USA (2008).
3. Sheldrick, G. M.: A short history of SHELX. *Acta Crystallogr.* **A71** (2015) 3–8.
4. Sheldrick, G. M.: Crystal structure refinement with SHELXL. *Acta Crystallogr.* **C71** (2015) 3–8.
5. Farrugia, L. J.: WinGX and ORTEP for Windows: an update. *J. Appl. Cryst.* **45** (2012) 849–854.
6. Jotani, M. M.; Yeo, C. I.; Tiekink, E. R. T.: A new monoclinic polymorph of *N*-(3-methylphenyl) ethoxycarbothioamide: crystal structure and Hirshfeld surface analysis. *Acta Crystallogr.* **E73** (2017) 1889–1897.
7. Yeo, C. I.; Tiekink, E. R. T.: Crystal structure of *N*-(2-methylphenyl)ethoxycarbothioamide, C₁₀H₁₃NOS. *Z. Kristallogr. – NCS* **233** (2018) 299–301.