Wan-Wan Wang, Xiao-Wei Deng* and Wei-Na Wu

Crystal structure of dichlorido-bis(*N*-benzyl-2-(quinolin-8-yloxy)acetamide- $\kappa^2 N,O$)copper(II) – ethyl acetate (1/1), $C_{38}H_{36}N_4O_6Cl_2Cu$

Table 1: Data collection and handling.

Crystal:	Green block
Size:	0.21 × 0.18 × 0.16 mm
Wavelength:	Mo K α radiation (0.71073 Å)
μ :	0.80 mm ⁻¹
Diffractometer, scan mode:	Bruker APEX-II, φ and ω
$\theta_{\max}$, completeness:	25.0°, 99%
$N(hkl)_{\text{measured}}$, $N(hkl)_{\text{unique}}$, R_{int} :	9637, 6346, 0.024
Criterion for I_{obs} , $N(hkl)_{\text{gt}}$:	$I_{\text{obs}} > 2 \sigma(I_{\text{obs}})$, 4503
$N(\text{param})_{\text{refined}}$:	460
Programs:	Bruker [1], SHELX [2]

contains the list of the atoms including atomic coordinates and displacement parameters.

Source of material

N-Phenyl-2-(quinolin-8-yloxy)acetamide [3] (0.280 g, 1 mmol) was dissolved in ethyl acetate (10 mL), then an ethyl acetate solution (10 mL) containing copper(II) chloride dihydrate (0.179 g, 1 mmol) was added dropwise at room temperature. After stirring for 2 h, the mixture was filtered and set aside to crystallize at room temperature for several days, giving green block crystals.

Experimental details

The hydrogen atoms were placed at calculated positions and refined as riding atoms with isotropic displacement parameters.

Comment

The structures and luminescent properties of the complexes with amide type ligands bearing quinoline ring have been investigated in our previous work [4–7]. As part of our ongoing work, the title complex was synthesized and characterized.

In the title complex, there exist one complex and one crystal ethyl acetate molecule. The six-coordinated Cu atom is in a distorted octahedral geometry coordinated by two O atoms and two N atoms from two independent ligands, and two chlorido ligands. In the crystal, a pair of intermolecular N–H...Cl hydrogen bonds between two complex molecules create a centrosymmetric dimer.

<https://doi.org/10.1515/ncrs-2019-0023>

Received January 8, 2019; accepted March 11, 2019; available online April 22, 2019

Abstract

$C_{38}H_{36}N_4O_6Cl_2Cu$, triclinic, $P\bar{1}$ (no. 2), $a = 10.749(5)$ Å, $b = 11.074(5)$ Å, $c = 15.951(7)$ Å, $\alpha = 80.471(8)^\circ$, $\beta = 80.987(8)^\circ$, $\gamma = 77.283(8)^\circ$, $V = 1812.3(14)$ Å³, $Z = 2$, $R_{\text{gt}}(F) = 0.0465$, $wR_{\text{ref}}(F^2) = 0.1279$, $T = 296(2)$ K.

CCDC no.: 1902365

The asymmetric unit of the title crystal structure is shown in the figure. Table 1 contains crystallographic data and Table 2

*Corresponding author: Xiao-Wei Deng, School of Physics and Electronic Information Engineering, Henan Polytechnic University, Jiaozuo 454000, P.R. China, e-mail: 383082433@qq.com

Wan-Wan Wang: College of Chemistry and Chemical Engineering, Henan Polytechnic University, Jiaozuo 454000, P.R. China

Wei-Na Wu: School of Physics and Electronic Information Engineering, Henan Polytechnic University, Jiaozuo 454000, P.R. China

Table 2: Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²).

Atom	x	y	z	<i>U</i> _{iso} */ <i>U</i> _{eq}
Cu1	0.60611(4)	0.37282(4)	0.29338(2)	0.04625(15)
O1	0.4715(2)	0.3647(2)	0.42825(14)	0.0539(6)
O2	0.2351(2)	0.3063(2)	0.42852(16)	0.0642(7)
O3	0.7468(2)	0.3616(2)	0.15939(14)	0.0568(6)
O4	1.0215(3)	0.5015(3)	0.0964(2)	0.0994(11)
O5	1.0138(5)	0.0845(4)	0.2071(3)	0.1529(18)
O6	1.2116(4)	0.0257(4)	0.2350(3)	0.1226(13)
N1	0.6389(2)	0.1930(3)	0.34953(18)	0.0475(7)
N2	0.1637(3)	0.4015(3)	0.54707(18)	0.0530(7)
H2A	0.1830	0.4518	0.5764	0.064*
N3	0.5059(3)	0.3329(3)	0.20803(17)	0.0488(7)
N4	0.8138(3)	0.5916(3)	0.13613(19)	0.0617(8)
H4A	0.7401	0.5730	0.1557	0.074*
C1	0.7159(4)	0.1060(4)	0.3095(3)	0.0634(10)
H1A	0.7379	0.1234	0.2507	0.076*
C2	0.7667(4)	−0.0120(4)	0.3508(4)	0.0798(13)
H2B	0.8205	−0.0714	0.3197	0.096*
C3	0.7372(4)	−0.0385(4)	0.4357(3)	0.0771(13)
H3A	0.7717	−0.1162	0.4640	0.093*
C4	0.6546(3)	0.0504(4)	0.4816(3)	0.0612(10)
C5	0.6195(4)	0.0307(5)	0.5709(3)	0.0780(13)
H5A	0.6516	−0.0454	0.6022	0.094*
C6	0.5404(4)	0.1205(5)	0.6113(3)	0.0795(14)
H6A	0.5194	0.1063	0.6704	0.095*
C7	0.4887(4)	0.2356(4)	0.5658(2)	0.0664(11)
H7A	0.4341	0.2970	0.5949	0.080*
C8	0.5182(3)	0.2575(3)	0.4798(2)	0.0505(9)
C9	0.6049(3)	0.1665(3)	0.4357(2)	0.0473(8)
C10	0.3570(3)	0.4451(3)	0.4602(2)	0.0591(10)
H10A	0.3380	0.5192	0.4188	0.071*
H10B	0.3693	0.4710	0.5130	0.071*
C11	0.2453(3)	0.3764(3)	0.4766(2)	0.0504(8)
C12	0.0501(3)	0.3563(3)	0.5795(2)	0.0496(8)
C13	0.0026(4)	0.3685(5)	0.6630(3)	0.0822(14)
H13A	0.0453	0.4037	0.6961	0.099*
C14	−0.1102(5)	0.3278(6)	0.6979(3)	0.1067(18)
H14A	−0.1440	0.3376	0.7543	0.128*
C15	−0.1715(4)	0.2739(5)	0.6505(4)	0.0894(15)
H15A	−0.2457	0.2447	0.6748	0.107*
C16	−0.1248(4)	0.2628(4)	0.5684(3)	0.0756(12)
H16A	−0.1676	0.2266	0.5360	0.091*
C17	−0.0144(3)	0.3042(3)	0.5315(3)	0.0603(10)
H17A	0.0165	0.2969	0.4744	0.072*
C18	0.3868(4)	0.3214(4)	0.2319(2)	0.0651(10)
H18A	0.3562	0.3164	0.2900	0.078*
C19	0.3026(4)	0.3163(4)	0.1739(3)	0.0774(13)
H19A	0.2177	0.3102	0.1934	0.093*
C20	0.3464(4)	0.3201(4)	0.0900(3)	0.0765(12)
H20A	0.2921	0.3159	0.0510	0.092*
C21	0.4741(4)	0.3306(3)	0.0615(2)	0.0584(9)
C22	0.5243(5)	0.3402(4)	−0.0258(3)	0.0704(11)
H22A	0.4743	0.3334	−0.0668	0.084*
C23	0.6451(5)	0.3593(4)	−0.0497(3)	0.0764(12)
H23A	0.6770	0.3666	−0.1076	0.092*
C24	0.7234(4)	0.3684(4)	0.0097(2)	0.0643(10)
H24A	0.8058	0.3833	−0.0087	0.077*

Table 2 (continued)

Atom	x	y	z	<i>U</i> _{iso} */ <i>U</i> _{eq}
C25	0.6793(3)	0.3556(3)	0.0946(2)	0.0492(8)
C26	0.5520(3)	0.3382(3)	0.1227(2)	0.0480(8)
C27	0.8804(3)	0.3661(4)	0.1417(3)	0.0677(11)
H27A	0.9175	0.3212	0.0935	0.081*
H27B	0.9222	0.3224	0.1909	0.081*
C28	0.9104(4)	0.4945(4)	0.1219(2)	0.0647(10)
C29	0.8170(4)	0.7186(4)	0.1234(2)	0.0647(10)
C30	0.7116(6)	0.8039(5)	0.1427(4)	0.114(2)
H30A	0.6359	0.7769	0.1659	0.137*
C31	0.7108(7)	0.9296(6)	0.1296(5)	0.134(2)
H31A	0.6348	0.9854	0.1436	0.161*
C32	0.8159(8)	0.9732(6)	0.0975(4)	0.124(2)
H32A	0.8141	1.0587	0.0875	0.149*
C33	0.9230(9)	0.8923(8)	0.0802(6)	0.181(4)
H33A	0.9972	0.9225	0.0576	0.218*
C34	0.9300(5)	0.7672(6)	0.0937(5)	0.146(3)
H34A	1.0087	0.7132	0.0836	0.176*
C35	1.0649(5)	0.1060(6)	0.3441(4)	0.126(2)
H35A	1.1432	0.0849	0.3699	0.188*
H35B	1.0319	0.1943	0.3412	0.188*
H35C	1.0029	0.0616	0.3779	0.188*
C36	1.0908(6)	0.0710(5)	0.2572(5)	0.1075(18)
C37	1.2403(7)	−0.0079(9)	0.1474(5)	0.181(3)
H37A	1.1998	−0.0764	0.1432	0.217*
H37B	1.2086	0.0630	0.1065	0.217*
C38	1.3799(8)	−0.0448(9)	0.1300(6)	0.226(5)
H38A	1.4028	−0.0675	0.0733	0.338*
H38B	1.4187	0.0237	0.1344	0.338*
H38C	1.4100	−0.1150	0.1709	0.338*
Cl1	0.52520(9)	0.57859(9)	0.25784(6)	0.0611(3)
Cl2	0.77810(9)	0.39807(9)	0.35042(6)	0.0622(3)

References

1. Bruker. SMART and SAINT. Bruker AXS Inc., Madison, WI, USA (2007).
2. Sheldrick, G. M.: Crystal refinement with SHELX. *Acta Crystallogr.* **C71** (2015) 3–8.
3. Chen, L.; Wu, W.-N.: Crystal structure of dichlorido-bis(*N*-phenyl-2-(quinolin-8-yloxy)acetamide-κ²*N,O*) – acetone (2/1), C_{35.5}H₃₁N₄O_{4.5}Cl₂Cu. *Z. Kristallogr. NCS* **233** (2018) 275–277.
4. Wu, W.-N.; Yuan, W.-B.; Tang, N.; Yang, R.-D.; Yan, L.; Xu, Z.-H.: Synthesis, characterizations and luminescent properties of three novel aryl amide type ligands and their lanthanide complexes. *Spectrochim. Acta, Part A* **65** (2006) 912–918.
5. Wu, W.-N.; Tang, N.; Yan, L.: Syntheses, characterization and fluorescent properties of six novel lanthanide complexes with *N,N*-diphenyl-2-(quinolin-8-yloxy)acetamide. *J. Fluoresc.* **18** (2008) 101–107.
6. Cai, H.-X.; Wu, W.-N.; Wang, Y.: Synthesis and crystal structures of two copper(II) complexes containing amide type ligands. *Chin. J. Inorg. Chem.* **18** (2013) 845–849.
7. Lin, L.; Li, X.-H.; Zhang, B.; Zhang, Z.-Y.; Wu, W.-N.; Wang, Y.: Syntheses, crystal structures and fluorescence properties of Cu(II)/Zn(II)/Ag(I) complexes with an amide type ligand. *Chin. J. Inorg. Chem.* **32** (2016) 1653–1658.