Yee Seng Tan and Edward R.T. Tiekink*

Crystal structure of *catena*-{poly[bis(*O,O'*-diethyldithiophosphato-*S*)-(μ₂-1,2-bis(4-pyridylmethylene)hydrazine-*N,N'*)-zinc(II)] di-acetonitrile solvate}, {C₂₀H₃₀N₄O₄P₂S₄Zn · 2 C₂H₃N}_n

Table 1: Data collection and handling.

Crystal:	Yellow prism
Size:	0.21 × 0.11 × 0.11 mm
Wavelength:	Cu Kα radiation (1.54184 Å)
μ:	4.59 mm ⁻¹
Diffractometer, scan mode:	XtaLAB Synergy, ω
θ _{max} , completeness:	67.1°, >99%
N(hkl) _{measured} , N(hkl) _{unique} , R _{int} :	36948, 5995, 0.024
Criterion for I _{obs} , N(hkl) _{gt} :	I _{obs} > 2 σ(I _{obs}), 5730
N(param) _{refined} :	376
Programs:	CrysAlis ^{PRO} [1], SHELX [2, 3], WinGX/ORTEP [4]

A part of the coordination polymer is shown in the figure. Table 1 contains crystallographic data and Table 2 contains the list of the atoms including atomic coordinates and displacement parameters.

Source of material

The Zn[S₂P(OEt)₂]₂ precursor was prepared in high yield from the *in situ* reaction of Zn(NO₃)₂·6 H₂O (Alfa Aesar; 14.87 g, 0.05 mol), EtOH (Merck; 12.25 mL, 0.21 mol), P₂S₅ (Sigma-Aldrich; 11.11 g, 0.05 mol) and 50° w/w NaOH solution (Merck; 8.80 mL, 0.11 mol). The ligand, 1,2-bis(4-pyridylmethylene)hydrazine, was prepared in high yield from the 2:1 reflux reaction of 4-picolyamine (Aldrich; 2.03 mL, 0.02 mol) and hydrazinium hydroxide (Merck; 0.49 mL, 0.01 mol) in ethanol solution (Merck; 5 mL). The title compound was obtained by mixing the suspension of Zn[S₂P(OEt)₂]₂ (0.50 g, 1.15 mmol) and 1,2-bis(4-pyridylmethylene)hydrazine (0.25 g, 1.20 mmol) in dimethylformamide (Merck; 5 mL), followed by stirring for 30 min at 373 K. The solution was filtered and the filtrate was collected in a sample vial containing acetonitrile (Merck; 1 mL). Yellow blocks formed after one day. Yield: 0.55 g, (66%, based on Zn[S₂P(OEt)₂]₂). M. pt. (Biobase Automatic Melting Apparatus MP45): 387.15–388.95 K. IR (Bruker Vertex 70 V equipped with Platinum ATR from 400 to 80 cm⁻¹, cm⁻¹): 1015(s) ν(C–O), 932(s) ν(P–O), 661(s) ν(P–S)_{asym}, 522(m) ν(P–S)_{sym}. ¹H NMR (Bruker Ascend 400 MHz, d₆-DMSO): δ [p.p.m.] 8.75 (dd, 4H, 2-aryl-H, ³J_{HH} = 6.04 Hz, ⁴J_{HH} = 2.92 Hz), 8.70 (s, 2H, N=CH), 7.83 (dd, 4H, 3-aryl-H, ³J_{HH} = 6.08 Hz, ⁴J_{HH} = 2.92 Hz), 3.82 (dq, 8H, OCH₂, ³J_{PH} = 9.08 Hz, ³J_{HH} = 7.10 Hz), 2.08

<https://doi.org/10.1515/ncrs-2019-0116>

Received February 15, 2019; accepted March 16, 2019; available online July 9, 2019

Abstract

C₂₄H₃₆N₆O₄P₂S₄Zn, triclinic, *P* $\bar{1}$ (no. 2), *a* = 11.2847(2) Å, *b* = 12.6967(3) Å, *c* = 13.6595(3) Å, α = 64.738(2)°, β = 72.181(2)°, γ = 76.681(2)°, *V* = 1673.68(7) Å³, *Z* = 2, *R*_{gt}(*F*) = 0.0225, *wR*_{ref}(*F*²) = 0.0582, *T* = 100(2) K.

CCDC no.: 1903612

*Corresponding author: Edward R.T. Tiekink, Research Centre for Crystalline Materials, School of Science and Technology, Sunway University, 47500 Bandar Sunway, Selangor Darul Ehsan, Malaysia, e-mail: edwardt@sunway.edu.my

Yee Seng Tan: Research Centre for Crystalline Materials, School of Science and Technology, Sunway University, 47500 Bandar Sunway, Selangor Darul Ehsan, Malaysia

Table 2: Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²).

Atom	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> _{iso} [*] / <i>U</i> _{eq}
Zn	0.36954(2)	0.36824(2)	0.39290(2)	0.01202(6)
S1	0.23479(4)	0.25914(3)	0.39044(3)	0.01609(9)
S2	0.44385(4)	0.31226(4)	0.14029(3)	0.02028(9)
S3	0.30974(4)	0.42341(3)	0.54632(3)	0.01496(9)
S4	0.38117(4)	0.13012(3)	0.67850(3)	0.01831(9)
P1	0.27842(4)	0.27156(3)	0.23001(3)	0.01397(9)
P2	0.26007(4)	0.26812(3)	0.66833(3)	0.01314(9)
O1	0.17459(11)	0.36124(10)	0.17050(10)	0.0194(2)
O2	0.24593(10)	0.15270(9)	0.23711(9)	0.0173(2)
O3	0.22195(11)	0.30000(9)	0.77577(9)	0.0170(2)
O4	0.13372(10)	0.24194(9)	0.65770(9)	0.0171(2)
N1	0.54321(12)	0.27137(11)	0.39660(11)	0.0141(3)
N2	0.95302(13)	0.04823(12)	0.48961(12)	0.0184(3)
N3	0.38850(12)	0.53304(11)	0.27162(11)	0.0134(3)
N4	0.49900(13)	0.94340(11)	0.04248(11)	0.0170(3)
N5	0.13990(18)	0.66076(17)	0.96912(18)	0.0445(5)
N6	1.11947(17)	−0.07987(16)	0.07853(17)	0.0408(4)
C1	0.04195(16)	0.34375(16)	0.21888(16)	0.0241(4)
H1A	0.032927	0.273662	0.289388	0.029*
H1B	0.007302	0.331212	0.166792	0.029*
C2	−0.0273(2)	0.44934(18)	0.2404(2)	0.0418(5)
H2A	−0.117089	0.441069	0.267596	0.063*
H2B	−0.013376	0.519058	0.171264	0.063*
H2C	0.002783	0.457531	0.296666	0.063*
C3	0.33081(16)	0.04584(14)	0.26961(14)	0.0189(3)
H3A	0.344472	0.027818	0.343627	0.023*
H3B	0.412775	0.054799	0.214865	0.023*
C4	0.27153(17)	−0.05114(15)	0.27361(15)	0.0235(4)
H4A	0.326712	−0.124904	0.295402	0.035*
H4B	0.258642	−0.032382	0.199909	0.035*
H4C	0.190628	−0.059253	0.328134	0.035*
C5	0.17875(19)	0.20868(16)	0.88552(14)	0.0248(4)
H5A	0.229800	0.132644	0.890510	0.030*
H5B	0.090138	0.199753	0.896697	0.030*
C6	0.19155(17)	0.24412(16)	0.97293(14)	0.0228(4)
H6A	0.162991	0.184123	1.046802	0.034*
H6B	0.140450	0.319232	0.967611	0.034*
H6C	0.279576	0.252198	0.961565	0.034*
C7	0.02414(17)	0.32960(16)	0.64874(17)	0.0266(4)
H7A	0.048823	0.406397	0.591597	0.032*
H7B	−0.013492	0.338386	0.720901	0.032*
C8	−0.06881(17)	0.29051(18)	0.61691(17)	0.0293(4)
H8A	−0.140728	0.351011	0.605847	0.044*
H8B	−0.097240	0.217280	0.676360	0.044*
H8C	−0.029327	0.277624	0.547664	0.044*
C9	0.56162(16)	0.16826(14)	0.38555(14)	0.0176(3)
H9	0.498765	0.148442	0.365428	0.021*
C10	0.66789(16)	0.08988(14)	0.40218(14)	0.0181(3)
H10	0.677403	0.017762	0.393576	0.022*
C11	0.76067(15)	0.11732(14)	0.43154(13)	0.0149(3)
C12	0.74283(15)	0.22512(14)	0.44183(14)	0.0171(3)
H12	0.805142	0.247775	0.460179	0.021*
C13	0.63360(15)	0.29795(14)	0.42496(14)	0.0166(3)
H13	0.621354	0.370306	0.433719	0.020*
C14	0.87041(15)	0.03032(14)	0.45404(13)	0.0167(3)

Table 2 (continued)

Atom	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> _{iso} [*] / <i>U</i> _{eq}
H14	0.879863	−0.039831	0.441983	0.020*
C15	0.32950(15)	0.58031(13)	0.18764(13)	0.0150(3)
H15	0.278337	0.534234	0.182018	0.018*
C16	0.34066(15)	0.69352(14)	0.10925(13)	0.0156(3)
H16	0.297096	0.724345	0.051494	0.019*
C17	0.41625(15)	0.76200(13)	0.11561(13)	0.0139(3)
C18	0.47594(15)	0.71347(14)	0.20364(13)	0.0152(3)
H18	0.527312	0.757754	0.211425	0.018*
C19	0.45930(15)	0.60071(14)	0.27886(13)	0.0151(3)
H19	0.499589	0.568848	0.338864	0.018*
C20	0.43083(15)	0.88201(14)	0.03285(13)	0.0155(3)
H20	0.390281	0.913673	−0.026980	0.019*
C21	0.21585(19)	0.59990(17)	0.93470(17)	0.0291(4)
C22	0.31250(18)	0.52305(17)	0.89175(16)	0.0300(4)
H22A	0.339860	0.456145	0.953821	0.045*
H22B	0.383920	0.566160	0.842211	0.045*
H22C	0.279202	0.494782	0.850125	0.045*
C23	1.0354(2)	−0.02649(18)	0.11918(17)	0.0332(5)
C24	0.9283(2)	0.04170(19)	0.16674(18)	0.0394(5)
H24A	0.861349	−0.008259	0.213589	0.059*
H24B	0.897335	0.108283	0.106446	0.059*
H24C	0.954489	0.070652	0.211969	0.059*

(s, 6H, CH₃CN), 1.14 (t, 12H, CH₃, ³*J*_{HH} = 7.08 Hz). ¹³C{¹H} NMR (Bruker Ascend 400 MHz, d₆-DMSO): δ [p.p.m.] 159.6 (C= N), 150.5 (2-C₅H₄N), 140.6 (1-C₅H₄N), 122.2 (3-C₅H₄N), 118.1 (CH₃CN), 60.0 (d, OCH₂, ²*J*_{PC} = 6.69 Hz), 16.1 (d, CH₃, ³*J*_{PC} = 8.75 Hz), 1.2 (CH₃CN). ³¹P{¹H} NMR (Bruker Ascend 400 MHz, d₆-DMSO): δ [p.p.m.] 112.2.

Experimental details

The C-bound H atoms were geometrically placed (C–H = 0.95–0.99 Å) and refined as riding with *U*_{iso}(H) = 1.2–1.5*U*_{eq}(C). Analysis of the residual electron density revealed that one of the two acetonitrile molecules (C23, C24, N6) is disordered over two positions with occupancy factors of 0.9 and 0.1. Refinement of a heavily restrained/constrained disordered model was unsatisfactory and was not included in the final refinement.

Comment

Four-coordinate ZnS₄ geometries and supramolecular association leading to dimers and chains characterize the structural chemistry of the homoleptic zinc dithiophosphates, {Zn[S₂P(OR)₂]₂}_n [5]. The addition of potentially bridging bipyridyl donors might be expected to increase the dimensionality of the resultant species; however, four-coordination prevails, leading to binuclear species or polymeric chains only [6]. The adoption of zero- or one-dimensional species is regulated by steric effects associated

with the span of the bridge provided by the bipyridyl donor and/or the steric bulk of the remote R group in the dithiophosphate ligand [6–8]. A key example of this phenomenon is found when R = cyclohexyl in $Zn[S_2P(OR)_2]_2$. Thus, when the bipyridyl donor is 4,4'-bipyridine, only the binuclear structure, *i.e.* $\{Zn[S_2P(OCy)_2]_2\}_2(4,4'\text{-bipyridine})$, could be isolated despite the stoichiometry of the respective reagents employed in the reaction [8]. When the bridging ligand was 1,2-bis(4-pyridylmethylene)hydrazine, *i.e.* with a significantly greater distance between the bridging-nitrogen atoms, a zigzag chain for $\{Zn[S_2P(OCy)_2]_2(1,2\text{-bis}(4\text{-pyridylmethylene)hydrazine})\}_2$ ensued [9]. In continuation of systematic studies in this area [7–10], herein the crystal and molecular structures of polymeric $\{Zn[S_2P(OEt)_2]_2(1,2\text{-bis}(4\text{-pyridylmethylene)hydrazine})\}_n$, isolated with two molecules of acetonitrile per repeat unit, are described.

The asymmetric unit is shown in the figure (70% displacement ellipsoids; symmetry operations i: $2 - x, -y, 1 - z$ and ii: $1 - x, 2 - y, -z$) and has been extended to show the full bridging 1,2-bis(4-pyridylmethylene)hydrazine molecules (each disposed about a centre of inversion) and to highlight the zigzag topology of the chain thus formed. The zinc(II) centre is coordinated by two monodentate dithiophosphate ligands [$Zn-S1, S3 = 2.2964(4)$ and $2.3468(4)$ Å] and two nitrogen atoms derived from two bridging 1,2-bis(4-pyridylmethylene)hydrazine ligands [$Zn-N1, N3 = 2.0624(13)$ and $2.0493(13)$ Å]. As noted earlier [10], the disparity in the P–S bond lengths [$P1-S1, S2 = 2.0372(6)$ and $1.9385(6)$ Å; $P2-S3, S4 = 2.0278(5)$ and $1.9396(6)$ Å], with the short P–S bonds associated with the non-coordinating sulfur atoms, is as expected. The relatively narrow range of angles subtended at the zinc atom, $S3-Zn-N3$ [$97.00(4)^\circ$] to $S1-Zn-N3$ [$118.58(4)^\circ$] indicated minor deviations from a tetrahedral geometry for the N_2S_2 donor set.

In the crystal, the chains are approximately orientated along $[1 \bar{2} 1]$. These are connected into a three-dimensional architecture by pyridyl-C–H...S [$C10-H10 \cdots S4^i$: $H10 \cdots S4^i = 2.73$ Å, $C10 \cdots S4^i = 3.6106(19)$ Å with angle at $H10 = 154^\circ$ for symmetry operation i: $1 - x, -y, 1 - z$], methyl-C–H...N(hydrazine) [$C4-H4c \cdots N2^i$: $H4c \cdots N2^i = 2.56$ Å, $C4 \cdots N2^i = 3.453(2)$ Å, angle = 151°] and acetonitrile-C–H...S [$C22-H22a \cdots S2^{ii}$: $H22a \cdots S2^{ii} = 2.84$ Å, $C22 \cdots S2^{ii} = 3.792(2)$ Å and angle = 164° and $C22-H22b \cdots S2^{iii}$: $H22b \cdots S2^{iii} = 2.86$ Å, $C22 \cdots S2^{iii} = 3.641(2)$ Å and angle = 137° for ii: $x, y, 1 + z$ and iii: $1 - x, 1 - y,$

$1 - z$] interactions. The latter contacts prove the crucial role of the N5-acetonitrile solvent molecule in assembling the chains and lead to the formation of centrosymmetric eight-membered $\{\cdots S \cdots HCH\}_2$ synthons. However, no specific role is noted for the second independent N6-acetonitrile molecule thus also explaining why this molecule is disordered in the crystal.

Acknowledgements: Sunway University is thanked for supporting crystal engineering studies of metal 1,1-dithiolates.

References

1. Agilent Technologies: CrysAlis PRO. Agilent Technologies, Santa Clara, CA, USA (2017).
2. Sheldrick, G. M.: A short history of SHELX. *Acta Crystallogr. A* **64** (2008) 112–122.
3. Sheldrick, G. M.: Crystal structure refinement with SHELXL. *Acta Crystallogr. C* **71** (2015) 3–8.
4. Farrugia, L. J.: WinGX and ORTEP for Windows: an update. *J. Appl. Crystallogr.* **45** (2012) 849–854.
5. Tiekink, E. R. T.: Exploring the topological landscape exhibited by binary zinc-triad 1,1-dithiolates. *Crystals* **8** (2018) 292.
6. Tiekink, E. R. T.: Perplexing coordination behaviour of potentially bridging bipyridyl-type ligands in the coordination chemistry of zinc and cadmium 1,1-dithiolate compounds. *Crystals* **8** (2018) 18.
7. Lai, C. S.; Liu, T.; Tiekink, E. R. T.: Steric control over polymer formation and topology in adducts of zinc dithiophosphate formed with bridging bipyridine ligands. *CrystEngComm* **6** (2004) 221–226.
8. Chen, D. Lai, C. S.; Tiekink, E. R. T.: Supramolecular aggregation in diimine adducts of zinc(II) dithiophosphates: controlling the formation of monomeric, dimeric, polymeric (zig-zag and helical), and 2-D motifs. *CrystEngComm* **8** (2006) 51–58.
9. Chen, D. Lai, C. S.; Tiekink, E. R. T.: *catena*-Poly[[bis(O, O' -dicyclohexyldithiophosphato- κ^2, S, S')zinc(II)]- μ -1,2-bis(4-pyridylmethylene)hydrazine- $\kappa^2 N: N'$]. *Acta Crystallogr. E* **61** (2005) m2052–m2054.
10. Tan, Y. S.; Tiekink, E. R. T.: Crystal structure of *catena*-poly[(μ -4,4'-bipyridyl- N, N')-bis(O, O' -dimethyldithiophosphato- S)-zinc(II)], $\{C_{14}H_{20}N_2O_4P_2S_4Zn\}_n$. *Z. Kristallogr. NCS* **234** (2019) doi: 10.5115/ncrs-2019-0085.
11. Tiekink, E. R. T.; Wardell, J. L.; Welte, W. B.: *catena*-Poly[[bis(O, O' -diethyl dithiophosphato- $\kappa^2 S, S'$)zinc(II)]- μ -1,2-di-4-pyridylethane- $\kappa^2 N: N'$]. *Acta Crystallogr. E* **63** (2007) m818–m820.
12. Tiekink, E. R. T.; Wardell, J. L.; Welte, W. B.: *catena*-Poly[[bis(O, O' -diethyl dithiophosphato- $\kappa^2 S, S'$)zinc(II)]- μ -1,2-di-4-pyridylethane- $\kappa^2 N: N'$]. *Acta Crystallogr. E* **63** (2007) m818–m820.