Mohammad Hayal Alotaibi, Bakr F. Abdel-Wahab, Obaid Fahad Aldosari, Amany S. Hegazy, Benson M. Kariuki* and Gamal A. El-Hiti*

The crystal structure of *N*-(7-(4-fluorobenzylidene)-3-(4-fluorophenyl)-3,3*a*,4,5,6,7-hexahydro-2*H*-indazole-2-carbonothioyl)benzamide, $C_{28}H_{23}F_2N_3OS$

$V = 4773.8(4) \text{ \AA}^3$, $Z = 8$, $R_{\text{gt}}(F) = 0.0489$, $wR_{\text{ref}}(F^2) = 0.1543$, $T = 296(2) \text{ K}$.

CCDC no.: 1920669

The crystal structure is shown in the figure. Table 1 contains crystallographic data and Table 2 contains the list of the atoms including atomic coordinates and displacement parameters.

Table 1: Data collection and handling.

Crystal:	Colourless block
Size:	$0.27 \times 0.26 \times 0.12 \text{ mm}$
Wavelength:	Mo $K\alpha$ radiation ($0.71073 \text{ \AA}$)
μ :	0.18 mm^{-1}
Diffractometer, scan mode:	SuperNova, ω -scans
θ_{max} , completeness:	29.7° , $>93\%$ (up to 25.2° , $>99\%$)
$N(hkl)_{\text{measured}}$, $N(hkl)_{\text{unique}}$, R_{int} :	39559, 6270, 0.032
Criterion for I_{obs} , $N(hkl)_{\text{gt}}$:	$I_{\text{obs}} > 2 \sigma(I_{\text{obs}})$, 3722
$N(\text{param})_{\text{refined}}$:	264
Programs:	CrysAlis ^{PRO} [1], SHELX [2, 3], WinGX and ORTEP [4]

<https://doi.org/10.1515/ncrs-2019-0281>

Received April 17, 2019; accepted June 4, 2019; available online July 19, 2019

Abstract

$C_{28}H_{23}F_2N_3OS$, monoclinic, $I2/a$ (no. 15), $a = 20.3481(8) \text{ \AA}$, $b = 10.2647(4) \text{ \AA}$, $c = 23.6975(11) \text{ \AA}$, $\beta = 105.317(5)^\circ$,

*Corresponding authors: Benson M. Kariuki, School of Chemistry, Cardiff University, Main Building, Park Place, Cardiff CF10 3AT, UK, e-mail: kariukib@cardiff.ac.uk; and Gamal A. El-Hiti, Department of Optometry, College of Applied Medical Sciences, King Saud University, P.O. Box 10219, Riyadh 11433, Saudi Arabia, e-mail: gelhiti@ksu.edu.sa

Mohammad Hayal Alotaibi: National Center for Petrochemicals Technology, King Abdulaziz City for Science and Technology, P.O. Box 6086, Riyadh 11442, Saudi Arabia

Bakr F. Abdel-Wahab: Department of Chemistry, College of Science and Humanities, Shaqra University, Duwadimi, Saudi Arabia; and Applied Organic Chemistry Department, National Research Centre, Dokki, Giza, Egypt

Obaid Fahad Aldosari: Department of Chemistry, College of Science and Human Studies at Hautat Sudair, Majmaah University, P.O. Box 66, 11952 Majmaah, Saudi Arabia; and Chemistry Department, College of Science and Humanities, Prince Sattam bin Abdulaziz University, P.O. Box 83, 11942 Alkhajj, Saudi Arabia

Amany S. Hegazy: School of Chemistry, Cardiff University, Main Building, Park Place, Cardiff CF10 3AT, UK

Source of materials

The title compound was synthesized from reaction of an equimolar mixture of 7-(4-fluorobenzylidene)-3-(4-fluorophenyl)-3,3*a*,4,5,6,7-hexahydro-2*H*-indazole and benzoyl isothiocyanate in anhydrous ethanol under reflux for 2 h. The crude product was recrystallized from dimethylformamide to give colourless crystals in 82% yield (Mp. $246\text{--}248^\circ\text{C}$).

Experimental details

All hydrogen atoms were placed in calculated positions and refined using a riding model. The N–H bond was fixed at $0.86 \text{ \AA}$ (AFIX 43 instruction in SHELXL [2, 3]), with displacement parameters set to 1.2 times $U_{\text{eq}}(\text{N})$. C–H distances for sp^2 carbon atoms were set to $0.93 \text{ \AA}$ (AFIX 43) and $U_{\text{iso}}(\text{H})$ set to 1.2 times $U_{\text{eq}}(\text{C})$. The methine C–H distance was set to $0.98 \text{ \AA}$ (AFIX 13) and $U_{\text{iso}}(\text{H})$ set to 1.2 times $U_{\text{eq}}(\text{C})$. The methylene C–H distances were set to $0.97 \text{ \AA}$ (AFIX 23) and $U_{\text{iso}}(\text{H})$ set to 1.2 times $U_{\text{eq}}(\text{C})$. The phenyl ring is disordered and was refined with restrained geometry to form regular hexagons and restrained displacement parameters

Table 2: Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²).

Atom	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> _{iso} */ <i>U</i> _{eq}
C1	0.29286(13)	−0.5248(3)	0.24724(15)	0.1024(9)
C2	0.30439(13)	−0.3979(3)	0.25847(12)	0.0997(8)
H2	0.303263	−0.362985	0.294425	0.120*
C3	0.31807(12)	−0.3193(3)	0.21515(11)	0.0848(6)
H3	0.326086	−0.230924	0.222449	0.102*
C4	0.32004(10)	−0.3690(2)	0.16194(10)	0.0709(5)
C5	0.30884(14)	−0.5001(2)	0.15249(13)	0.0966(8)
H5	0.310596	−0.536500	0.116986	0.116*
C6	0.29482(16)	−0.5791(3)	0.19591(17)	0.1128(10)
H6	0.286958	−0.667787	0.189478	0.135*
C7	0.33851(10)	−0.28349(19)	0.11630(9)	0.0673(5)
H7	0.334939	−0.333695	0.080434	0.081*
C8	0.41031(9)	−0.2236(2)	0.13749(9)	0.0687(5)
H8	0.426561	−0.237276	0.179833	0.082*
C9	0.39820(9)	−0.08113(19)	0.12669(9)	0.0656(5)
C10	0.45453(9)	0.0119(2)	0.13546(9)	0.0672(5)
C11	0.52028(10)	−0.0441(2)	0.12734(11)	0.0796(6)
H11A	0.558128	0.000816	0.153653	0.095*
H11B	0.522371	−0.025831	0.087694	0.095*
C12	0.52988(10)	−0.1910(2)	0.13815(11)	0.0801(6)
H12A	0.566937	−0.221034	0.122779	0.096*
H12B	0.542544	−0.207232	0.179959	0.096*
C13	0.46581(10)	−0.2688(2)	0.10973(10)	0.0767(6)
H13A	0.452123	−0.253247	0.067875	0.092*
H13B	0.474220	−0.361245	0.116397	0.092*
C14	0.22965(9)	−0.16202(18)	0.07109(9)	0.0630(5)
C15	0.14029(8)	0.00507(18)	0.03167(9)	0.0616(5)
C16 ^a	0.1329(3)	0.1480(3)	0.0181(3)	0.061(2)
C17 ^a	0.0671(2)	0.1981(6)	0.0045(4)	0.064(2)
H17 ^a	0.030541	0.143861	0.004547	0.076*
C18 ^a	0.05606(16)	0.3294(6)	−0.0090(3)	0.081(4)
H18 ^a	0.012057	0.362997	−0.018094	0.097*
C19 ^a	0.1107(2)	0.4106(4)	−0.0090(2)	0.0849(16)
H19 ^a	0.103339	0.498396	−0.018125	0.102*
C20 ^a	0.17651(17)	0.3604(3)	0.0045(2)	0.0951(17)
H20 ^a	0.213106	0.414661	0.004487	0.114*
C21 ^a	0.18759(17)	0.2291(3)	0.0181(2)	0.0785(14)
H21 ^a	0.231592	0.195525	0.027129	0.094*
C16A ^b	0.1313(3)	0.1489(5)	0.0294(4)	0.0619(19)
C17A ^b	0.0700(3)	0.1954(8)	−0.0060(5)	0.067(2)
H17A ^b	0.038356	0.137217	−0.027968	0.080*
C18A ^b	0.0557(3)	0.3266(8)	−0.0090(4)	0.081(4)
H18A ^b	0.014729	0.356630	−0.033024	0.097*
C19A ^b	0.1018(3)	0.4130(5)	0.0235(3)	0.0840(17)
H19A ^b	0.091479	0.501391	0.022123	0.101*
C20A ^b	0.1631(2)	0.3696(4)	0.0580(3)	0.0930(19)
H20A ^b	0.194948	0.428809	0.078834	0.112*
C21A ^b	0.1774(2)	0.2374(4)	0.0616(3)	0.0762(15)
H21A ^b	0.218433	0.208039	0.085959	0.091*
C22	0.44478(10)	0.1351(2)	0.15006(9)	0.0687(5)
H22	0.400694	0.153165	0.152123	0.082*
C23	0.49145(10)	0.2462(2)	0.16333(8)	0.0669(5)
C24	0.47536(12)	0.3456(2)	0.19732(9)	0.0787(6)
H24	0.435538	0.339254	0.209383	0.094*
C25	0.51670(14)	0.4534(2)	0.21371(11)	0.0895(7)

Table 2 (continued)

Atom	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> _{iso} */ <i>U</i> _{eq}
H25	0.505255	0.518707	0.236592	0.107*
C26	0.57466(13)	0.4613(2)	0.19542(11)	0.0864(7)
C27	0.59160(11)	0.3698(2)	0.16056(11)	0.0842(6)
H27	0.630873	0.379194	0.147888	0.101*
C28	0.55015(10)	0.2624(2)	0.14399(10)	0.0762(6)
H28	0.561429	0.200044	0.119646	0.091*
N1	0.29616(7)	−0.16351(15)	0.10253(7)	0.0657(4)
N2	0.33473(7)	−0.04890(15)	0.10742(7)	0.0660(4)
N3	0.20597(7)	−0.03631(15)	0.05803(7)	0.0664(4)
H3A	0.236086	0.024224	0.067574	0.080*
O1	0.09265(7)	−0.06788(14)	0.01683(8)	0.0878(5)
S1	0.18611(3)	−0.29703(5)	0.05185(3)	0.0797(2)
F1	0.27951(10)	−0.6031(2)	0.28933(9)	0.1500(8)
F2	0.61599(8)	0.56611(15)	0.21233(8)	0.1208(6)

Occupancies: ^a = 0.521(5), ^b = 0.479(5).

(DFIX, SIMU) to give final occupancies of 0.479(5) and 0.521(5) for the two components.

Discussion

Many pyrazole derivatives have been synthesized and found to display antimicrobial, antiviral, anti-inflammatory, insecticidal and herbicidal activities [5–8]. The structure of the title compound has been obtained as part of a study of these materials [9].

The asymmetric unit consists of one molecule of the title compound (see the figure). The molecule consists of two fluorophenyl groups [**A** (F1, C1–C6) and **D** (F2, C23–C28)], a hexahydroindazolyl group [**B** (N1, N2, C7–C13)] and a phenyl group [**C** (C16–C21)]. Ring **C** is disordered by a rotation of 35.3(3)° about the C15–C16 bond. The cyclohexane segment (C8–C13) of **B** assumes a chair conformation. The pyrazolyl ring of **B** is planar with a maximum deviation of 0.037(3) Å from the least squares plane of the ring. Related structures include *cis*-1-benzoyl-2-chloroacetyl-4,5-hexamethylenepyrazolidine [10] and 2-benzyl 3-ethyl 6a-methyl 1-(benzoylcarbamothioyl)hexahydrocyclopenta[c]pyrazole-2,3,6a(1H)-tricarboxylate [11] in which the pyrazolyl rings are in envelope conformation. An intramolecular N–H...N contact is observed (N3...N2 distance of 2.575(2) Å; N3–H3A...N2 angle of 112.8°). Two molecules related by inversion symmetry are linked by weak non-classical C–H...O interactions (C17...O1 distance of 3.427(3) Å and C17–H17...O1 angle of 158.4°) to form R²₂(10) rings. Comparably weak C–H...S interactions also occur in the structure with C27...S1 distances of 3.676(3) Å and C27–H27...S1 angles of 139.7°.

Acknowledgements: The authors would like to thank the Deanship of Scientific Research at Majmaah University for

supporting this work under Project Number No. 1440-127, and to Cardiff University for continued support.

References

1. Rigaku Oxford Diffraction. CrysAlis PRO. Rigaku Oxford Diffraction, Yarnton, England (2015).
2. Sheldrick, G. M.: A short history of SHELX. *Acta Crystallogr. A* **64** (2008) 112–122.
3. Sheldrick, G. M.: Crystal structure refinement with SHELXL. *Acta Cryst. C* **71** (2015) 3–8.
4. Farrugia, L. J.: WinGX and ORTEP for Windows: an update. *J. Appl. Crystallogr.* **45** (2012) 849–854.
5. Naim, M. J.; Alam, O.; Nawaz, F.; Alam, M. J.; Alam, P.: Current status of pyrazole and its biological activities. *J. Pharm. Bioallied Sci.* **8** (2016) 2–17.
6. Sayed, G. H.; Azab, M. E.; Anwer, K. E.; Abdel Raouf, M.; Negm, N. A.: Pyrazole, pyrazolone and enamionitrile pyrazole derivatives: Synthesis, characterization and potential in corrosion inhibition and antimicrobial applications. *J. Mol. Liq.* **252** (2018) 329–338.
7. Sidduri, A.; Budd, D. C.; Fuentes, M. E.; Lambros, T.; Ren, Y.; Roongta, V.; Schoenfeld, R. C.; Gillespie, P.; Stevenson, C. S.; Truitt, T.; Qian, Y.: Discovery of novel non-carboxylic acid 5-amino-4-cyanopyrazole derivatives as potent and highly selective LPA1R antagonists. *Bioorg. Med. Chem. Lett.* **24** (2014) 4450–4454.
8. Faria, J. V.; Vegi, P. F.; Miguita, A. G. C.; Dos Santos, M. S.; Boechat, N.; Bernardino, A. M. R.: Recently reported biological activities of pyrazole compounds. *Bioorg. Med. Chem.* **25** (2017) 5891–5903.
9. Alotaibi, M. H.; Hanan, M. A.; Bakr, F. A.-W.; Amany, H. S.; Kariuki, B. M.; El-Hiti, G. A.: Crystal structure of 5-(5-(4-chlorophenyl)-1-phenyl-1*H*-pyrazol-3-yl)-*N*-phenyl-2-amine, $C_{23}H_{16}ClN_5O$. *Z. Kristallogr. NCS* **234** (2019) 543–545.
10. Kapor, A.; Rakic, S.; Stancic, M.; Simon, L.; Talpas, G. S.; Bernath, G.; Engel, P.: Structure and conformation of *cis*-benzoyl-2-chloroacetyl-4,5-hexamethylenepyrazolidine: an x-ray study. *J. Mol. Struct.* **380** (1996) 85–91.
11. Gergely, J.; Morgan, J. B.; Overman, L. E.: Stereocontrolled synthesis of functionalized *cis*-cyclopentapyrazolidines by 1,3-dipolar cycloaddition reactions of azomethine imines. *J. Org. Chem.* **71** (2006) 9144–9152.