See Mun Lee, Kong Mun Lo and Edward R.T. Tiekink*

Crystal structure of (2,2'-bipyridyl)bis(4-chlorobenzyl)dichloridotin(IV), $C_{24}H_{20}Cl_4N_2Sn$

$V = 2420.92(4) \text{ \AA}^3$, $Z = 4$, $R_{\text{gt}}(F) = 0.0229$, $wR_{\text{ref}}(F^2) = 0.0603$, $T = 100(2) \text{ K}$.

CCDC no.: 1948401

The molecular structure is shown in the figure. Table 1 contains crystallographic data and Table 2 contains the list of the atoms including atomic coordinates and displacement parameters.

Table 1: Data collection and handling.

Crystal:	Colourless prism
Size:	$0.14 \times 0.09 \times 0.04 \text{ mm}$
Wavelength:	$\text{Cu K}\alpha$ radiation ($1.54178 \text{ \AA}$)
μ :	12.6 mm^{-1}
Diffractometer, scan mode:	XtaLAB Synergy, ω
θ_{max} , completeness:	67.1° , $>99\%$
$N(hkl)_{\text{measured}}$, $N(hkl)_{\text{unique}}$, R_{int} :	30103, 4329, 0.042
Criterion for I_{obs} , $N(hkl)_{\text{gt}}$:	$I_{\text{obs}} > 2 \sigma(I_{\text{obs}})$, 3987
$N(\text{param})_{\text{refined}}$:	280
Programs:	CrysAlis ^{PRO} [1], SHELX [2, 3], WinGX/ORTEP [4]

Source of material

The melting point of the compound was measured on a Melt-Temp II digital melting point apparatus and was uncorrected. The IR spectrum was measured on a Bruker Vertex 70v FTIR spectrophotometer from 4000 to 400 cm^{-1} . The ^1H and $^{13}\text{C}\{^1\text{H}\}$ NMR spectra were recorded in CDCl_3 solution on a Bruker Ascend 400 MHz NMR spectrometer with chemical shifts relative to tetramethylsilane.

Di(4-chlorobenzyl)tin dichloride was synthesized by the direct reaction of 4-chlorobenzyl chloride (Aldrich) and metallic tin powder (Merck) in toluene according to a literature procedure [5]. Di(4-chlorobenzyl)tin dichloride (0.44 g , 1.0 mmol) and 2,2'-bipyridine (Aldrich, 0.16 g , 1.0 mmol) were heated in 95% ethanol (30 mL) and stirred for 2 h. After filtration, the filtrate was evaporated slowly until colourless crystals were formed. Yield: 0.31 g (51.9%). **M.pt.** $471\text{--}473 \text{ K}$. IR (cm^{-1}): 1606 (m) $\nu(\text{C--N})$, 1483 , 1141 (s) $\nu(\text{C--N})$, 481 (m) $\nu(\text{Sn--N})$. $^1\text{H NMR}$ (CDCl_3 , ppm): δ $3.17 \text{ (s, 2H, Ph--CH}_2\text{)}$, $6.55\text{--}6.56 \text{ (dd, 2H, Ph--H)}$, $6.57\text{--}6.58 \text{ (dd, 2H, Ph--H)}$, $7.50\text{--}7.73 \text{ (m, 1H, bipy-H)}$, $7.86\text{--}7.88 \text{ (d, 1H, bipy-H)}$, $8.03\text{--}8.07 \text{ (t, 1H, bipy-H)}$.

<https://doi.org/10.1515/ncrs-2019-0508>

Received July 18, 2019; accepted August 21, 2019; available online September 9, 2019

Abstract

$C_{24}H_{20}Cl_4N_2Sn$, monoclinic, $P2_1/n$ (no. 14), $a = 7.7944(1) \text{ \AA}$, $b = 21.1245(2) \text{ \AA}$, $c = 14.7611(1) \text{ \AA}$, $\beta = 95.077(1)^\circ$,

*Corresponding author: Edward R.T. Tiekink, Research Centre for Crystalline Materials, School of Science and Technology, Sunway University, 47500 Bandar Sunway, Selangor Darul Ehsan, Malaysia, e-mail: edwardt@sunway.edu.my. <https://orcid.org/0000-0003-1401-1520>

See Mun Lee and Kong Mun Lo: Research Centre for Crystalline Materials, School of Science and Technology, Sunway University, 47500 Bandar Sunway, Selangor Darul Ehsan, Malaysia

Table 2: Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²).

Atom	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> _{iso} [*] / <i>U</i> _{eq}
Sn	0.56304(2)	0.69771(2)	0.57470(2)	0.02011(6)
Cl1	0.65370(8)	0.76490(3)	0.71076(4)	0.03300(14)
Cl2	0.46362(8)	0.59790(3)	0.64596(4)	0.03040(14)
Cl3	0.98078(10)	0.50596(4)	0.25670(5)	0.0559(2)
Cl4	0.16551(14)	0.98789(4)	0.39746(5)	0.0602(2)
N1	0.6389(2)	0.77422(9)	0.47038(12)	0.0220(4)
N2	0.4981(3)	0.66045(9)	0.42577(13)	0.0237(4)
C1	0.8256(3)	0.66250(12)	0.57962(17)	0.0293(5)
H1A	0.9052	0.6991	0.5846	0.035*
H1B	0.8487	0.6368	0.6354	0.035*
C2	0.8653(3)	0.62366(12)	0.50031(18)	0.0274(5)
C3	0.9441(3)	0.65035(13)	0.42807(17)	0.0295(6)
H3	0.9741	0.6939	0.4302	0.035*
C4	0.9793(3)	0.61458(14)	0.35337(18)	0.0347(6)
H4	1.0346	0.6332	0.3051	0.042*
C5	0.9331(3)	0.55163(14)	0.35000(19)	0.0361(6)
C6	0.8530(3)	0.52352(14)	0.4196(2)	0.0403(7)
H6	0.8211	0.4801	0.4163	0.048*
C7	0.8204(3)	0.55973(13)	0.4940(2)	0.0357(6)
H7	0.7659	0.5406	0.5421	0.043*
C8	0.3035(3)	0.73429(11)	0.57325(16)	0.0245(5)
H8A	0.2244	0.7033	0.5412	0.029*
H8B	0.2731	0.7367	0.6369	0.029*
C9	0.2704(3)	0.79740(11)	0.53029(16)	0.0234(5)
C10	0.2932(3)	0.85329(12)	0.58044(16)	0.0270(5)
H10	0.3307	0.8510	0.6434	0.032*
C11	0.2624(4)	0.91181(13)	0.54036(17)	0.0332(6)
H11	0.2767	0.9495	0.5753	0.040*
C12	0.2103(4)	0.91449(12)	0.44805(18)	0.0338(6)
C13	0.1897(3)	0.86048(13)	0.39608(16)	0.0293(5)
H13	0.1553	0.8632	0.3328	0.035*
C14	0.2197(3)	0.80247(11)	0.43712(17)	0.0245(5)
H14	0.2058	0.7651	0.4015	0.029*
C15	0.7005(3)	0.83109(12)	0.49631(17)	0.0290(5)
H15	0.7154	0.8406	0.5594	0.035*
C16	0.7434(4)	0.87653(13)	0.4348(2)	0.0368(6)
H16	0.7898	0.9162	0.4551	0.044*
C17	0.7174(4)	0.86304(13)	0.3438(2)	0.0408(7)
H17	0.7455	0.8935	0.3000	0.049*
C18	0.6503(4)	0.80501(12)	0.31598(19)	0.0339(6)
H18	0.6292	0.7957	0.2530	0.041*
C19	0.6136(3)	0.76016(11)	0.38062(15)	0.0227(5)
C20	0.5476(3)	0.69619(11)	0.35705(16)	0.0219(5)
C21	0.5388(3)	0.67292(13)	0.26900(16)	0.0301(5)
H21	0.5729	0.6985	0.2208	0.036*
C22	0.4795(3)	0.61185(14)	0.25209(17)	0.0336(6)
H22	0.4758	0.5949	0.1923	0.040*
C23	0.4262(3)	0.57587(13)	0.32175(18)	0.0340(6)
H23	0.3828	0.5343	0.3110	0.041*
C24	0.4377(3)	0.60198(12)	0.40811(18)	0.0307(6)
H24	0.4012	0.5774	0.4568	0.037*

bipy-H), 9.09–9.11 (d, 1H, bipy-H). ¹³C{¹H} NMR (CDCl₃, ppm): δ 50.4 (CH₂), 127.3, 129.1, 129.8, 138.2 (Ph–C), 121.0, 126.6, 127.0, 140.6, 148.5 (bipy-C).

Experimental details

The C-bound H atoms were geometrically placed (C–H = 0.95–0.99 Å) and refined as riding with *U*_{iso}(H) = 1.2*U*_{eq}(C).

Comment

In connection with the recent report of the crystal and molecular structures of (4-BrC₆H₄)₂SnBr₂(2,2'-bipyridyl) [6], herein the synthesis and structure of the all-chloro derivative, (I), are described. The molecule of (I) is illustrated in the figure (50% displacement ellipsoids). A distorted octahedral geometry for the tin atom is noted which is based on a C₂Cl₂N₂ donor set defined by two carbon atoms of the 4-chloro-substituted benzyl groups, two chloride atoms and two nitrogen atoms of the 2,2'-bipyridyl molecule. The benzyl-carbon atoms are mutually trans, and each of the chloride atoms is trans to a pyridyl-nitrogen atom. The Sn–Cl bond lengths are experimentally equivalent [Sn–Cl1 = 2.5093(6) Å and Sn–Cl2 = 2.5094(6) Å] as are the Sn–N bonds [Sn–N1 = 2.3439(19) Å and Sn–N2 = 2.3478(19) Å]. This contrasts the situation found in the all-bromo derivative where non-systematic differences in tin-ligand bond lengths were noted [6]. Such variations have been proven by DFT methods on related compounds to be dependent upon global molecular packing effects in crystals [7, 8].

In (I), the five-membered chelate ring formed by the 2,2'-bipyridyl ligand is responsible for the greatest distortion from the ideal octahedral geometry with the acute N1–Sn–N2 chelate angle being 70.25(7)°. The maximum deviation of a trans angles is seen in the N1–Sn–Cl2 angle of 163.73(5)°. As reported for the all-bromo derivative [6], the benzyl-phenyl rings are orientated to lie over the 2,2'-bipyridyl molecule. In the present case, the separation between the ring centroids of the chelate and C2-phenyl rings is 3.7784(14) Å; the separation between the N1-pyridyl and C9-phenyl rings is marginally longer at 3.7957(15) Å. Recent literature has highlighted the stabilising effect of π ··· π interactions where at least one of the rings is a chelate ring [9, 10].

The structure of the (4-ClC₆H₄CH₂)₂SnCl₂ precursor molecule is available in the literature for comparison [11]. The independent Sn–Cl bond length in this 2-fold symmetric structure is 2.3754(11) Å which is considerably shorter than those reported above for the title compound, reflecting, in part, the increased coordination number for tin.

While the precursor (4-XC₆H₄CH₂)₂SnX₂, X = Cl [11] and Br [12] species are isostructural, those of (4-XC₆H₄CH₂)₂SnX₂(2,2'-bipyridyl), X = Cl and Br [6], are not. However, the title compound appears to be isostructural with the 1,10-phenanthroline analogue [13]. The non-isostructural relationship between the (4-XC₆H₄CH₂)₂SnX₂(2,2'-bipyridyl), X = Cl and Br [6], compounds is reflected in the distinctive atom-to-atom points of contact between molecules.

Based on the distance criteria in PLATON [14], the only significant intermolecular interactions in the crystal are benzyl- and pyridyl- $C-H \cdots Cl$ (tin- and phenyl-bound) interactions [$C4-H4 \cdots Cl1^i$: $H4 \cdots Cl1^i = 2.77 \text{ \AA}$, $C4 \cdots Cl1^i = 3.642(3) \text{ \AA}$ with angle at $H4 = 153^\circ$, and $C22-H22 \cdots Cl4^{ii}$: $H22 \cdots Cl4^{ii} = 2.80 \text{ \AA}$, $C22 \cdots Cl4^{ii} = 3.545(3) \text{ \AA}$ with angle at $H22 = 136^\circ$ for symmetry operations (i) $1/2 + x, 3/2 - y, -1/2 + z$ and (ii) $1/2 - x, -1/2 + y, 1/2 - z$]. The molecules pack in columns aligned along the a -axis direction and with each molecule accepting and donating two $C-H \cdots Cl$ interactions, a three-dimensional architecture ensues.

Finally, an analysis of the calculated Hirshfeld surfaces was undertaken in order to evaluate the molecular packing further and to make a comparison with the packing in the crystal of the all-bromo analogue [6]. For this purpose, Crystal Explorer 17 [15] was employed following protocols established in the literature [16] to calculate the Hirshfeld surfaces and two-dimensional fingerprint plots (overall and delineated into specific contacts). In the crystal, $H \cdots H$ contacts make the greatest overall contribution at 42.3%, being greater than $H \cdots Cl/Cl \cdots H$ contacts at 38.7%. Other significant contacts are noted, that is, $H \cdots C/C \cdots H$ [12.4%], then $Cl \cdots C/C \cdots Cl$ [3.0%], $Cl \cdots Cl$ [2.2%] and $C \cdots C$ [2.0%]. A very similar pattern in surface contacts were calculated for the all-bromo derivative with the notable exception that $H \cdots Br/Br \cdots H$ [40.8%] were marginally more prevalent than $H \cdots H$ [40.2%] contacts [6].

Acknowledgements: Sunway University Sdn Bhd is thanked for financial support of this work through Grant no. STR-RCTR-RCCM-001-2019.

References

1. Rigaku Oxford Diffraction: CrysAlis^{PRO}. Rigaku Corporation, Oxford, UK (2018).
2. Sheldrick, G. M.: A short history of SHELX. *Acta Crystallogr. A* **64** (2008) 112–122.
3. Sheldrick, G. M.: Crystal structure refinement with SHELXL. *Acta Crystallogr. C* **71** (2015) 3–8.
4. Farrugia, L. J.: WinGX and ORTEP for Windows: an update. *J. Appl. Crystallogr.* **45** (2012) 849–854.
5. Sisido, K.; Takeda, Y.; Kinugawa, Z.: Direct synthesis of organotin compounds I. di- and tribenzyltin chlorides. *J. Am. Chem. Soc.* **83** (1961) 538–541.
6. Lee, S. M.; Lo, K. M.; Tiekink, E. R. T.: Crystal structure of (2,2'-bipyridyl)bis(4-bromobenzyl)dibromidotin(IV), $C_{24}H_{20}Br_4N_2Sn$. *Z. Kristallogr. NCS* **234** (2019) submitted, doi: org/10.1515/ncrs-2019-0507.
7. Buntine, M. A.; Hall, V. J.; Kosovel, F. J.; Tiekink, E. R. T.: Influence of crystal packing on molecular geometry: a crystallographic and theoretical investigation of selected diorganotin systems. *J. Phys. Chem. A* **102** (1998) 2472–2482.
8. Buntine, M. A.; Hall, V. J.; Tiekink, E. R. T.: The crystal and molecular structures of $R_2SnCl_2(1,10\text{-phenanthroline})$, $R = iPr$, Cy , CH_2Ph and $R_2 = Me$, Ph : a comparison between solid state and theoretical structures. *Z. Kristallogr.* **213** (1998) 669–678.
9. Malenov, D. P.; Janjić, G. V.; Medaković, V. B.; Hall, M. B.; Zarić, S. D.: Noncovalent bonding: stacking interactions of chelate rings of transition metal complexes. *Coord. Chem. Rev.* **345** (2017) 318–341.
10. Tiekink, E. R. T.: Supramolecular assembly based on “emerging” intermolecular interactions of particular interest to coordination chemists. *Coord. Chem. Rev.* **345** (2017) 209–228.
11. Kuang, D.-Z.; Feng, Y.-L.: Synthesis and crystal structure of bis(p -chlorobenzyl)tin dichloride. *Chin. J. Inorg. Chem.* **16** (2000) 603–606.
12. Lee, S. M.; Lo, K. M.; Tiekink, E. R. T.: Crystal structure of dibromidobis(4-bromobenzyl)tin(IV), $C_{14}H_{12}Br_4Sn$. *Z. Kristallogr. NCS* **234** (2019) 947–948.
13. Tan, C. L.; Lo, K. M.; Ng, S. W.: Dichloridobis(4-chloro-benzyl- κC)(1,10-phenanthroline- $\kappa^2 N, N'$)tin(IV). *Acta Crystallogr. E* **65** (2009) m717.
14. Spek, A. L.: Structure validation in chemical crystallography. *Acta Crystallogr. D* **65** (2009) 148–155.
15. Turner, M. J.; McKinnon, J. J.; Wolff, S. K.; Grimwood, D. J.; Spackman, P. R.; Jayatilaka, D.; Spackman, M. A.: Crystal Explorer v17. The University of Western Australia, Australia (2017).
16. Tan, S. L.; Jotani, M. M.; Tiekink, E. R. T.: Utilizing Hirshfeld surface calculations, non-covalent interaction (NCI) plots and the calculation of interaction energies in the analysis of molecular packing. *Acta Crystallogr. E* **75** (2019) 308–318.