Yee Seng Tan and Edward R.T. Tiekink*

Crystal structure of *catena*-poly[(μ_2 -1,2-bis(3-pyridylmethylene)hydrazine- $\kappa^2N:N'$)-bis(*O,O'*-dimethyl dithiophosphato- κ^2 -*S,S'*)cadmium(II)], {C₁₆H₂₂CdN₄O₄P₂S₄}_n

Abstract

C₁₆H₂₂CdN₄O₄P₂S₄, orthorhombic, *Pbca* (no. 61), *a* = 12.66410(10) Å, *b* = 13.00140(10) Å, *c* = 15.30720(10) Å, *V* = 2520.35(3) Å³, *Z* = 4, *R*_{gt}(*F*) = 0.0186, *wR*_{ref}(*F*²) = 0.0503, *T* = 100(2) K.

CCDC no.: 1959862

The crystal structure is shown in the figure. Table 1 contains crystallographic data and Table 2 contains the list of the atoms including atomic coordinates and displacement parameters.

Table 1: Data collection and handling.

Crystal:	Yellow prism
Size:	0.11 × 0.11 × 0.06 mm
Wavelength:	Cu <i>K</i> α radiation (1.54184 Å)
<i>μ</i> :	11.51 mm ^{−1}
Diffractometer, scan mode:	XtaLAB Synergy, <i>ω</i> -scans
<i>θ</i> _{max} , completeness:	67°, >99%
<i>N</i> (<i>hkl</i>) _{measured} , <i>N</i> (<i>hkl</i>) _{unique} , <i>R</i> _{int} :	15972, 2247, 0.026
Criterion for <i>I</i> _{obs} , <i>N</i> (<i>hkl</i>) _{gt} :	<i>I</i> _{obs} > 2 <i>σ</i> (<i>I</i> _{obs}), 2144
<i>N</i> (<i>param</i>) _{refined} :	144
Programs:	CrysAlis ^{PRO} [1], SHELX [2, 3], WinGX and ORTEP [4]

Source of material

The Cd[S₂P(OMe)₂]₂ precursor was prepared in high yield from the *in situ* reaction of Cd(NO₃)₂ · 4H₂O (Acro Organic; 15.42 g, 0.05 mol), MeOH (Merck; 8.90 mL, 0.22 mol), P₂S₅ (Sigma-Aldrich; 11.11 g, 0.05 mol) and 50% w/w NaOH solution (Merck; 8.80 mL, 0.11 mol). 1,2-Bis(3-pyridylmethylene)hydrazine was prepared in high yield from the 2:1 reaction (reflux) of 3-picolyamine (Aldrich; 2.04 mL, 0.02 mol) and hydrazinium hydroxide (Merck; 0.49 mL, 0.01 mol) in ethanol solution (Merck; 5 mL). The title compound was obtained by mixing a suspension of Cd[S₂P(OMe)₂]₂ (0.50 g, 1.17 mmol) and 1,2-bis(3-pyridylmethylene)hydrazine (0.25 g, 1.19 mmol) in dimethylformamide (Merck; 5 mL), followed by stirring for 30 min. at 373 K. The solution was filtered and the collected filtrate transferred to a sample vial containing acetonitrile (Merck; 1 mL). Yellow blocks formed after one day. Yield: 0.33 g, (44.3%, based on Cd[S₂P(OMe)₂]₂). **M. pt** (MelTemp Melting Point Apparatus): 431.2–433.2 K. **IR** (Bruker Vertex 70 V equipped with Platinum ATR from 400 to 80 cm^{−1}):

<https://doi.org/10.1515/ncrs-2019-0654>

Received September 5, 2019; accepted October 17, 2019; available online November 6, 2019

*Corresponding author: Edward R.T. Tiekink, Research Centre for Crystalline Materials, School of Science and Technology, Sunway University, 47500 Bandar Sunway, Selangor Darul Ehsan, Malaysia, e-mail: edwardt@sunway.edu.my. <https://orcid.org/0000-0003-1401-1520>

Yee Seng Tan: Research Centre for Crystalline Materials, School of Science and Technology, Sunway University, 47500 Bandar Sunway, Selangor Darul Ehsan, Malaysia

Table 2: Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ($\text{\AA}^2$).

Atom	x	y	z	U_{iso}^*/U_{eq}
Cd	0.500000	0.500000	0.500000	0.01276(8)
S1	0.50422(3)	0.39582(3)	0.65173(3)	0.01638(11)
S2	0.33651(3)	0.58052(3)	0.58891(2)	0.01796(10)
P1	0.38285(3)	0.48585(3)	0.68372(3)	0.01570(11)
O1	0.40685(9)	0.54459(9)	0.77269(7)	0.0217(3)
O2	0.28538(10)	0.42263(9)	0.72038(8)	0.0252(3)
N1	0.40015(10)	0.36017(10)	0.44326(8)	0.0142(3)
N2	0.49313(11)	0.05353(14)	0.49356(9)	0.0162(4)
C1	0.50294(13)	0.60421(15)	0.78070(13)	0.0255(4)
H1A	0.507665	0.632915	0.839765	0.038*
H1B	0.502156	0.660303	0.737940	0.038*
H1C	0.564029	0.559750	0.769809	0.038*
C2	0.22779(17)	0.35511(16)	0.66230(13)	0.0347(5)
H2A	0.166223	0.327039	0.692903	0.052*
H2B	0.273833	0.298631	0.643690	0.052*
H2C	0.204290	0.393915	0.610977	0.052*
C3	0.42940(12)	0.26373(12)	0.46251(9)	0.0137(3)
H3	0.493338	0.253436	0.493771	0.016*
C4	0.37027(13)	0.17807(12)	0.43873(9)	0.0136(3)
C5	0.27606(13)	0.19419(12)	0.39422(11)	0.0183(3)
H5	0.232323	0.137669	0.378782	0.022*
C6	0.24663(13)	0.29370(13)	0.37259(11)	0.0220(4)
H6	0.183340	0.306358	0.341064	0.026*
C7	0.31101(13)	0.37392(12)	0.39774(10)	0.0170(3)
H7	0.291233	0.441953	0.382081	0.020*
C8	0.40368(12)	0.07275(12)	0.45912(10)	0.0146(3)
H8	0.357338	0.017321	0.446280	0.017*

1173(w) $\nu(C-O)$; 1000(m) $\nu(P-O)$; 650 (s) $\nu(P-S)$; 392(m) $\nu(Cd-N)$; 263(w) $\nu(Cd-S)$.

Experimental details

H atoms were geometrically placed ($C-H = 0.95-0.98 \text{ \AA}$) and refined as riding with $U_{iso}(H) = 1.2-1.5U_{eq}(C)$.

Discussion

It is well-documented that very different structures/supramolecular architectures can be observed for the zinc-triad 1,1-dithiolates [5] and their adducts with bipyridyl-type ligands [6] when only small changes in the nature of a remote group are made. This has been very recently highlighted in the structures of $\{Cd[S_2P(OR)_2]_2(\text{pyrazine})\}_n$. Thus, in the case of $R = Et$, an octahedral *trans*- N_2S_4 donor set is found for cadmium(II) within a linear, one-dimensional coordination polymer [7]. By contrast, a *cis*- N_2S_4 donor set is found in the $R = isopropyl$ derivative, characterized as an acetonitrile solvate [8]. While the donor set still defines an octahedral geometry, the *cis*-disposition of the nitrogen donors gives rise to a one-dimensional coordination polymer with a zigzag topology [8]. Such intriguing results

vindicate systematic studies in the field. In keeping with this, herein, the crystal and molecular structures of a new coordination polymer, $\{Cd[S_2P(OEt)_2]_2(3\text{-PyAld})\}_n$, (I), are described; where 3-PyAld is 3-pyridylaldazine molecule, i.e. $3\text{-NC}_5\text{H}_4\text{C(H)=N=N=C(H)C}_5\text{H}_4\text{N-3}$.

The asymmetric unit of (I) comprises half a $Cd[S_2P(OEt)_2]_2$ entity as the cadmium(II) centre is located on a centre of inversion as well as half a 3-PyAld molecule as this is disposed about another centre of inversion. Referring to the upper view of the figure (70% probability displacement ellipsoids), the symmetry related dithiophosphate ligand is generated by the symmetry operation (i) $1 - x, 1 - y, 1 - z$, and the other half of the 3-PyAld ligand is related by (ii) $1 - x, -y, 1 - z$. A *trans*- N_2S_4 donor set is evident for the cadmium(II) atom as the dithiophosphate ligands are symmetrically chelating within a plane defined by the four S atoms. The $Cd-S1, S2 = 2.6892(4)$ and $2.6897(4) \text{ \AA}$ bond lengths are experimentally equivalent. The equivalence of the $Cd-S$ bonds is reflected in the experimental equivalence of the associated $P1-S1, S2$ bond lengths [i.e. $1.9930(6)$ and $1.9915(6) \text{ \AA}$]. The $Cd-N1$ bond length is $2.3788(13) \text{ \AA}$. The coordination geometry is based on an octahedron with the maximum deviation traced to the relatively acute $S1-Cd-S2$ chelate angle of $76.958(11)^\circ$.

As seen in the lower view of the figure, the resultant one-dimensional coordination polymer is linear. This resembles the situation observed for the $R = ethyl$ [9] and the $R = isopropyl$ [10] derivatives. Thus, the four structures of the general formula $\{Cd[S_2P(OEt)_2]_2(3\text{-PyAld})\}_n$, $R = methyl$ (I), *ethyl* [9], *isopropyl* [10, 11] and *cyclohexyl* [12] adopt the same structural motif in the solid-state.

In the crystal, the chains are aligned along $[0 1 0]$. Connections between chains are of the type $pyridyl-C-H \cdots O(\text{alkoxy})$ [$C6-H6 \cdots O1^{iii}$: $H6 \cdots O1^{iii} = 2.48 \text{ \AA}$, $C6 \cdots O1^{iii} = 3.246(2) \text{ \AA}$ with angle at $H6 = 138^\circ$ for (iii) $1/2 - x, 1 - y, -1/2 + z$]. These give rise to a three-dimensional architecture with rather large voids. The voids are occupied by a second three-dimensional architecture so that the structure is 3D-3D doubly interpenetrated.

Acknowledgements: Sunway University Sdn Bhd is thanked for financial support of this work through Grant no. STR-RCTR-RCCM-001-2019.

References

1. Rigaku Oxford Diffraction: CrysAlis PRO. Rigaku Corporation, Oxford, UK (2018).
2. Sheldrick, G. M.: A short history of SHELX. *Acta Crystallogr. A* **64** (2008) 112–122.
3. Sheldrick, G. M.: Crystal structure refinement with SHELXL. *Acta Crystallogr. C* **71** (2015) 3–8.

4. Farrugia, L. J.: WinGX and ORTEP for Windows: an update. *J. Appl. Cryst.* **45** (2012) 849–854.
5. Tiekink, E. R. T.: Exploring the topological landscape exhibited by binary zinc-triad 1,1-dithiolates. *Crystals* **8** (2018) article no. 292 (34 pages).
6. Tiekink, E. R. T.: Perplexing coordination behaviour of potentially bridging bipyridyl-type ligands in the coordination chemistry of zinc and cadmium 1,1-dithiolate compounds. *Crystals* **8** (2018) article no. 18 (29 pages).
7. Tan, Y. S.; Tiekink, E. R. T.: Crystal structure of *catena*-poly[(μ_2 -pyrazine-*N,N'*)-bis(*O,O'*-di-ethyldithiophosphato-*S,S'*)cadmium(II)], $[C_{12}H_{24}CdN_2O_4P_2S_4]_n$. *Z. Kristallogr. NCS* **235** (2020) 319–321.
8. Tan, Y. S.; Tiekink, E. R. T.: Crystal structure of *catena*-poly[(μ_2 -pyrazine-*N,N'*)-bis(*O,O'*-di-isopropyldithiophosphato-*S,S'*)cadmium(II) acetonitrile di-solvate], $[C_{16}H_{32}CdN_2O_4P_2S_4 \cdot 2(C_2H_3N)]_n$. *Z. Kristallogr. NCS* **235** (2020) 323–325.
9. Tan, Y. S.; Tiekink, E. R. T.: Crystal structure of *catena*-poly[(bis(*O,O'*-diethyl dithiophosphato- κ^2S,S')- μ_2 -1,2-bis(3-pyridylmethylene)hydrazine- κ^2N,N')cadmium(II)], $\{C_{20}H_{30}CdN_4O_4P_2S_4\}_n$. *Z. Kristallogr. NCS* **234** (2020) 343–345.
10. Lai, C. S.; Tiekink, E. R. T.: Polymeric topologies in cadmium(II) dithiophosphate adducts of the isomeric *n*-pyridinealdazines, $n = 2, 3$ and 4. *Z. Kristallogr. C. M.* **221** (2006) 288–293.
11. Tan, Y. S.; Tiekink, E. R. T.: A redetermination of the crystal structure of *catena*-poly[(bis(*O,O'*-isopropyl dithiophosphato- κ^2S,S')-(μ_2 -1,2-bis(3-pyridylmethylene)hydrazine- κ^2N,N')cadmium(II)], $\{C_{24}H_{38}CdN_4O_4P_2S_4\}_n$. *Z. Kristallogr. NCS* **235** (2020) 253–255.
12. Lai, C. S.; Tiekink, E. R. T.: Delineating the principles controlling polymer formation and topology in zinc(II)- and cadmium(II)-dithiophosphate adducts of diimine-type ligands. *J. Molec. Struct.* **796** (2006) 114–118.