Guido J. Reiss, Vlada B. Urlacher and U. Joost Luelf*

Enzyme-mediated synthesis and crystal structure of (2*R*,4*S*)-hydroxyketamine, C₁₃H₁₆ClNO₂

Abstract

C₁₃H₁₆ClNO₂, monoclinic, *P*₂₁ (no. 4), *a* = 7.4945(7) Å, *b* = 7.2336(6) Å, *c* = 11.5401(10) Å, β = 92.555(2)°, *V* = 624.99(10) Å³, *Z* = 2, *R*_{gt}(*F*) = 0.0415, *wR*_{ref}(*F*²) = 0.0947, *T* = 291 K.

CCDC no.: 1998665

The molecular structure is shown in the figure. Table 1 contains crystallographic data and Table 2 contains the list of the atoms including atomic coordinates and displacement parameters.

Table 1: Data collection and handling.

Crystal:	Colorless block
Size:	0.90 × 0.80 × 0.56 mm
Wavelength:	Mo Kα radiation (0.71073 Å)
μ:	0.30 mm ^{−1}
Diffractometer, scan mode:	Bruker APEX-II, φ and ω
θ _{max} , completeness:	30.0°, >99%
<i>N</i> (<i>hkl</i>) _{measured} , <i>N</i> (<i>hkl</i>) _{unique} , <i>R</i> _{int} :	6118, 3527, 0.087
Criterion for <i>I</i> _{obs} , <i>N</i> (<i>hkl</i>) _{gt} :	<i>I</i> _{obs} > 2 σ(<i>I</i> _{obs}), 3225
<i>N</i> (<i>param</i>) _{refined} :	164
Programs:	Bruker [1], SHELX [2, 3], Diamond [4]

Source of material

An amount of 360 μmol (98.7 mg) of (*R*)-ketamine hydrochloride was converted to the title compound by an *Escherichia coli*-based whole-cell biocatalyst expressing the double mutant I238G/M388A of the cytochrome P450 monooxygenase CYP154E1 from *Thermobifida fusca* YX and two redox partner proteins as recently described by Bokel et al. [5]. The reaction was carried out in a total volume of 180 mL with an educt concentration of 2 mM. After incubation at 25 °C, 250 revolutions per minute for 20 h, the product mixture was extracted using ethyl acetate after addition of sodium carbonate. The title compound (2*R*,4*S*)-hydroxyketamine {systematic name: (2*R*,4*S*)-2-(2-chlorophenyl)-4-hydroxy-2-methylamino)-cyclohexanone} was purified by flash chromatography on silica as previously described [5]. The solvent was evaporated and the residue was dissolved in methanol. Crystals were obtained by slow evaporation overnight from methanol.

<https://doi.org/10.1515/ncrs-2020-0157>

Received March 23, 2020; accepted April 23, 2020; available online June 18, 2020

***Corresponding author: U. Joost Luelf**, Institut für Biochemie, Heinrich-Heine-Universität Düsseldorf, Universitätsstrasse 1, 40225 Düsseldorf, Germany, e-mail: joost.luelf@hhu.de.
<https://orcid.org/0000-0002-6296-2278>

Guido J. Reiss: Institut für Anorganische Chemie und Strukturchemie, Lehrstuhl II: Material- und Strukturforchung, Heinrich-Heine-Universität Düsseldorf, Universitätsstrasse 1, 40225 Düsseldorf, Germany

Vlada B. Urlacher: Institut für Biochemie, Heinrich-Heine-Universität Düsseldorf, Universitätsstrasse 1, 40225 Düsseldorf, Germany

Table 2: Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²).

Atom	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl1	0.05378(9)	0.25991(10)	0.07451(5)	0.05572(19)
O1	0.1888(4)	−0.3498(3)	0.3458(2)	0.0702(7)
H1O	0.150(5)	−0.448(6)	0.350(3)	0.069(11)*
O2	0.0475(2)	0.2845(2)	0.34835(15)	0.0480(4)
N1	0.3678(3)	0.0021(3)	0.3870(2)	0.0507(5)
H1N	0.341(4)	−0.102(5)	0.403(3)	0.044(8)*
C1	0.0783(3)	0.1234(3)	0.32893(17)	0.0354(4)
C2	0.2580(3)	0.0585(3)	0.28374(18)	0.0342(4)
C3	0.2383(3)	−0.1139(3)	0.2045(2)	0.0411(5)
H12	0.198160	−0.073719	0.127520	0.049*
H13	0.355195	−0.169700	0.198221	0.049*
C4	0.1107(4)	−0.2606(3)	0.2448(2)	0.0503(6)
H44	0.092957	−0.352762	0.183122	0.060*
C5	−0.0678(4)	−0.1767(4)	0.2699(3)	0.0561(7)
H5A	−0.146834	−0.272738	0.296116	0.067*
H5B	−0.121082	−0.124178	0.199197	0.067*
C6	−0.0492(4)	−0.0262(4)	0.3625(2)	0.0480(5)
H6A	−0.165444	0.027927	0.374096	0.058*
H6B	−0.006709	−0.081142	0.435259	0.058*
C7	0.3512(3)	0.2111(3)	0.2179(2)	0.0374(4)
C8	0.5293(3)	0.2550(5)	0.2475(3)	0.0586(6)
H8	0.587639	0.193452	0.308931	0.070*
C9	0.6209(5)	0.3883(5)	0.1871(5)	0.0807(12)
H9	0.738859	0.415501	0.209126	0.097*
C10	0.5397(5)	0.4799(5)	0.0958(4)	0.0785(12)
H10	0.602036	0.568930	0.055869	0.094*
C11	0.3633(5)	0.4393(4)	0.0628(3)	0.0612(8)
H11	0.306591	0.500376	0.000529	0.073*
C12	0.2731(3)	0.3065(3)	0.1242(2)	0.0414(5)
C13	0.3658(5)	0.1213(5)	0.4897(3)	0.0738(10)
H13A	0.388078	0.246823	0.467645	0.111*
H13B	0.456806	0.081628	0.545421	0.111*
H13C	0.251167	0.113289	0.523331	0.111*

Experimental details

Coordinates of hydrogen atoms attached to nitrogen and oxygen were refined without constraints or restraints. The carbon-bound hydrogen atoms were placed using a riding model (AFIX 13/23/43/137) implemented in the SHELXL system using the standard parameters for the constrained $U_{\text{iso}}(\text{H})$ values [3]. The absolute structure determination succeeded as the derived Flack parameter is found to be near zero [0.01(5) from 1306 selected quotients] using Parsons's method [6]. The classical calculation of the Flack parameter showed a slightly worse result 0.02(8) using all reflections [3, 6].

Comment

The anesthetic and antidepressant ketamine (2-(2-chlorophenyl)-2-(methylamino)-cyclohexanone) is mainly metabolized by cytochrome P450s (CYPs) in the liver leading to, *inter alia*, several hydroxynorketamines and hydroxyketamines [7–11].

In order to synthesize some of these metabolites, which are considered to be the pharmacologically active compounds in treatment of depression [10–13], protein engineering of the cytochrome P450 CYP154E1 from the thermotolerant soil bacterium *Thermobifida fusca* YX was applied as described previously [5]. CYP154E1 was chosen as an appropriate candidate because it has been found to possess high substrate promiscuity and substrate-dependent high regioselectivity [5, 14–16]. Mutagenesis and screening among a number of CYP154E1 mutants revealed the double mutant I238G/M388A that enabled the regio- and stereoselective hydroxylation at position 4 of (R)-ketamine as well as of norketamine (not shown). To the best of our knowledge, the only hydroxylated at position 4 metabolites of ketamine reported so far are stereoisomers of 4-hydroxynorketamine.

Description and comparison. There is one hydroxyketamine molecule in the asymmetric unit of the title structure (*cf.* the Figure). As the compound crystallizes in the non-centrosymmetric space group $P2_1$ it is quite clear that the investigated crystal contains (2R,4S)-hydroxyketamine only. All bond lengths and angles derived from our diffraction experiment are in the expected ranges [17, 18]. The title molecule possesses a conformational flexibility by a rotation about the C2–C7 and C2–N1 single bonds, respectively. It is evident that these potential rotations will not take place independently from one another. Furthermore, the intramolecular $\text{NH} \cdots \text{O}$ hydrogen bond [$\text{N1} \cdots \text{O1} = 2.907(3)$; see the Figure] reduces the possibilities. Thus it is of interest to compare the conformation of the title structure with examples found in the literature. In the title structure C2–N1 and the chlorophenyl moiety are almost coplanar (rms deviation of fitted atoms N1, C2, C7–C12 = 0.045 Å). For the parent compound ketamine, structural reports for the racemic single crystal [17] as well as for both enantiomeric forms [18] are described. For all these ketamine structures the chlorophenyl moiety is almost coplanar with the aforementioned central C–N single bond. A CSD database survey [19] yielded structures of all four possible stereoisomers of the closely related 4-hydroxynorketamine [20]. In accordance with our results, the mean plane of the chlorophenyl group is almost coplanar with the C–N bond in the structures of the (2S,4S)- and (2R,4R)-stereoisomers. However, the stereoisomers (2R,4S)- and (2S,4R)-hydroxynorketamine were crystallized as hydrochlorides showing a rotation about C5–C7 as well as the other chair conformation of the cyclohexanone ring; the amino and hydroxy group are found in equatorial positions. Thus, an intramolecular hydrogen bond is not possible. In this context it should not be kept unmentioned that during our database survey we identified a ketamine hydrochloride, which shows the same rotation of the mean plane of the chlorophenyl moiety (rotation about C5–C7) and

the other chair conformation, too [21]. Protonation at N1 and formation of a salt with some more hydrogen bonds seems to seriously affect the general conformation. This crucial aspect should be investigated in the future.

Supramolecular aspects. Adjacent molecules are connected *via* intermolecular OH...O hydrogen bonds [O1...O2' = 2.850(3) Å; ' = x, -1 + y, z; see the Figure] to form a chain structure along the crystallographic *b* axis. Similar to other simple molecules containing almost rigid moieties [22], the packing scheme is a compromise between the conformational possibilities and hydrogen bonding interactions.

Acknowledgements: We gratefully acknowledge support by the Ministry of Innovation, Science and Research of North-Rhine Westphalia and the German Research Foundation (DFG) for financial support (D8/APEX2 diffractometer; INST 208/589-1). Generous financial support by the Federal Ministry of Education and Research to Heinrich Heine University Düsseldorf (grant number 031A223A) is also gratefully acknowledged. We also thank Ansgar Bokel (Heinrich Heine University Düsseldorf, Germany) for experimental support.

References

1. Bruker. APEX2, SAINT, SADABS. Bruker AXS Inc., Madison, WI, USA (2012).
2. Sheldrick, G. M.: SHELXT – Integrated space-group and crystal-structure determination. *Acta Crystallogr.* **A71** (2015) 3–8.
3. Sheldrick, G. M.: Crystal structure refinement with SHELXL. *Acta Crystallogr.* **C71** (2015) 3–8.
4. Brandenburg, K.: DIAMOND. Visual Crystal Structure Information System. Ver. 4.0. Crystal Impact, Bonn, Germany (2015).
5. Bokel, A.; Rühlmann, A.; Hutter, M. C.; Urlacher, V. B.: Enzyme-mediated two-step regio- and stereoselective synthesis of potential rapid-acting antidepressant (2S,6S)-hydroxynorketamine. *ACS Catal.* **10** (2020) 4151–4159.
6. Parsons, S.; Flack, H. D.; Wagner, T.: Use of intensity quotients and differences in absolute structure refinement. *Acta Crystallogr.* **B69** (2013) 249–259.
7. Adams Jr., J. D.; Baillie, T. A.; Trevor, A. J.; Castagnoli Jr., N.: Studies on the biotransformation of ketamine. 1-identification of metabolites produced *in vitro* from rat liver microsomal preparations. *Biomed. Mass Spectrom.* **8** (1981) 527–538.
8. Moaddel, R.; Venkata, S. L. V.; Tanga, M. J.; Bupp, J. E.; Green, C. E.; Iyer, L.; Furimsky, A.; Goldberg, M. E.; Torjman, M. C.; Wainer, I. W.: A parallel chiral-achiral liquid chromatographic method for the determination of the stereoisomers of ketamine and ketamine metabolites in the plasma and urine of patients with complex regional pain syndrome. *Talanta* **82** (2010) 1892–1904.
9. Desta, Z.; Moaddel, R.; Ogburn, E. T.; Xu, C.; Ramamoorthy, A.; Venkata, S. L. V.; Sanghvi, M.; Goldberg, M. E.; Torjman, M. C.; Wainer, I. W.: Stereoselective and regiospecific hydroxylation of ketamine and norketamine. *Xenobiotica* **42** (2012) 1076–87.
10. Zarate Jr, C. A.; Brutsche, N.; Laje, G.; Luckenbaugh, D. A.; Venkata, S. L. V.; Ramamoorthy, A.; Moaddel, R.; Wainer, I. W.: Relationship of ketamine's plasma metabolites with response, diagnosis, and side effects in major depression. *Biol. Psychiatry* **72** (2012) 331–338.
11. Zanos, P.; Moaddel, R.; Morris, P. J.; Riggs, L. M.; Highland, J. N.; Georgiou, P.; Pereira, E. F. R.; Albuquerque, E. X.; Thomas, C. J.; Zarate Jr, C. A.; Gould, T. D.: Ketamine and ketamine metabolite pharmacology: insights into therapeutic mechanisms. *Pharmacol. Rev.* **70** (2018) 621–660.
12. Hashimoto, K.: Rapid-acting antidepressant ketamine, its metabolites and other candidates: a historical overview and future perspective. *Psychiat. Clin. Neuros.* **73** (2019) 613–627.
13. Zanos, P.; Moaddel, R.; Morris, P. J.; Georgiou, P.; Fischell, J.; Elmer, G. I.; Alkondon, M.; Yuan, P.; Pribut, H. J.; Singh, N. S.; Dossou, K. S. S.; Fang, Y.; Huang, X.-P.; Mayo, C. L.; Wainer, I. W.; Albuquerque, E. X.; Thompson, S. M.; Thomas, C. J.; Zarate Jr., C. A.; Gould, T. D.: NMDAR inhibition-independent antidepressant actions of ketamine metabolites. *Nature* **533** (2016) 481–486.
14. von Bühler, C.; Le-Huu, P.; Urlacher, V. B.: Cluster screening: an effective approach for probing the substrate space of uncharacterized cytochrome P450s. *ChemBioChem* **14** (2013) 2189–2198.
15. Bogazkaya, A. M.; von Bühler, C.; Kriening, S.; Busch, A.; Seifert, A.; Pleiss, J.; Laschat, S.; Urlacher, V. B.: Selective allylic hydroxylation of acyclic terpenoids by CYP154E1 from *Thermobifida fusca* YX. *Beilstein J. Org. Chem.* **10** (2014) 1347–1353.
16. Rühlmann, A.; Antovic, D.; Müller, T. J. J.; Urlacher, V. B.: Regioselective hydroxylation of stilbenes by engineered cytochrome P450 from *Thermobifida fusca* YX. *Adv. Synth. Catal.* **359** (2017) 984–994.
17. Kassir, V.-E. H.; Géraudy, R.; Toupay, T.; Collin, D.; Salvadeo, E.; Toussaint, F.; Van Hecke, K.; Monbaliu, J.-C. M.: Expedient preparation of active pharmaceutical ingredient ketamine under sustainable continuous flow conditions. *Green Chem.* **21** (2019) 2952–2966.
18. Chankvetadze, B.; Burjanadze, N.; Breitzkreutz, J.; Bergander, K.; Bergenthal, D.; Kataeva, O.; Fröhlich, R.; Luftmann, H.; Blaschke, G.: Mechanistic study on the opposite migration order of the enantiomers of ketamine with α - and β -cyclodextrin in capillary electrophoresis. *J. Sep. Sci.* **25** (2002) 1155–1166.
19. Groom, C. R.; Allen, F. H.: The Cambridge Structural Database in retrospect and prospect. *Angew. Chem. Int. Ed.* **53** (2014) 662–671.
20. Morris, P. J.; Moaddel, R.; Zanos, P.; Moore, C. E.; Gould, T.; Zarate Jr, C. A.; Thomas, C. J.: Synthesis and *N*-methyl-*D*-aspartate (NMDA) receptor activity of ketamine metabolites. *Org. Lett.* **19** (2017) 4572–4575.
21. Haakey, P.; Ouellette, W.; Zubieta, J.; Korter, T.: (S)-(+)-Ketamine hydrochloride. *Acta Crystallogr.* **E64** (2008) o1487.
22. Reiss, G. J.: The crystal structure of 3-((1*R*,2*S*)-1-methylpyrrolidin-1-ium-2-yl)pyridin-1-ium tetrachloridomanganate(II), C₁₀H₁₆Cl₄MnN₂. *Z. Kristallogr. NCS* **235** (2020) 415–417.