Walid Guerrab, Mohcine Missioui, Younes Zaoui, Joel T. Mague and Youssef Ramli*

Synthesis and crystal structure of 2-azido-*N*-phenylacetamide, C₈H₈N₄O

Table 1: Data collection and handling.

Crystal:	Colourless block
Size:	0.40 × 0.32 × 0.30 mm
Wavelength:	Cu K α radiation (1.54178 Å)
μ :	0.79 mm ⁻¹
Diffractometer, scan mode:	D8 VENTURE PHOTON 100, ω
$\theta_{\max}$, completeness:	72.5°, 99%
$N(hkl)_{\text{measured}}$	8365, 3383, 0.032
$N(hkl)_{\text{unique}}$, R_{int} :	
Criterion for I_{obs} , $N(hkl)_{\text{gt}}$:	$I_{\text{obs}} > 2 \sigma(I_{\text{obs}})$, 3204
$N(\text{param})_{\text{refined}}$:	236
Programs:	Bruker [1], SHELX [2–4, 6], Diamond [5]

<https://doi.org/10.1515/ncrs-2020-0409>

Received July 27, 2022; accepted September 14, 2020;
published online October 5, 2020

Abstract

C₈H₈N₄O, orthorhombic, $P2_12_12_1$ (no. 19), $a = 5.8085(13)$ Å, $b = 16.533(4)$ Å, $c = 18.171(4)$ Å, $V = 1745.0(7)$ Å³, $Z = 8$, $R_{\text{gt}}(F) = 0.0399$, $wR_{\text{ref}}(F^2) = 0.1078$, $T = 296(2)$ K.

CCDC no.: 2031786

The molecular structure is shown in the Figure. Table 1 contains crystallographic data and Table 2 contains the list of the atoms including atomic coordinates and displacement parameters.

***Corresponding author: Youssef Ramli**, Laboratory of Medicinal Chemistry, Drug Sciences Research Center, Faculty of Medicine and Pharmacy, Mohammed V University, Rabat, Morocco, e-mail: y.ramli@um5s.net.ma. <https://orcid.org/0000-0002-6885-5692>

Walid Guerrab, Mohcine Missioui and Younes Zaoui, Laboratory of Medicinal Chemistry, Drug Sciences Research Center, Faculty of Medicine and Pharmacy, Mohammed V University, Rabat, Morocco, e-mail: guerrab.walid@gmail.com (W. Guerrab), mohcinemissioui@gmail.com (M. Missioui), younes.zaoui@gmail.com (Y. Zaoui)

Joel T. Mague, Department of Chemistry, Tulane University, New Orleans, LA 70118, USA, e-mail: joelt@tulane.edu

Source of material

2-Chloro-*N*-(4-nitrophenyl)acetamide (0.006 mol) and sodium azide (0.0085 mol) were dissolved in a mixture of ethanol/water (70:30, v:v) then refluxed for 24 h at 80 °C. After cooling the title compound precipitated and was filtered off, dried and recrystallized from ethanol, yield: 77%. A portion of the product was dissolved in hot ethanol, the solution was filtered and the filtrate was left undisturbed for 5 days to form clear, colorless block crystals.

Experimental details

Hydrogen atoms were included as riding contributions in idealized positions. The Flack-Parsons parameter was calculated as -0.04(10) based on 1251 quotients [6].

Comment

Azides have found valuable applications in medicinal chemistry [7], molecular biology [8], and an increasing interest in the field of organic synthesis as intermediates for the preparation of heterocycles such as tetrazoles, triazolines, triazoles, etc. [9–11]. As part of our ongoing research in this area, we report the synthesis and molecular structure of the title azide.

Table 2: Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²).

Atom	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> _{iso} [*] / <i>U</i> _{eq}
O1	0.2355 (3)	0.29894 (8)	0.23380 (8)	0.0657 (4)
N1	0.3120 (3)	0.19233 (9)	0.30918 (9)	0.0535 (4)
H1	0.272547	0.143385	0.319042	0.064*
N2	−0.1241 (5)	0.22663 (13)	0.17114 (14)	0.0871 (7)
N3	−0.2933 (4)	0.19339 (12)	0.14523 (12)	0.0707 (5)
N4	−0.4536 (5)	0.17043 (18)	0.11761 (17)	0.0993 (9)
C1	0.4922 (3)	0.22590 (11)	0.35168 (10)	0.0505 (4)
C2	0.5078 (4)	0.30827 (12)	0.36577 (12)	0.0625 (5)
H2	0.397649	0.343636	0.347233	0.075*
C3	0.6891 (5)	0.33725 (16)	0.40772 (14)	0.0754 (7)
H3	0.700620	0.392420	0.417011	0.090*
C4	0.8527 (5)	0.28540 (18)	0.43585 (12)	0.0741 (7)
H4	0.975150	0.305343	0.463358	0.089*
C5	0.8324 (5)	0.20377 (17)	0.42272 (13)	0.0729 (6)
H5	0.941466	0.168440	0.441990	0.087*
C6	0.6533 (5)	0.17367 (14)	0.38152 (11)	0.0638 (5)
H6	0.640454	0.118259	0.373706	0.077*
C7	0.1968 (3)	0.22964 (11)	0.25490 (10)	0.0510 (4)
C8	0.0165 (4)	0.17728 (12)	0.21914 (12)	0.0617 (5)
H8A	0.089937	0.134667	0.190952	0.074*
H8B	−0.079095	0.152361	0.256584	0.074*
O2	0.6474 (3)	0.46627 (9)	0.63830 (10)	0.0737 (5)
N5	0.4595 (3)	0.58698 (10)	0.62794 (10)	0.0587 (4)
H5A	0.457556	0.635003	0.646123	0.070*
N6	0.8264 (4)	0.50883 (14)	0.77325 (14)	0.0826 (7)
N7	1.0343 (4)	0.50238 (11)	0.76906 (11)	0.0679 (5)
N8	1.2245 (6)	0.4922 (2)	0.7732 (2)	0.1183 (12)
C9	0.3063 (4)	0.57312 (12)	0.56913 (11)	0.0561 (5)
C10	0.3275 (6)	0.51010 (14)	0.52004 (13)	0.0753 (7)
H10	0.450885	0.474455	0.523344	0.090*
C11	0.1625 (7)	0.50050 (18)	0.46569 (16)	0.0985 (11)
H11	0.174874	0.457034	0.433444	0.118*
C12	−0.0139 (7)	0.5518 (2)	0.45813 (16)	0.0967 (10)
H12	−0.122525	0.543667	0.421252	0.116*
C13	−0.0341 (6)	0.6168 (2)	0.50528 (16)	0.0950 (9)
H13	−0.154569	0.653305	0.499807	0.114*
C14	0.1267 (5)	0.62718 (18)	0.56094 (13)	0.0778 (7)
H14	0.113588	0.670847	0.592930	0.093*
C15	0.6087 (4)	0.53509 (12)	0.65948 (12)	0.0565 (5)
C16	0.7281 (5)	0.57151 (14)	0.72602 (15)	0.0742 (7)
H16A	0.849637	0.607510	0.709639	0.089*
H16B	0.618243	0.603172	0.754047	0.089*

The asymmetric unit consists of two independent molecules differing primarily in the orientation of the azide moiety. Thus in the molecule containing O1, the dihedral angle between the mean planes defined by N1/C7/C8/O1 and C8/N2/N3/N4 is 10.9(2)° (see the Figure), while the corresponding angle in the molecule containing O2 is 80.9(2)°. Also, in the molecule containing O1, the dihedral angle between the mean planes defined by C1⋯C6 and N1/C7/C8/O1 is 28.4(1)°, while the corresponding angle in the other molecule is 16.3(2)°. In the crystal, the two

independent molecules are linked in alternating fashion by N1–H1⋯O2 and N5–H5A⋯O1 hydrogen bonds into helical chains extending along the *b*-axis direction (Table 2). The chains are joined into layers parallel to the *bc* plane by π interactions between the N3/N4 moiety of the azide substituent and the C1⋯C6 ring at $-x, y - 1/2, -z + 1/2$ (N4⋯centroid = 3.462(3) Å, N3⋯centroid = 3.695(3) Å, N3/N4⋯centroid = 93.0(2)°).

Author contribution: All the authors have accepted responsibility for the entire content of this submitted manuscript and approved submission.

Funding information: The support of NSF-MRI Grant #1228232 for the purchase of the diffractometer and Tulane University for support of the Tulane Crystallography Laboratory are gratefully acknowledged.

Conflict of interest statement: The authors declare no conflicts of interest regarding this article.

References

1. Bruker. *APEX3 – Diffractometer Control and Data Reduction Software Package*; Bruker AXS Inc.: Madison, WI, USA, 2016.
2. Krause L., Herbst-Irmer R., Sheldrick G. M., Stalke D. Comparison of silver and molybdenum microfocus X-ray sources for single-crystal structure determination. *J. Appl. Crystallogr.* 2015, 48, 3–10.
3. Sheldrick G. M. SHELXT – integrated space-group and crystal-structure determination. *Acta Crystallogr.* 2015, A71, 3–8.
4. Sheldrick G. M. Crystal structure refinement with *SHELXL*. *Acta Crystallogr.* 2015, C71, 3–8.
5. Brandenburg K. *DIAMOND. Visual Crystal Structure Information System*. Version 4.0; Crystal Impact: Bonn, Germany, 2015.
6. Parsons S., Flack H. D., Wagner T. Use of intensity quotients and differences in absolute structure refinement. *Acta Crystallogr.* 2013, B69, 249–259.
7. Contin M., Sepulveda C., Fascio M., Stortz C. A., Damonte E. B., D'Accorso N. B. Modified ribavirin analogues as antiviral agents against Junin virus. *Bioorg. Med. Chem. Lett.* 2019, 29, 556–559.
8. Ahmed S., Abdallah N. A. Dansyl azide as a selective fluorescence tagging probe for click chemistry reactions and its application to monitor rasagiline in pharmacokinetic studies. *J. Pharm. Biomed. Anal.* 2019, 165, 357–365.
9. Chauhan U. B., Tomich A. W., Lavallo V. The first example of a click reaction with a carboranyl azide and an olefin. *Tetrahedron* 2019, 75, 1323–1325.
10. Bakulev V., Shafran Y., Dehaen W. Progress in intermolecular and intramolecular reactions of thioamides with diazo compounds and azides. *Tetrahedron Lett.* 2019, 60, 513–523.
11. Abad N., Hajji M., Ramli Y., Belkhiria M., Elmgirhi S. M. H., Habib M. A., Guerfel T., Mague J. T., Essassi E. M. A newly synthesized nitrogen-rich derivative of bicyclic quinoxaline-structural and conceptual DFT reactivity study. *J. Phys. Org. Chem.* 2020, 33, e4055.