Mamolatelolo J. Moremi*, Orbett T. Alexander, Banele Vatsha, Katlego Makgopa and Amanda-Lee E. Manicum

The crystal structure of *fac*-tricarbonyl(4,4-dimethyl-2,2-dipyridyl- κ^2N,N')-(pyrazole- κN)rhenium(I) nitrate, $C_{18}H_{16}O_3N_4Re$

Table 1: Data collection and handling.

Crystal:	Yellow block
Size:	0.20 × 0.17 × 0.10 mm
Wavelength:	Mo K α radiation (0.71073 Å)
μ :	6.35 mm ⁻¹
Diffractometer, scan mode:	Bruker APEX-II, φ and ω
$\theta_{\max}$, completeness:	26.1°, >99%
$N(hkl)_{\text{measured}}$, $N(hkl)_{\text{unique}}$, R_{int} :	25,596, 3854, 0.064
Criterion for I_{obs} , $N(hkl)_{\text{gt}}$:	$I_{\text{obs}} > 2 \sigma(I_{\text{obs}})$, 3315
$N(\text{param})_{\text{refined}}$:	273
Programs:	Bruker [1], SIR97 [2], Olex2 [3], SHELX [4], Diamond [5]

<https://doi.org/10.1515/ncrs-2020-0458>

Received August 27, 2020; accepted October 12, 2020;

published online November 6, 2020

Abstract

$C_{18}H_{16}O_3N_4Re$, monoclinic, $P2_1/c$ (no. 14), $a = 9.8409(6)$ Å, $b = 14.0933(9)$ Å, $c = 13.9153(9)$ Å, $\beta = 90.558(2)^\circ$, $V = 1929.8(2)$ Å³, $Z = 4$, $R_{\text{gt}}(F) = 0.0266$, $wR_{\text{ref}}(F^2) = 0.0584$, $T = 100(2)$ K.

CCDC no.: 2036274

The molecular structure is shown in the Figure. Table 1 contains crystallographic data, and Table 2 contains the list of the atoms including atomic coordinates and displacement parameters.

Source of material

The starting complex was synthesized according to a published procedure [14]: *fac*-[Re(Ddp)(CO)₃(H₂O)]⁺ (30.00 mg; 0.063 mmol) was dissolved in methanol (5 ml) and pyrazole (4.316 mg, 0.063 mmol) was added; followed by refluxing for 24 h. The bright yellow solid was obtained by evaporation of the solvent, and the product was recrystallized using cold methanol.

IR (FTIR cm⁻¹): $\nu_{\text{CO}} = 2021.6, 1888.1, 1618$.

Experimental details

All the hydrogen atoms were positioned geometrically and refined discernibly using a riding model, with fixed C-H_{methyl} = 0.96 Å; C-H_{Aromatic} = 0.97 Å. The H atoms isotropic displacement parameters were fixed; $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ for aromatic and $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$ for methyl, allowing them to ride on the parent atom. The graphics were obtained using the DIAMOND program with 50% probability ellipsoids. All the H-atoms on the title structure were omitted for clarity.

*Corresponding author: Mamolatelolo J. Moremi, Department of Chemistry, Tshwane University of Technology, Pretoria, 0001, South Africa, E-mail: joshmolatelolo2@gmail.com. <https://orcid.org/0000-0003-4837-213X>

Orbett T. Alexander, Department of Chemistry, University of the Free State, Bloemfontein 9301, South Africa. <https://orcid.org/0000-0003-4926-8342>

Banele Vatsha, Department of Chemistry, University of Johannesburg, Auckland Park 2006, South Africa

Katlego Makgopa and Amanda-Lee E. Manicum, Department of Chemistry, Tshwane University of Technology, Pretoria 0001, South Africa

Table 2: Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²).

Atom	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> _{iso} */ <i>U</i> _{eq}
Re1	0.36523 (2)	0.36667 (2)	0.15167 (2)	0.01396 (7)
O1	0.4160 (3)	0.3151 (2)	−0.0595 (2)	0.0264 (7)
O2	0.5057 (3)	0.1840 (2)	0.2198 (2)	0.0262 (7)
O3	0.6380 (3)	0.4703 (2)	0.1279 (2)	0.0243 (7)
O4	0.6958 (3)	0.3629 (2)	0.3347 (2)	0.0241 (7)
O5	0.6997 (3)	0.5165 (2)	0.3356 (2)	0.0257 (7)
O6	0.8885 (3)	0.4380 (2)	0.3395 (3)	0.0359 (9)
N1	0.1627 (3)	0.3057 (2)	0.1449 (2)	0.0137 (7)
N2	0.2300 (3)	0.4857 (2)	0.1288 (2)	0.0145 (7)
N3	0.3273 (3)	0.4026 (3)	0.3010 (2)	0.0160 (7)
N4	0.4256 (3)	0.4002 (3)	0.3699 (2)	0.0177 (8)
H4	0.511688	0.387514	0.359227	0.021*
N5	0.7630 (3)	0.4398 (3)	0.3361 (2)	0.0200 (8)
C1	0.3933 (4)	0.3347 (3)	0.0188 (3)	0.0193 (10)
C2	0.4571 (4)	0.2529 (3)	0.1925 (3)	0.0189 (9)
C3	0.5366 (4)	0.4316 (3)	0.1429 (3)	0.0186 (10)
C4	0.1367 (4)	0.2118 (3)	0.1458 (3)	0.0160 (9)
H4A	0.209268	0.169252	0.159136	0.019*
C5	0.0098 (4)	0.1753 (3)	0.1285 (3)	0.0179 (9)
H5	−0.003982	0.108613	0.129190	0.021*
C6	−0.0989 (4)	0.2360 (3)	0.1097 (3)	0.0163 (9)
C7	−0.2374 (4)	0.1983 (3)	0.0841 (3)	0.0257 (11)
H7A	−0.303757	0.250042	0.086171	0.039*
H7B	−0.262891	0.148950	0.130138	0.039*
H7C	−0.235731	0.171266	0.019210	0.039*
C8	−0.0722 (4)	0.3320 (3)	0.1119 (3)	0.0152 (9)
H8	−0.144425	0.375800	0.101995	0.018*
C9	0.0581 (4)	0.3657 (3)	0.1285 (3)	0.0148 (9)
C10	0.0936 (4)	0.4671 (3)	0.1268 (3)	0.0118 (8)
C11	−0.0010 (4)	0.5396 (3)	0.1253 (3)	0.0152 (9)
H11	−0.095228	0.525011	0.123859	0.018*
C12	0.0410 (4)	0.6336 (3)	0.1258 (3)	0.0164 (9)
C13	−0.0610 (4)	0.7129 (3)	0.1215 (3)	0.0208 (10)
H13A	−0.073396	0.733384	0.054624	0.031*
H13B	−0.028121	0.766444	0.160198	0.031*
H13C	−0.148006	0.690739	0.146723	0.031*
C14	0.1789 (5)	0.6515 (3)	0.1301 (3)	0.0197 (10)
H14	0.210879	0.715056	0.132844	0.024*
C15	0.2698 (4)	0.5775 (3)	0.1306 (3)	0.0169 (9)
H15	0.364265	0.591403	0.132226	0.020*
C16	0.2138 (4)	0.4253 (3)	0.3478 (3)	0.0165 (9)
H16	0.127329	0.432742	0.317796	0.020*
C17	0.2380 (4)	0.4367 (3)	0.4455 (3)	0.0178 (9)
H17	0.174469	0.452672	0.493854	0.021*
C18	0.3742 (4)	0.4196 (3)	0.4565 (3)	0.0191 (10)
H18	0.423711	0.421314	0.515383	0.023*

Comment

The applications of the *fac*-[M(H₂O)₃(CO)₃]⁺ synthon, (M = Tc-99m, Re) and the introduction of nanotechnology merged with coordination chemistry, generally referred to

as nano-medicine is a highly investigated field due to their prospective and existing applications in radiochemistry [6–9]. Nanoparticles are the major system used in nano-medicine as theranostic agents because of their high molecular specificity which permit them to absorb high quantities of drugs [10–12].

The molecular structure of the title complex comprises three facial tricarbonyl ligands, a 4,4-dimethyl-2,2-dipyridyl bidentate ligand in the equatorial plane which is *trans* to two of the carbonyl ligands and a *N*-coordinated pyrazole monodentate ligand in the axial position. The complex was obtained from the [2 + 1] mixed ligand system on the *fac*-[M(H₂O)₃(CO)₃]⁺ moiety.

The model involves the substitution of the two labile water molecules in the equatorial plane, followed by the coordination of the mono-dentate aqua ligand in the axial position; and the complex is neutralized by a nitrate counter-ion. The polyhedron geometry of the title complex reveals an octahedral distortion around rhenium(I). The distortion is evident in the angles of 167.56(14)° and 171.99(14)° for C2–Re1–N2 and C3–Re1–N1 respectively. The bite angle of the chelate ligand of 74.76(12)° from the title structure correlates well with those observed in literature reported similar structures [13–14]. The bond distances between rhenium and the coordinated N atoms in the first coordination sphere ranges between 2.162(3) Å and 2.174(3) Å whereas the bond distances between the rhenium and the carbonyl carbon atoms ranges between 1.923(5) Å and 1.926(5) Å.

Author contribution: All the authors have accepted responsibility for the entire content of this submitted manuscript and approved submission.

Research funding: The authors would like to express the gratitude towards NRF (Thuthuka grant specific number 113629) for financial assistance. We would also like to acknowledge Tshwane University of Technology for Institutional support.

Conflict of interest: The authors declare no conflicts of interest regarding this article.

References

1. Bruker. *SAINT-Plus* (Version 7.12) and *SADABS* (Version 2004/1); Bruker AXS Inc.: Madison, Wisconsin, USA, 2004.
2. Altomare A., Burla M. C., Camalli M., Cascarano G. L., Giacovazzo C., Guagliardi A., Moliterni A. G. G., Polidori G., Spagna R. A new tool for crystal structure determination and refinement. *J. Appl. Cryst.* 1999, 32, 115–119.
3. Dolomanov O. V., Bourhis L. J., Gildea R. J., Howard J. A. K., Puschmann H. OLEX2: a complete structure solution, refinement and analysis program. *J. Appl. Cryst.* 2009, 42, 339–341.

4. Sheldrick G. M. A short history of *SHELX*. *Acta Crystallogr.* 2008, *A64*, 112–122.
5. Brandenbug K., Putz H. *DIAMOND*. Visual Crystal Structure Information System. Ver. 3; Oc Crystal Impact GbR: Bonn, Germany, 2005.
6. Alberto R., Schibli R., Egi A., Schubiger A. P., Abram U., Kaden T. A. A novel organometallic aqua complex of technetium for the labelling of biomolecules: synthesis of $[^{99}Tc(OH_2)(CO)_3]^+$ from $[^{99m}Tc(OH)_4]^-$ in aqueous solution and its reaction with a bifunctional ligand. *J. Am. Chem. Soc.* 1998, *120*, 7987–7988.
7. Jurisson S., Lydon J. Potential technetium small molecule radiopharmaceuticals. *Chem. Rev.* 1999, *99*, 2205–2218.
8. Manicum A., Smith M. S., Alexander O. T., Twigge L., Roodt A., Visser H. G. First kinetic data of the CO substitution in *fac*- $[Re(L, L'-Bid)(CO)_3(X)]$ complexes by tertiary phosphines PTA and PPh_3 : synthesis and crystal structures of water soluble rhenium(I) tri- and dicarbonyl complexes with 1,3,5-triaza-7-phosphaadamantane (PTA). *Inorg. Chem. Comm.* 2019, *101*, 93–98.
9. Bao G., Mitragotri S., Tong S. Multifunctional nanoparticles for drug delivery and molecular imaging. *Annu. Rev. Biomed. Eng.* 2013, *15*, 253–282.
10. Celardo I., Pedersen J. Z., Traversa E., Ghibelli L. Pharmacological potential of cerium oxide nanoparticles. *Nanoscale* 2011, *3*, 1411–1420.
11. De Jong W. H., Borm P. J. Drug delivery and nanoparticles: applications and hazards. *Int. J. Nanomed.* 2008, *3*, 133–149.
12. Borm P. J., Robbins D., Haubold S. The potential risks of nanomaterials: a review carried out for ECETOC. *Part. Fibre Toxicol.* 2006, *3*, 11–13.
13. Yazdani A., Jansen N., Banevicius L., Czorny S., Valliant J. F. Imidazole-based $[2+1]$ $Re(I)$ / ^{99m}Tc complexes as isostructural nuclear and optical probes. *Inorg. Chem.* 2015, *54*, 1728–1736.
14. Ramoba V. L., Alexander T. O., Visser H. G., Manicum A. The crystal structure of *fac*-tricarbonyl (1,10-phenanthroline- κ^2N, N')-(pyrazole- κN) rhenium(I)nitrate, $C_{18}H_{12}O_3N_4Re$. *Z. Kristallogr. NCS* 2020, *235*, 1203–1205.