Siyi Ding, Min Zhu and Zong-Cheng Miao

The crystal structure of 2-(4-bromophenyl)-2,3-dihydro-1*H*-naphtho[1,8-*de*][1,3,2]diazaborinine, C₁₆H₁₂BBrN₂

Table 1: Data collection and handling.

Crystal:	Colorless needle
Size:	0.10 × 0.08 × 0.06 mm
Wavelength:	Ga K α radiation (1.34139 Å)
μ :	2.72 mm ⁻¹
Diffractometer, scan mode:	Bruker APEX-II, φ and ω
$\theta_{\max}$, completeness:	53.9°, 98%
$N(hkl)_{\text{measured}}$, $N(hkl)_{\text{unique}}$, R_{int} :	3991, 2427, 0.044
Criterion for I_{obs} , $N(hkl)_{\text{gt}}$:	$I_{\text{obs}} > 2 \sigma(I_{\text{obs}})$, 2188
$N(\text{param})_{\text{refined}}$:	181
Programs:	Bruker [1], Olex2 [2], SHELX [3, 4]

Source of material

In air, 2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-2,3-dihydro-1*H*-naphtho[1,8-*de*][1,3,2]diazaborinine (Bpin-B(dan)) (0.1 mmol, 1.0 eq.), 4-bromoaniline (0.2 mmol, 2.0 eq.), TBAI (tetrabutylammonium iodide 0.01 eq.), NaOAc (0.15 eq.), and BPO (0.01 eq.) were sequentially weighed and added to a screw-capped Schlenk tube containing a magnetic stir bar. The vessel was evacuated and refilled with nitrogen. Tert-butyl nitrite (*t*BuONO) (0.2 eq.) and MeCN (0.6 mL) were added in turn under N₂ atmosphere using syringes through a septum which was temporarily used to replace the screw cap. The reaction mixture was then vigorously stirred at 80 °C for the indicated time. The resulting mixture was filtered through a pad of Celite®, and the filter cake was washed with ethyl acetate (3 mL × 2). The combined filtrate was evaporated under vacuum to dryness and the residue was purified by column chromatography to yield the desired product as colorless solid.

Experimental details

All the H atoms on the benzene rings were placed geometrically and refined without any constraints or restraints.

Comment

Organoboron compounds, which have been broadly used in synthetic chemistry, drug discovery and materials

<https://doi.org/10.1515/ncrs-2020-0556>

Received October 30, 2020; accepted November 26, 2020;

published online December 10, 2020

Abstract

C₁₆H₁₂BBrN₂, monoclinic, $P2_1$ (no. 4), $a = 4.7061(3)$ Å, $b = 10.4639(6)$ Å, $c = 13.5675(8)$ Å, $\beta = 97.512(2)^\circ$, $V = 662.39(7)$ Å³, $Z = 2$, $R_g(F) = 0.0384$, $wR_{\text{ref}}(F^2) = 0.0995$, $T = 189.98$ K.

CCDC no.: 2040532

The molecular structure is shown in the figure. Table 1 contains crystallographic data and Table 2 contains the list of the atoms including atomic coordinates and displacement parameters.

***Corresponding author: Zong-Cheng Miao**, Xi'an Key Laboratory of Advanced Photo-electronics Materials and Energy Conversion Device, School of Science, Xijing University, Xi'an, Shaanxi 710123, China, E-mail: miaocongcheng@xijing.edu.cn

Siyi Ding, Min Zhu and Zong-Cheng Miao, Xi'an Key Laboratory of Advanced Photo-electronics Materials and Energy Conversion Device, School of Science, Xijing University, Xi'an, Shaanxi 710123, China, E-mail: dingsiyi2009@163.com (S. Ding), miaocongcheng@xijing.edu.cn (Z.-C. Miao). <https://orcid.org/0000-0001-5446-0254> (S. Ding). <https://orcid.org/0000-0003-0454-3187> (M. Zhu)

Table 2: Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²).

Atom	x	y	z	U _{iso} [*] /U _{eq}
Br01	1.04997 (8)	0.87888 (9)	0.07431 (3)	0.0290 (2)
N1	0.1275 (7)	0.3841 (8)	0.1996 (3)	0.0263 (8)
H1	0.172177	0.371879	0.139324	0.032*
N2	0.1809 (8)	0.5011 (4)	0.3546 (3)	0.0233 (8)
H2	0.259294	0.563560	0.392208	0.028*
C1	−0.0108 (10)	0.4214 (5)	0.3931 (4)	0.0221 (10)
C2	−0.0826 (11)	0.4377 (6)	0.4887 (4)	0.0279 (11)
H2A	0.001973	0.504833	0.529613	0.033*
C3	−0.2794 (10)	0.3552 (5)	0.5242 (3)	0.0287 (11)
H3	−0.328301	0.367807	0.589212	0.034*
C4	−0.4039 (9)	0.2566 (5)	0.4677 (3)	0.0258 (10)
H4	−0.537117	0.201951	0.493733	0.031*
C5	−0.3332 (9)	0.2358 (5)	0.3697 (4)	0.0230 (9)
C6	−0.4550 (10)	0.1351 (5)	0.3088 (4)	0.0271 (10)
H6	−0.585788	0.077775	0.333244	0.032*
C7	−0.3850 (12)	0.1202 (5)	0.2151 (4)	0.0302 (11)
H7	−0.469792	0.052713	0.174603	0.036*
C8	−0.1906 (10)	0.2023 (5)	0.1773 (4)	0.0264 (10)
H8	−0.143673	0.189401	0.112073	0.032*
C9	−0.0665 (9)	0.3020 (5)	0.2343 (3)	0.0210 (9)
C10	0.4664 (9)	0.5826 (5)	0.2126 (3)	0.0207 (9)
C11	0.5168 (11)	0.5747 (5)	0.1132 (4)	0.0287 (11)
H11	0.427910	0.508311	0.072431	0.034*
C12	0.6916 (11)	0.6605 (6)	0.0728 (4)	0.0287 (11)
H12	0.724365	0.652767	0.005451	0.034*
C13	0.8178 (9)	0.7573 (4)	0.1313 (3)	0.0219 (9)
C14	0.7790 (10)	0.7694 (5)	0.2296 (4)	0.0270 (10)
H14	0.869439	0.836221	0.269481	0.032*
C15	0.6050 (10)	0.6817 (5)	0.2689 (4)	0.0255 (10)
H15	0.578574	0.689028	0.336895	0.031*
C16	−0.1353 (9)	0.3201 (5)	0.3329 (3)	0.0199 (9)
B1	0.2587 (10)	0.4874 (6)	0.2567 (4)	0.0214 (10)

science, have attracted significant attention in recent years [5]. They can react with various functional groups to construct new carbon-carbon bonds or carbon-heteroatom bonds, which can rapidly construct the complex structures of target molecules [6, 7]. When the naphthalene-1,8-diaminato (dan) ligand is used as the masking group, which has two nitrogen atoms that may donate their lone pair electrons to the vacant *p*-orbital of the boron atoms, the formed naphthalene-1,8-diamino boronamide (Bdan) compounds are robust enough to avoid undesirable organic reactions, such as Suzuki-Miyaura coupling reaction. Moreover, such compounds can be easily transformed to their corresponding boronic acids by simple treatment under aqueous acidic conditions. These features enable

their wide applications as modular synthetical building blocks [8–10], especially in the iterative cross-coupling reactions and the application of di-boron compounds. So here we report the crystal structure of the title compound (*cf.* figure).

There is one molecule in the symmetric unit (see the figure). All bonds and angles in the crystal structure are within the normal range [11]. In conclusion, we have developed a facile process for the synthesis of 2-(4-bromophenyl)-2,3-dihydro-1*H*-naphtho[1,8-*de*][1,3,2]diazaborinine, C₁₆H₁₂BBrN₂.

Author contribution: All the authors have accepted responsibility for the entire content of this submitted manuscript and approved submission.

Research funding: The National Natural Science Foundation of China (No. 21901214) and the Natural Science Basic Research Plan in Shaanxi Province of China (No. 2020JQ-918).

Conflict of interest statement: The authors declare no conflicts of interest regarding this article.

References

1. Bruker. *APEX-2, SAINT+, Version 6.02 (Includes XPREF and SADABS)*; Bruker AXS Inc.: Madison, WI, USA, 2016.
2. Dolomanov O. V., Bourhis L. J., Gildea R. J., Howard J. A. K., Puschmann H. OLEX2: a complete structure solution, refinement and analysis program. *J. Appl. Crystallogr.* 2009, 42, 339–341.
3. Sheldrick G. M. A short history of SHELX. *Acta Crystallogr.* 2008, A64, 112–122.
4. Sheldrick G. M. Crystal structure refinement with SHELXL. *Acta Crystallogr.* 2015, C71, 3–8.
5. Hall D. J. *Boronic acids*; Wiley: Weinheim, Germany, 2011.
6. Lennox A. J., Lloyd-Jones G. C. Selection of boron reagents for Suzuki–Miyaura coupling. *Chem. Soc. Rev.* 2014, 43, 412–413.
7. Fyfe J. W. B., Watson A. J. B. Recent development in organoboron chemistry: old dogs, new tricks. *Chem* 2017, 3, 31–55.
8. Xu L., Zhang S., Li P. Boron-selective reactions as powerful tools for modular synthesis of synthesis of diverse complex molecules. *Chem. Soc. Rev.* 2015, 44, 8848–8858.
9. Takaya J., Iwasawa N. Catalytic, direct synthesis of bis(boronate) compounds. *ACS Catal.* 2012, 2, 1993–2006.
10. Xu L., Li P. Differentiated di- and polyboron compounds: synthesis and application in successive Suzuki–Miyaura coupling. *Synlett* 2014, 25, 1799–1802.
11. Akerman M. P., Robinson R. S., Slabber C. A. 2-(4-Chlorophenyl)naphtho[1,8-*de*][1,3,2]diazaborinine. *Acta Crystallogr.* 2011, E67, o1873.