Tian-Xu Qiao, Yu-Xiao Chen, Jing-Yi An and Feng-Mei Nie*

Crystal structure of {tris((1*H*-benzo[*d*]imidazol-2-yl)methyl)amine- $\kappa^4 N, N', N'', N'''$ }-succinato- $\kappa^2 O, O'$ nickel(II) – methanol (1/4), $C_{32}H_{41}N_7NiO_8$

Table 1: Data collection and handling.

Crystal:	Blue block
Size:	0.34 × 0.32 × 0.30 mm
Wavelength:	Mo K α radiation (0.71073 Å)
μ :	0.64 mm ⁻¹
Diffractometer, scan mode:	Xcalibur, ω
$\theta_{\max}$, completeness:	26.0°, >99%
$N(hkl)_{\text{measured}}$, $N(hkl)_{\text{unique}}$, R_{int} :	16175, 6632, 0.055
Criterion for I_{obs} , $N(hkl)_{\text{gt}}$:	$I_{\text{obs}} > 2 \sigma(I_{\text{obs}})$, 4640
$N(\text{param})_{\text{refined}}$:	448
Programs:	CrysAlis ^{PRO} [1], Olex2 [2], SHELX [3]

<https://doi.org/10.1515/ncrs-2020-0608>

Received November 21, 2020; accepted December 15, 2020;

published online January 12, 2021

Abstract

$C_{32}H_{41}N_7NiO_8$, monoclinic, $P2_1/c$ (no. 14), $a = 10.0076(5)$ Å, $b = 18.6561(10)$ Å, $c = 19.0893(18)$ Å, $\beta = 108.532(6)^\circ$, $V = 3378.4(4)$ Å³, $Z = 4$, $R_g(F) = 0.0607$, $wR_{\text{ref}}(F^2) = 0.1340$, $T = 111.9(1)$ K.

CCDC no.: 2050294

The molecular structure is shown in the figure (four solvent molecules which belong to the asymmetric unit are not shown). Table 1 contains crystallographic data and Table 2 contains the list of the atoms including atomic coordinates and displacement parameters.

Source of material

Tris(benzimidazol-2-yl-methyl)amine (ntb) was synthesized according to a literature procedure [4]. A methanol

solution (50 mL) of $Ni(ClO_4)_2 \cdot 6H_2O$ (73 mg, 0.2 mmol) and ntb (82 mg, 0.2 mmol) was stirred for 10 min. Then a methanol solution (40 mL) of disodium succinate hexahydrate (54 mg, 0.2 mmol) was added dropwise to the aforementioned solution. After being stirred at room temperature for 2 h, the solution was filtered for slow evaporation. Light-blue crystals formed.

Experimental details

The structure was solved with the Olex2 program [2]. The methyl groups were idealized and refined using rigid groups allowed to rotate about the N–C bond (with the SHELX program [3]). The U_{iso} values of the hydrogen atoms of methyl groups were set to 1.5 U_{eq} and the values of all other hydrogen atoms were set to 1.2 U_{eq} . The disordered C27 and C28 atoms of succinate including H atoms on them were located at two sites with occupancies to be 0.503(5) for C27, C28 and 0.497(5) for C27', C28' (Table 2).

Comment

The ligand ntb is often used in the construction of a variety of metal complexes [5]. Some mononuclear nickel(II) complexes have been reported which use monocarboxylate (isonicotinate, picolinate and 3, 5-dinitrobenzoate), azide

*Corresponding author: Feng-Mei Nie, Department of Chemistry, Capital Normal University, Beijing, 100048, P.R. China, E-mail: niefm@mail.cnu.edu.cn. <https://orcid.org/0000-0001-9150-0729>

Tian-Xu Qiao, Yu-Xiao Chen and Jing-Yi An, Department of Chemistry, Capital Normal University, Beijing, 100048, P.R. China

Table 2: Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²).

Atom	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> _{iso} */ <i>U</i> _{eq}
C1	−0.2079 (3)	0.35410 (19)	0.13583 (18)	0.0209 (7)
H1A	−0.273228	0.395496	0.124452	0.025*
H1B	−0.265262	0.309762	0.125465	0.025*
C2	−0.1232 (3)	0.35589 (17)	0.21544 (18)	0.0156 (7)
C3	0.0529 (4)	0.35991 (18)	0.31572 (19)	0.0225 (8)
C4	−0.0697 (4)	0.35832 (18)	0.33607 (19)	0.0223 (8)
C5	−0.0652 (5)	0.3596 (2)	0.4096 (2)	0.0378 (10)
H5A	−0.148847	0.358306	0.422916	0.045*
C6	0.0653 (5)	0.3629 (3)	0.4619 (2)	0.0605 (15)
H6	0.072830	0.363675	0.512809	0.073*
C7	0.1876 (5)	0.3651 (3)	0.4415 (2)	0.0640 (16)
H7A	0.276241	0.367728	0.479218	0.077*
C8	0.1849 (4)	0.3636 (2)	0.3690 (2)	0.0385 (11)
H8	0.269026	0.364986	0.356023	0.046*
C9	−0.1369 (4)	0.42405 (18)	0.04423 (19)	0.0233 (8)
H9A	−0.236349	0.439825	0.030601	0.028*
H9B	−0.112959	0.416519	−0.001719	0.028*
C10	−0.0420 (4)	0.47997 (18)	0.09122 (19)	0.0216 (8)
C11	0.1407 (3)	0.52716 (18)	0.17066 (19)	0.0205 (8)
C12	0.0514 (4)	0.58354 (18)	0.1359 (2)	0.0220 (8)
C13	0.0850 (4)	0.65469 (19)	0.1541 (2)	0.0287 (9)
H13	0.023488	0.692540	0.130617	0.034*
C14	0.2115 (4)	0.6679 (2)	0.2079 (2)	0.0367 (10)
H14	0.238549	0.716000	0.221546	0.044*
C15	0.3014 (4)	0.6119 (2)	0.2430 (2)	0.0338 (9)
H15	0.388310	0.623173	0.279604	0.041*
C16	0.2673 (4)	0.54087 (19)	0.2258 (2)	0.0261 (8)
H16	0.327635	0.503115	0.250581	0.031*
C17	−0.1357 (4)	0.29061 (19)	0.0414 (2)	0.0321 (9)
H17A	−0.094032	0.297107	0.001200	0.039*
H17B	−0.236986	0.279055	0.019120	0.039*
C18	−0.0617 (4)	0.23234 (19)	0.0923 (2)	0.0279 (9)
C19	0.0899 (4)	0.18070 (18)	0.1843 (2)	0.0243 (8)
C20	−0.0109 (4)	0.12876 (19)	0.1504 (2)	0.0280 (9)
C21	−0.0043 (4)	0.05951 (19)	0.1778 (2)	0.0314 (9)
H21	−0.072449	0.024329	0.154316	0.038*
C22	0.1054 (4)	0.0443 (2)	0.2404 (2)	0.0359 (10)
H22	0.113444	−0.002661	0.260660	0.043*
C23	0.2057 (4)	0.0961 (2)	0.2753 (3)	0.0382 (10)
H23	0.279570	0.083534	0.318870	0.046*
C24	0.2001 (4)	0.1647 (2)	0.2480 (2)	0.0356 (10)
H24	0.268694	0.199605	0.271730	0.043*
C25	0.4248 (4)	0.33511 (18)	0.2196 (2)	0.0246 (8)
C26	0.2643 (5)	0.3365 (2)	0.0444 (2)	0.0411 (11)
C27 ^a	0.4526 (10)	0.2907 (5)	0.1503 (5)	0.0196 (14)
H27A ^a	0.509596	0.321924	0.128771	0.024*
H27B ^a	0.511185	0.248353	0.171320	0.024*
C27 ^b	0.4153 (10)	0.2781 (5)	0.1700 (5)	0.0196 (14)
H27C ^b	0.327782	0.250541	0.163723	0.024*
H27D ^b	0.496597	0.245384	0.189502	0.024*
C28 ^a	0.3269 (7)	0.2649 (4)	0.0875 (4)	0.0204 (11)
H28A ^a	0.257068	0.241001	0.106518	0.024*
H28B ^a	0.356324	0.231049	0.055197	0.024*
C28 ^b	0.4148 (7)	0.3092 (4)	0.0950 (4)	0.0204 (11)
H28C ^b	0.448615	0.271925	0.067790	0.024*

Table 2: (continued)

Atom	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> _{iso} */ <i>U</i> _{eq}
H28D ^b	0.482006	0.349771	0.104165	0.024*
C29	0.6529 (5)	0.5039 (2)	0.3587 (2)	0.0454 (11)
H29A	0.599986	0.532994	0.316275	0.068*
H29B	0.732629	0.480966	0.348218	0.068*
H29C	0.688129	0.534561	0.402420	0.068*
C30	0.5114 (5)	0.2784 (4)	−0.0970 (3)	0.0788 (19)
H30A	0.505314	0.307451	−0.140611	0.118*
H30B	0.422805	0.252236	−0.105007	0.118*
H30C	0.589201	0.244113	−0.088428	0.118*
C31	−0.4254 (7)	0.0810 (6)	0.0464 (4)	0.173 (5)
H31A	−0.515824	0.062527	0.014271	0.259*
H31B	−0.442210	0.118898	0.078069	0.259*
H31C	−0.372033	0.041951	0.077159	0.259*
C32	0.4594 (4)	0.4982 (3)	0.1160 (2)	0.0477 (12)
H32A	0.366858	0.475068	0.097356	0.072*
H32B	0.447573	0.550377	0.112812	0.072*
H32C	0.504937	0.484324	0.167646	0.072*
N1	−0.1174 (3)	0.35652 (14)	0.08711 (15)	0.0203 (6)
N2	0.0154 (3)	0.35792 (14)	0.23932 (14)	0.0165 (6)
N3	−0.1805 (3)	0.35595 (14)	0.27041 (15)	0.0183 (6)
H3	−0.270865	0.354720	0.265559	0.022*
N4	0.0787 (3)	0.46250 (15)	0.14095 (15)	0.0193 (6)
N5	−0.0646 (3)	0.55125 (15)	0.08568 (15)	0.0217 (7)
H5	−0.138937	0.573146	0.055888	0.026*
N6	0.0550 (3)	0.24494 (15)	0.14593 (17)	0.0245 (7)
N7	−0.1068 (4)	0.16396 (16)	0.09182 (17)	0.0336 (8)
H7	−0.182998	0.145281	0.060194	0.040*
Ni1	0.10551 (5)	0.35324 (2)	0.15663 (3)	0.02142 (14)
O1	0.3085 (2)	0.35461 (14)	0.22464 (16)	0.0396 (7)
O2	0.5401 (2)	0.35651 (13)	0.26313 (12)	0.0253 (6)
O3	0.1566 (3)	0.35167 (13)	0.05988 (15)	0.0345 (7)
O4	0.2918 (3)	0.36258 (15)	−0.00957 (14)	0.0347 (7)
O5	0.5632 (3)	0.45025 (15)	0.37227 (14)	0.0376 (7)
H5B	0.543087	0.420215	0.337695	0.056*
O6	0.5354 (4)	0.3229 (4)	−0.0354 (3)	0.148 (3)
H6A	0.457870	0.335828	−0.031219	0.222*
O7	−0.3525 (5)	0.1073 (2)	0.00527 (18)	0.0793 (13)
H7B	−0.383861	0.090430	−0.037661	0.119*
O8	0.5417 (4)	0.4769 (3)	0.0743 (3)	0.118 (2)
H8A	0.492191	0.453989	0.037272	0.178*

^aOccupancy: 0.497(5), ^bOccupancy: 0.503(5).

and 2-(2'-pyridyl)-4, 4, 5, 5-tetramethylimidazoline-1-oxyl-3-oxide (NIT2Py) [6–8] as coligands, respectively. Two dinuclear nickel(II) complexes of ntb were prepared and characterized using dicarboxylate (terephthalate and fumarate) as bridging ligands by our group [9]. Succinic acid is a saturated aliphatic dicarboxylic acid. It shows conformational freedom and coordination versatility due to the single-bonded carbon chains, coordinating metal ions in various modes [10–13]. As an extension of our work, succinate was selected for the construction of a nickel(II) complex

of ntb. At this time, the effort to get a dinickel(II) complex failed, while a mononuclear complex was obtained.

The title complex was obtained by the reaction of $Ni(ClO_4)_2 \cdot 6H_2O$, ntb and disodium succinate hexahydrate in the ratio of 2:2:1 in methanol. The asymmetric unit of the title structure consists of one $[Ni(ntb)(suc)]$ unit and four methanol molecules. The central Ni(II) is six-coordinate in a N_4O_2 donor set formed by four nitrogen atoms of ntb and two oxygen atoms of succinate. The equatorial plane is defined by N2, N4, N6 of ntb and O3 of succinate. Equatorial bond angles are in the range $86.68(10)$ – $91.14(11)^\circ$. The apical positions are occupied by O1 of succinate and tertiary amine N1 of ntb with O1–Ni1–N1 bond angle to be 176.64° , showing a distorted octahedral environment. As expected, the tripodal ntb ligand coordinates nickel(II) with the three benzimidazole groups in a T-shaped fashion. This can be observed almost in all nickel(II) complexes of ntb [7–9, 14, 15]. The three benzimidazole Ni–N bond lengths are in the range of $2.054(3)$ – $2.077(3)$ Å, while the Ni–N(tertiary amine) bond length is $2.205(3)$ Å. The succinate anion shows a bidentate chelating mode. Each carboxylate provides one oxygen atom (O1, O3) to coordinate with nickel(II) to form a seven-membered ring. The two Ni–O bond lengths are $2.033(3)$ Å for Ni1–O1 and $2.067(2)$ Å for Ni1–O3, respectively. The O1–Ni–O3 bond angle is $95.16(11)^\circ$. This coordination is different to previously reported dicarboxylato-nickel(II) complexes of ntb. In the complexes $[Ni_2(ntb)_2(\mu\text{-tp})(H_2O)_2](NO_3)_2 \cdot 4CH_3OH \cdot H_2O$ and $[Ni_2(ntb)_2(\mu\text{-fum})(H_2O)(CH_3OH)](NO_3)_2 \cdot 6CH_3OH \cdot H_2O$, two $[Ni(ntb)]^{2+}$ units are bridged by terephthalate and fumarate in bis(monodentate) mode [7]. Thus, we showed that the single-bonded carbon chain of succinate is flexible enough to adopt the bidentate chelating mode.

In the complex, two of the $[Ni(ntb)(suc)]$ units are interlinked by H-bonds to form a dimolecular unit $[N5 \cdots H5 \cdots O4, d(D \cdots A) = 2.786(4)$ Å]. These dimolecular units are interacted to form a chain structure by H-bonds through the carboxylate (O2) of succinate and the benzimidazole $NH(N3)$ groups of ntb $[N3 \cdots H3 \cdots O2, d(D \cdots A) = 2.756(3)$ Å]. The chains are further expanded into a 2D network structure by weak π - π interaction between benzimidazole groups.

Author contributions: All the authors have accepted responsibility for the entire content of this submitted manuscript and approved submission.

Research funding: Scientific Research Base Development Program of the Beijing Municipal Commission of Education (KM201910028013).

Conflict of interest statement: The authors declare no conflicts of interest regarding this article.

References

- Agilent Technologies. *CrysAlis^{PRO}*; Agilent Technologies: Santa Clara, CA, USA, 2017.
- Dolomanov O. V., Bourhis L. J., Gildea R. J., Howard J. A. K., Puschmann H. OLEX2: a complete structure solution, refinement and analysis program. *J. Appl. Crystallogr.* 2009, 42, 339–341.
- Sheldrick G. M. A short history of SHELX. *Acta Crystallogr.* 2008, A64, 112–122.
- Thompson L. K., Ramaswamy B. S., Seymour E. A. Cobalt(II) and zinc(II) complexes of ‘tripod’ ligand tris(2-benzimidazolymethyl)amine. some five-coordinated derivatives and some with mixed stereochemistries. *Can. J. Chem.* 1977, 55, 878–888.
- Contreras R., Flores-Parra A., Migangos E., Téllez F., López-Sandoval H., Barba-Behrens N. From mono to polydentate azole and benzazole derivatives, versatile ligands for main group and transition metal atoms. *Coord. Chem. Rev.* 2009, 253, 1979–1999.
- Zhang C. X., Zhang Y. Y., Zhang H. L., Zhao Y., Sun Y. Q. Synthesis, structures and magnetic properties of nickel(II), manganic(II) and zinc(II) complexes containing pyridyl-substituted nitronyl nitroxide and tris(2-benzimidazolymethyl)amine. *Inorg. Chim. Acta.* 2009, 362, 5231–5236.
- Nie F. M., Lu F., Shang W. Y., Chen J. Synthesis and characterization of picolinato and nicotinato cobalt(II) complexes containing tris(2-benzimidazolymethyl)amine. *J. Coord. Chem.* 2010, 63, 1711–1719.
- Tian J. L., Xie M. J., Zhan Q., Yan S. P., Liao D. Z., Jiang Z. H. Syntheses, properties and X-ray structures of $[Ni(ntb)(N_3)_2] \cdot CH_3OH$ and $[Cu(ntb)(SCN)]Cl \cdot 3CH_3OH$ (ntb = tris(2-benzimidazolymethyl)amine). *J. Coord. Chem.* 2005, 58, 833–840.
- Nie F. M., Chen J., Lu F. Synthesis, crystal structures and magnetic studies of terephthalato- and fumarato-bridged dinickel(II) complexes with tripodal poly-benzimidazole ligand. *Inorg. Chim. Acta.* 2011, 365, 190–195.
- Zhou Z. H., Yang J. M., Wan H. L. Diamine substitution reactions of tetrahydrate succinato nickel, cobalt, and zinc coordination polymers. *Cryst. Growth Des.* 2005, 5, 1102–1108.
- Montney M. R., Mallika Krishnan S., Patel N. M., Supkowski R. M., LaDuca R. L. Divalent metal succinate/4,4'-dipyridylamine extended solids: from helical chains to diverse three-dimensional networks featuring double, quadruple, or self-interpenetration. *Cryst. Growth Des.* 2007, 7, 1145–1153.
- Ou G. C., Yuan X. Y., Li Z. Z. Synthesis and crystal structure of two succinato-bridged macrocyclic nickel(II) complexes. *Z. Anorg. Allg. Chem.* 2013, 639, 158–162.
- Choubey S., Roy S., Bhar K., Mitra P., Ribas J., Ghosh B. K. Varied dicarboxylate bridges in dinuclear trigonal prismatic manganese(II) and octahedral nickel(II) compounds containing tetradentate N-donor Schiff bases: syntheses, structures and magnetic behaviors. *Polyhedron* 2014, 74, 134–143.
- Li X. L., Luo Z. P., Zhang J. C., Nie F. M. Crystal structure of $\{[tris((1H\text{-benzo}[d]imidazolyl-2\text{-yl)methyl)amine-}\kappa^4 N,N',N'',N''']\text{-}(\text{nitrito-}\kappa^2 O,O') \text{nickel(II) perchlorate-ethanol (1/1), } C_{26}H_{27}ClN_8NiO_7\}$. *Z. Kristallogr. NCS* 2020, 235, 767–769.
- Ma J. C., Liu Y. Y., Yang J., Liu Y. Y., Ma J. F. A series of lanthanide-transition metal coordination polymers with mixed ligands: syntheses, structures, photoluminescence and magnetic properties. *CrystEngComm* 2011, 13, 3498–3505.