Raphael Wirthensohn and Maik Finze*

The crystal structure of trimethylsulfonium tris(trifluoromethylsulfonyl)methanide, $C_7H_9F_9O_6S_4$

Table 1: Data collection and handling.

Crystal:	Colourless block
Size:	0.50 × 0.33 × 0.28 mm
Wavelength:	Cu K α radiation (1.54184 Å)
μ :	6.57 mm ⁻¹
Diffractometer, scan mode:	XtaLAB Synergy,
$\theta_{\max}$, completeness:	77.1°, >99%
$N(hkl)_{\text{measured}}$, $N(hkl)_{\text{unique}}$, R_{int} :	106,974, 3454, 0.052
Criterion for I_{obs} , $N(hkl)_{\text{gt}}$:	$I_{\text{obs}} > 2 \sigma(I_{\text{obs}})$, 3451
$N(\text{param})_{\text{refined}}$:	239
Programs:	CrysAlis ^{PRO} [1], SHELX [2, 3], Diamond [4]

<https://doi.org/10.1515/ncrs-2020-0612>

Received November 23, 2020; accepted December 10, 2020;

published online January 8, 2021

Abstract

$C_7H_9F_9O_6S_4$, orthorhombic, $P2_12_12_1$ (no. 19), $a = 8.80180(10)$ Å, $b = 10.96580(10)$ Å, $c = 16.91360(10)$ Å, $V = 1632.48(3)$ Å³, $Z = 4$, $R_{\text{gt}}(F) = 0.0222$, $wR_{\text{ref}}(F^2) = 0.0604$, $T = 100$ K.

CCDC no.: 2049434

The molecular structure is shown in the figure. Table 1 contains crystallographic data and Table 2 contains the list of the atoms including atomic coordinates and displacement parameters.

Source of material

Trifluoromethylsulfonyl fluoride (0.2 g, 1.3 mmol) was transferred to diethyl ether (5 mL) in vacuum at -196 °C.

***Corresponding author: Maik Finze**, Institut für Anorganische Chemie, Julius-Maximilians-Universität Würzburg, Am Hubland, D-97074 Würzburg, Germany; and Institute for Sustainable Chemistry & Catalysis with Boron (ICB), Julius-Maximilians-Universität Würzburg, Am Hubland, D-97074 Würzburg, Germany, E-mail: maik.finze@uni-wuerzburg.de. <https://orcid.org/0000-0002-6098-7148>

Raphael Wirthensohn, Institut für Anorganische Chemie, Julius-Maximilians-Universität Würzburg, Am Hubland, D-97074 Würzburg, Germany; and Institute for Sustainable Chemistry & Catalysis with Boron (ICB), Julius-Maximilians-Universität Würzburg, Am Hubland, D-97074 Würzburg, Germany

The mixture was warmed to 0 °C and a solution of MeMgCl in THF (1.52 mL, 3 mol L⁻¹, 4.55 mmol) was added. The reaction mixture was allowed to warm to room temperature over the course of 12 h. A few crystals of the title compound were obtained by slow evaporation of the reaction mixture within 14 days.

Experimental details

Hydrogen atoms were located from difference Fourier maps. For the refinement, a riding model was applied and their U_{iso} values were set to 1.5 U_{eq} of the parent atoms. Using a hole-in-one method a Flack parameter of 0.459(16) [3] was calculated. Using Parsons' method [5] using 1456 selected quotients the parameter was very similar [0.461(8)]. The structure was finally refined as a 2-component inversion twin with a refined twin ratio of 0.459(6)/0.541(6).

Comment

Weakly coordinating anions (WCAs) have a low tendency to coordinate because the negative charge is delocalized and they are chemically and electrochemically robust [6–11]. Thus, WCAs have been applied for the stabilization of highly reactive cations and they serve as important

Table 2: Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²).

Atom	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> _{iso} [*] / <i>U</i> _{eq}
C1_1	0.3067 (3)	0.3541 (2)	0.42660 (14)	0.0249 (5)
S1_1	0.35961 (6)	0.49696 (5)	0.37304 (3)	0.02142 (13)
F1_1	0.42950 (19)	0.29494 (15)	0.45107 (11)	0.0359 (4)
O1_1	0.3952 (2)	0.57813 (18)	0.43680 (11)	0.0280 (4)
C2_1	0.4757 (3)	0.4897 (2)	0.15300 (14)	0.0279 (5)
S2_1	0.50395 (6)	0.38004 (5)	0.23488 (3)	0.02170 (13)
F2_1	0.22620 (19)	0.38664 (16)	0.48996 (9)	0.0346 (4)
O2_1	0.2323 (2)	0.52015 (17)	0.32299 (10)	0.0259 (4)
C3_1	0.7838 (3)	0.3973 (2)	0.40938 (15)	0.0270 (5)
S3_1	0.69775 (7)	0.51786 (5)	0.34747 (3)	0.02185 (13)
F3_1	0.22247 (19)	0.27995 (15)	0.38364 (9)	0.0309 (3)
O3_1	0.3649 (2)	0.31357 (17)	0.23848 (11)	0.0281 (4)
C4_1	0.5214 (3)	0.4663 (2)	0.31936 (14)	0.0214 (4)
O4_1	0.6429 (2)	0.31989 (17)	0.21490 (11)	0.0285 (4)
F4_1	0.3499 (2)	0.55323 (18)	0.16235 (10)	0.0430 (4)
O5_1	0.7951 (2)	0.52201 (17)	0.27979 (10)	0.0268 (4)
F5_1	0.4628 (3)	0.42691 (18)	0.08633 (10)	0.0449 (5)
O6_1	0.6911 (2)	0.61938 (17)	0.39992 (11)	0.0283 (4)
F6_1	0.5911(2)	0.56497 (18)	0.14605 (11)	0.0454 (5)
F7_1	0.7794 (2)	0.28927 (14)	0.37505 (10)	0.0351 (4)
F8_1	0.92796 (19)	0.42713 (18)	0.42247 (11)	0.0399 (4)
F9_1	0.71355 (18)	0.38993 (16)	0.47885 (9)	0.0321 (3)
C1_2	0.0209 (3)	0.2908 (3)	0.22124 (15)	0.0297 (5)
H1A_2	−0.089631	0.289526	0.228302	0.044*
H1AB_2	0.066205	0.347712	0.259168	0.044*
H1AC_2	0.061834	0.208845	0.230161	0.044*
S1_2	0.06499 (7)	0.33877 (5)	0.12289 (4)	0.02495 (13)
C2_2	−0.0184 (4)	0.4872 (2)	0.12157 (17)	0.0349 (6)
H2A_2	−0.013737	0.520471	0.067837	0.052*
H2AB_2	0.037455	0.540894	0.157657	0.052*
H2AC_2	−0.124672	0.481822	0.138490	0.052*
C3_2	−0.0694 (4)	0.2522 (3)	0.06682 (16)	0.0345(6)
H3A_2	−0.067806	0.279482	0.011649	0.052*
H3AB_2	−0.171373	0.264242	0.088786	0.052*
H3AC_2	−0.042768	0.165545	0.069320	0.052*

building blocks for many applications in materials science. For example, they are used as components of the electrolytes applied in electrochemical devices such as batteries, supercapacitors, and dye-sensitized solar cells (DSSCs). The tris(trifluoromethylsulfonyl)methanide anion ([Tf₃C][−]) is a WCA that is accessible from trifluoromethylsulfonyl fluoride and methylmagnesium bromide [12]. So far, only few crystallographic studies on salts of the [Tf₃C][−] anion with different types of counterions have been described [12–15].

In the course of our studies on reactions of trifluoromethylsulfonyl fluoride with methylmagnesium bromide single crystals of the title compound trimethylsulfonium tris(trifluoromethylsulfonyl)methanide ([SMe₃][Tf₃C]) have

been obtained. The [SMe₃]⁺ cation formed during storage of a reaction mixture for 14 days.

The asymmetric unit of the title compound contains one [Tf₃C][−] anion and one [SMe₃]⁺ cation. The C–S₃ unit of the anion is planar. The SO₂CF₃ groups have different orientations with respect to the central C–S₃ core with two groups facing up and one facing down with regard to the CS₃ plane. The bond lengths and angles are similar to those reported earlier for the [Tf₃C][−] anion [12–15].

The [SMe₃]⁺ cation reveals the typical pyramidal structure and the bond parameters are close to values reported for other trimethylsulfonium salts, e.g. [SMe₃][1-H₂N-6-F-*closo*-1-CB₁₁I₁₀] [16].

Author contribution: All the authors have accepted responsibility for the entire content of this submitted manuscript and approved submission.

Research funding: This publication was supported by the Open Access Publication Fund of the University of Wuerzburg.

Conflict of interest statement: The authors declare no conflicts of interest regarding this article.

References

1. Oxford Diffraction Ltd. *CrysAlis^{PRO}*; Yarnton: Oxfordshire, England, 2020.
2. Sheldrick G. M. SHELXT – Integrated space-group and crystal-structure determination. *Acta Crystallogr.* 2015, *A71*, 3–8.
3. Sheldrick G. M. Crystal structure refinement with *SHELXL*. *Acta Crystallogr.* 2015, *C71*, 3–8.
4. Brandenburg K. *DIAMOND. Visual crystal structure information system* (Ver. 4.5.1); Crystal Impact GbR: Bonn, Germany, 2018.
5. Parsons S., Flack H. D., Wagner T. Use of intensity quotients and differences in absolute structure refinement. *Acta Crystallogr.* 2013, *B69*, 249–259.
6. Beck W., Sünkel K. Metal complexes of weakly coordinating anions. Precursors of strong cationic organometallic Lewis acids. *Chem. Rev.* 1988, *88*, 1405–1421.
7. Seppelt K. “Noncoordinating” anions, II. *Angew. Chem. Int. Ed. Engl.* 1993, *32*, 1025–1027; *Angew. Chem.*, 1993, *105*, 1074–1077.
8. Strauss S. H. The search for larger and more weakly coordinating anions. *Chem. Rev.* 1993, *93*, 927–942.
9. Krossing I., Raabe I. Noncoordinating anions – fact or fiction? A survey of likely candidates. *Angew. Chem. Int. Ed.* 2004, *43*, 2066–2090. *Angew. Chem.*, 2004, *117*, 2116–2142.
10. Riddlestone I. M., Kraft A., Schaefer J., Krossing I. Taming the cationic beast: novel developments in the synthesis and application of weakly coordinating anions. *Angew. Chem. Int. Ed.* 2018, *57*, 13982–14024. *Angew. Chem.*, 2018, *130*, 14178–14221.
11. Huang Z., Wang S., Dewhurst R. D., Ignat’ev N. V., Finze M., Braunschweig H. Boron: its role in energy related research and applications. *Angew. Chem. Int. Ed.* 2020, *59*, 8800–8816. *Angew. Chem.*, 2020, *132*, 8882–8900.
12. Turowsky L., Seppelt K. Tris[(trifluoromethyl)sulfonyl]methane, $\text{C}(\text{SO}_2\text{CF}_3)_3$. *Inorg. Chem.* 1988, *27*, 2135–2137.
13. Schlueter J. A., Geiser U., Wang H. H., Aravinda M. K., Ward B. H., Parakka J. P., Daugherty R. G., Kelly M. E., Nixon P. G., Winter R. W., Gard G. L., Montgomery L. K., Koo H. -J., Whangbo M. -H. Trifluoromethylsulfonyl-based salts of BEDT-TTF: crystal and electronic structures and physical properties. *J. Solid State Chem.* 2002, *168*, 524–534.
14. Waller F. J., Barrett A. G. M., Braddock D. C., Ramprasad D., McKinnell R. M., White A. J. P., Williams D. J., Ducray R. Tris(trifluoromethanesulfonyl)methide (“triflide”) anion: convenient preparation, X-ray crystal structures, and exceptional catalytic activity as a counterion with ytterbium(III) and scandium(III). *J. Org. Chem.* 1999, *64*, 2910–2913.
15. Schulz C., Daniels J., Bredow T., Beck J. The electrochemical synthesis of polycationic clusters. *Angew. Chem. Int. Ed.* 2016, *55*, 1173–1177. *Angew. Chem.*, 2016, *128*, 1188–1192.
16. Finze M., Reiss G. J. Trimethylsulfonium 1-amino-6-fluoro-2,3,4,5,7,8,9,10,11,12-decaiodo-1-carba-closo-dodecaborate. *Acta Crystallogr.* 2012, *E68*, o640.