Daniel M. Orang'o, Martin O. Onani*, Paul K. Tarus, Samuel T. Lutta and Roger A. Lalancette

Crystal structure of 2,2'-(propane-1,3-dilylbis(azaneylylidene))bis(methanylylidene)bis(4-methylphenol), $C_{19}H_{22}N_2O_2$

<https://doi.org/10.1515/ncrs-2021-0279>

Received July 10, 2021; accepted August 16, 2021;

published online August 31, 2021

Abstract

$C_{19}H_{22}N_2O_2$, monoclinic, $P2_1/c$ (no. 14), $a = 19.3063(4)$ Å, $b = 5.83200(10)$ Å, $c = 14.7996(3)$ Å, $\beta = 92.715(1)^\circ$, $V = 1664.48(6)$ Å³, $Z = 4$, $R_{gt}(F) = 0.0423$, $wR_{ref}(F^2) = 0.1102$, $T = 100(2)$ K.

CCDC no.: 1911770

The molecular structure is shown in the figure. Table 1 contains crystallographic data and Table 2 contains the list of the atoms including atomic coordinates and displacement parameters.

Table 1: Data collection and handling.

Crystal:	Yellow block
Size:	$0.44 \times 0.39 \times 0.33$ mm
Wavelength:	Cu $K\alpha$ radiation (1.54178 Å)
μ :	0.64 mm^{-1}
Diffractometer, scan mode:	Bruker SMART Apex-II, φ and ω
$\theta_{\max}$, completeness:	66.8° , 96%
$N(hkl)_{\text{measured}}$, $N(hkl)_{\text{unique}}$:	17,078, 2818, 0.041
R_{int} :	
Criterion for I_{obs} , $N(hkl)_{\text{gt}}$:	$I_{\text{obs}} > 2 \sigma(I_{\text{obs}})$, 2575
$N(\text{param})_{\text{refined}}$:	213
Programs:	Bruker [1], SHELX [2, 3], Diamond [4]

Source of material

The salicylaldimine compound was synthesized and obtained in good yields from a 2:1 reaction ratio of substituted salicylaldehyde 2-hydroxy-5-methylbenzaldehyde with 1,3-diaminopropane in dry methanol under reflux. The compound was obtained as a yellow solid with yields of 96%.

Experimental details

Single crystals suitable for X-ray diffraction (XRD) studies for the compound were obtained after four weeks by slow diffusion and evaporation of hexane into a concentrated solution of the compound in dichloromethane (DCM). Crystal evaluation and data collection were done on a Bruker Smart APEX2 diffractometer [1] Crystal evaluation and data collection were done on a Bruker SMART 1000

***Corresponding author: Martin O. Onani**, Department of Chemistry, Faculty of Natural Sciences, University of the Western Cape, Private Bag X17, Bellville 7535, Republic of South Africa, E-mail: monani@uwc.ac.za. <https://orcid.org/0000-0002-4735-3669>

Daniel M. Orang'o, Department of Chemistry, Faculty of Natural Sciences, University of the Western Cape, Private Bag X17, Bellville 7535, Republic of South Africa; and Department of Chemistry and Biochemistry, School of Science, University of Eldoret, P. O. Box 1125–30100 Eldoret, Kenya

Paul K. Tarus and Samuel T. Lutta, Department of Chemistry and Biochemistry, School of Science, University of Eldoret, P. O. Box 1125–30100 Eldoret, Kenya

Roger A. Lalancette, Department of Chemistry, Rutgers State University, 73 Warren St. Rutgers, Newark, NJ 07102, USA

Table 2: Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²).

Atom	x	y	z	<i>U</i> _{iso} */ <i>U</i> _{eq}
C1	0.81206 (8)	0.6886 (3)	0.30440 (10)	0.0287 (3)
C2	0.87955 (8)	0.7683 (2)	0.29892 (10)	0.0327 (4)
H2A	0.887618	0.914471	0.273133	0.039*
C3	0.93485 (8)	0.6371 (3)	0.33055 (10)	0.0296 (4)
H3	0.980550	0.694931	0.325968	0.036*
C4	0.92561 (7)	0.4208 (2)	0.36927 (9)	0.0256 (3)
C5	0.98661 (8)	0.2776 (3)	0.40219 (11)	0.0326 (4)
H5A	0.971548	0.118938	0.410774	0.049*
H5B	1.005590	0.339013	0.459826	0.049*
H5C	1.022393	0.281639	0.357412	0.049*
C6	0.85811 (7)	0.3428 (2)	0.37430 (9)	0.0248 (3)
H6	0.850499	0.196294	0.400062	0.030*
C7	0.80083 (7)	0.4712 (2)	0.34299 (9)	0.0243 (3)
C8	0.73110 (7)	0.3780 (3)	0.34833 (10)	0.0287 (3)
H8	0.725134	0.231154	0.374611	0.034*
C9	0.60952 (8)	0.3859 (3)	0.32523 (13)	0.0438 (5)
H9A	0.613306	0.249316	0.364905	0.053*
H9B	0.592026	0.334995	0.264545	0.053*
C10	0.55888 (8)	0.5540 (3)	0.36317 (11)	0.0322 (4)
H10A	0.554359	0.688536	0.322515	0.039*
H10B	0.577304	0.608376	0.422955	0.039*
C11	0.48792 (8)	0.4486 (3)	0.37346 (12)	0.0371 (4)
H11A	0.470656	0.384241	0.314726	0.044*
H11B	0.491539	0.321829	0.417903	0.044*
C12	0.37444 (7)	0.5928 (3)	0.38494 (10)	0.0267 (3)
H12	0.359468	0.460389	0.352108	0.032*
C13	0.32289 (7)	0.7585 (2)	0.41226 (9)	0.0242 (3)
C14	0.34274 (8)	0.9615 (3)	0.45794 (11)	0.0319 (4)
C15	0.29194 (9)	1.1161 (3)	0.48124 (13)	0.0381 (4)
H15	0.304830	1.253449	0.512165	0.046*
C16	0.22283 (8)	1.0720 (3)	0.45988 (11)	0.0324 (4)
H16	0.188915	1.180246	0.476460	0.039*
C17	0.20162 (7)	0.8730 (2)	0.41466 (10)	0.0256 (3)
C18	0.12648 (8)	0.8250 (3)	0.39069 (11)	0.0316 (4)
H18A	0.117236	0.849437	0.325719	0.047*
H18B	0.097430	0.928573	0.424766	0.047*
H18C	0.115783	0.665793	0.405955	0.047*
C19	0.25266 (7)	0.7184 (2)	0.39178 (9)	0.0248 (3)
H19	0.239309	0.581013	0.361196	0.030*
N1	0.67781 (7)	0.4894 (2)	0.31845 (9)	0.0340 (3)
N2	0.43897 (6)	0.6204 (2)	0.40381 (9)	0.0291 (3)
O1	0.75882 (6)	0.8209 (2)	0.27355 (8)	0.0400 (3)
H1	0.721521	0.747671	0.276516	0.060*
O2	0.41009 (6)	1.0089 (2)	0.47933 (10)	0.0473 (4)
H2	0.435110	0.903102	0.460415	0.071*

CCD diffractometer with Mo K α radiation ($\lambda = 0.71073$ Å) equipped with an Oxford Cryostream low temperature apparatus operating at 100(1) K. The structure was solved by direct method using the SHELXS [2] program and refined with SHELXL [3].

All hydrogen atoms were placed in idealized positions and refined in riding models with U_{iso} assigned the values

of 1.2 times or 1.5 times those of their parent atoms and the distances of C–H were constrained to 0.95 Å for all the aromatic H atoms, 0.98 Å for the CH₃ protons and 0.9 for CH₂ protons or 0.84 Å for the hydroxy group H atoms, respectively. The idealized tetrahedral OH refined as rotating group. The visual crystal structure information was performed using Diamond [4].

Comment

N,O-salicylaldimines are Schiff base ligands with the ability to coordinate transition metal ions *via* the hard nitrogen and oxygen donor atoms. This leads to a stabilization of the metal complexes against reduction and eventually give good thermal stability [5]. It is also easy to manipulate their steric and electronic properties. The early transition metal complexes have been shown to exhibit high oxophilic nature [6, 7]. Despite this shortcoming, a lot of research has been conducted on salicylaldimine complexes and their application as catalysts in polymerization reactions. For instance, group four metal-complexes of titanium and zirconium have been synthesized and tested, which showed significant (co) polymerization of ethylene [8–10]. Their salicylaldimine complexes have also been shown to exhibit stability, catalytic activity and greater tolerance for polar monomers in olefin polymerization [11].

The asymmetric unit of the title compound contains one salicylaldimine molecule. In the structure, the 2-(imino-methyl)-4-methylphenol moieties are on either sides of the propyl linker and enclose a dihedral angle of 52.40(3) comparable to that of in the structure of 4,4'-Dichloro-2,2'-[(1*E*,1'*E*)-propane-1,3-diylbis(nitrilomethylidyne)]diphenol [12].

In the crystal, this compound displays two intramolecular hydrogen bonds O–H $\cdots$ N with an N $\cdots$ O distance of 2.593(2) Å. The bond distances and angles are comparable to those of related compounds [13–16].

Author contributions: All the authors have accepted responsibility for the entire content of this submitted manuscript and approved submission.

Research funding: We sincerely appreciate the funding from the University of Eldoret Research grant of 2016 and the University of the Western Cape as well as the National Research Foundation of South Africa (Grant number 105894).

Conflict of interest statement: The authors declare no conflicts of interest regarding this article.

References

1. Bruker. APEXII; Bruker AXS Inc: Madison, Wisconsin, USA, 2009.
2. Sheldrick G. M. *SHELXTL* – integrated space-group and crystal-structure determination. *Acta Crystallogr.* 2015, *A71*, 3–8.
3. Sheldrick G. M. Crystal structure refinement with SHELXL. *Acta Crystallogr.* 2015, *C71*, 3–8.
4. Brandenburg K. *DIAMOND*. Visual Crystal Structure Information System (Ver. 4.0); Crystal Impact: Bonn, Germany, 2015.
5. Li C. Q., Wang F. F., Gao R., Sun P., Zhang N., Wang J. Bidentate iron complexes based on hyperbranched salicylaldimine ligands and their catalytic behavior toward ethylene oligomerization. *Transition Met. Chem.* 2017, *42*, 339–346.
6. Britovsek G. J. P., Bruce M., Gibson V. C., Kimberley B. S., Maddox P. J., Mastroianni S., McTavish S. J., Redshaw C., Solan G. A., Stromberg S., White A. J. P., Williams D. J. Iron and cobalt ethylene polymerization catalysts bearing 2,6-bis(imino)pyridyl ligands: synthesis, structures, and polymerization studies. *J. Am. Chem. Soc.* 1999, *121*, 8728–8740.
7. Gibson V. C., Spitzmesser S. K. Advances in non-metallocene olefin polymerization catalysis. *Chem. Rev.* 2003, *103*, 283–315.
8. Salata M. R., Marks T. J. Synthesis, characterization, and marked polymerization selectivity characteristics of binuclear phenoxyiminato organozirconium catalysts. *J. Am. Chem. Soc.* 2008, *130*, 12–13.
9. Salata M. R., Marks T. J. Catalyst nuclearity effects in olefin polymerization. enhanced activity and comonomer enchainment in ethylene + olefin copolymerizations mediated by bimetallic group 4 phenoxyiminato catalysts. *Macromolecules* 2009, *42*, 1920–1933.
10. Han S., Yao E., Qin W., Zhang S., Ma Y. Binuclear heteroligated titanium catalyst based on phenoxyimine ligands: synthesis, characterization, and ethylene (co)polymerization. *Macromolecules* 2012, *45*, 4054–4059.
11. Suo H., Solan G. A., Ma Y., Sun W. H. Developments in compartmentalized bimetallic transition metal ethylene polymerization catalysts. *Coord. Chem. Rev.* 2018, *372*, 101–116.
12. Lei S., Zhu-Ping X., Zhong Z., Zhao-Zhao Z., Hai-Liang Z. Integrated space-group and crystal-structure determination. *Acta Crystallogr.* 2007, *E63*, o4726.
13. Nazir H., Arici C., Emregül K. C., Atakol O. A crystallographic and spectroscopic study on the imine-amine tautomerism of 2-hydroxyaldimine compounds. *Z. für Kristallogr. - Cryst. Mater.* 2006, *221*, 699–704.
14. Xue L. W., Li X. W., Zhao G. Q., Yang W. C. Synthesis, crystal structure, and thermal analysis of a dioxomolybdenum(VI) complex derived from *N,N'*-bis(5-methylsalicylidene)-1,3-diaminopropane. *Synth. React. Inorg. Metal-Org. Nano-Metal Chem.* 2013, *43*, 1514–1517.
15. Steiner T. The hydrogen bond in the solid state. *Angew. Chem. Int. Ed.* 2002, *41*, 48–76.
16. Bilge S., Kiliç Z., Hayvali Z., Hökelek T., Safran S. Intramolecular hydrogen bonding and tautomerism in Schiff bases: part VI. syntheses and structural investigation of salicylaldimine and naphthaldimine derivatives. *J. Chem. Sci.* 2009, *121*, 989–1001.