NOTIZEN

 ^1H and ^{13}C NMR Spectroscopic Evidence for the Complexation of Potassium Cations by Cyclohexanedioxydiacetamides

Whei Oh Lin

Seção de Química, Instituto Militar de Engenharia,
Praia Vermelha, URCA, Rio de Janeiro, RJ,
Brasil, 22290

Helmut G. Alt*

Laboratorium für Anorganische Chemie,
Universität Bayreuth, D-8580 Bayreuth,
Federal Republic of GermanyZ. Naturforsch. **40b**, 441–442 (1985);
received October 29, 1984Noncyclic, Ionophors, Complexation of K^+ Cations,
 ^1H NMR Spectra, ^{13}C NMR SpectraThe complexation of cyclohexanedioxydiacetamides by K^+ cations may be studied by ^1H and ^{13}C NMR spectroscopy.

Neutral compounds featuring ether and amide groups (I, II) are potential chelating ligands for group IIA cations [1–3]. Recently we have studied the binding modes of ligands I by using ^1H and ^{13}C NMR spectroscopy and found that the ether and the carbonyl oxygens act as chelating centers [4]. We

now report ^1H and ^{13}C NMR spectroscopic evidence for the complexation of potassium cations by compounds II. So far such a complexation has neither been observed by direct UV titration nor by picrate extraction methods [3].

Analogous to the ionophors I the *trans*-cyclohexanedioxydiacetamides II are able to coordinate Ca^{2+} and K^+ cations to form preferably 2:1 complexes. Upon complexation the ^1H NMR spectra change typically: in the ^1H NMR spectrum of free *trans*-cyclohexanedioxydiacetic acid morpholinoamide (in acetone- d_6 , at room temperature) the methylene protons H_a and H_b appear as a singlet at δ 4.27 ppm. The addition of K^+ cations at room temperature results in a shift to lower field and a splitting of the former singlet into an AB pattern (Fig. 1).

Fig. 1. ^1H NMR signals of the methylene protons H_a and H_b of 1,2-cyclohexanedioxydiacetic acid morpholinoamide and its K^+ complex in acetone- d_6 .

* Reprint requests to Priv.-Doz. Dr. H. G. Alt.
0340-5087/85/0300-0441/\$ 01.00/0

A downfield shift is expected upon complexation [4, 5]. An AB pattern for H_a and H_b should be observed even in the free ligand because the neighbouring cyclohexane ring renders these hydrogens diastereotopic. However, the appearance of an AB pattern at room temperature indicates a conformation change upon complexation. Studies with molecular models show that the chelate complexation locks the cyclohexane ring in the equatorial conformation that is already favored in the free ionophor (*ca.* 99%).

This interpretation is confirmed by the fact that the splitting of the methylene protons H_a and H_b cannot be observed upon complexation of I.

The complexation of II by K^+ cations can also be confirmed by ^{13}C NMR spectroscopy. The three carbon atoms that are directly bonded to the two oxygen atoms of a side chain are shifted to lower field ($\Delta\delta \approx 3$ ppm) while the signals of all the other carbons remain nearly unchanged. Independent of the ratio $[II]/[K^+]$ we only observe one species in the 1H and ^{13}C NMR spectra. This is indicative of a fast exchange mechanism between free K^+ cations and K^+/II -complexes. It is well documented that the NMR spectrum of a complex is dependent on the lifetime of the individual metal-ligand bonds [6].

The complexation of ligands of the type II with monovalent cations has been observed for the first time. Our studies have shown that NMR spectroscopy is a valuable tool for the determination of coordination modes of neutral ionophors, and for gaining information about intramolecular processes in ionophor-cation complexes.

We thank the Volkswagen Foundation for financial support and Prof. Dr. H. Gerlach for helpful discussions.

- [1] D. Ammann, E. Pretsch, and W. Simon, *Helv. Chim. Acta* **56**, 1780 (1973).
- [2] J. Readio, I. J. Borowitz, N. Pollack, J. Porter, L. Weiss, and G. B. Borowitz, *J. Coord. Chem.* **11**, 135 (1981).
- [3] I. J. Borowitz, W. O. Lin, T. C. Wun, R. Bittmann, L. Weiss, V. Diakiw, and G. B. Borowitz, *Tetrahedron* **33**, 1697 (1977).
- [4] W. O. Lin and H. G. Alt, *Z. Naturforsch.* **39b**, 1375 (1984).
- [5] a) R. U. Lemieux and J. W. Lown, *Can. J. Chem.* **42**, 893 (1964);
b) J. L. Sudmeier and C. N. Reilley, *Anal. Chem.* **36**, 1707 (1964).
- [6] R. J. Day and C. N. Reilley, *Anal. Chem.* **37**, 1326 (1965).