Organotin-Induced Hyperglycemia in the Crab, *Oziotelphusa senex senex* Fabricius

G. Purna Chandra Nagaraju, Md. Riyaz Basha and P. Sreenivasula Reddy*

Department of Biotechnology, Sri Venkateswara University, Tirupati - 517 502, India.

Fax: +91-8574-49601. E-mail: psreddy@vsnl.com

* Author for correspondence and reprint requests

Z. Naturforsch. **56c**, 315–317 (2001);
received March 21/November 13, 2000

Hyperglycemia, *Oziotelphusa senex senex*, Organotins

Injection of three different organotin compounds such as tripalmitin, fentin and fenbutatin produced a significant increase in the hemolymph sugar level of intact crabs at *Oziotelphusa senex senex* apparently by stimulating release of the hyperglycemic hormone (HGH).

Organotins and their degradation products which leach out from the painted surfaces occur at detectable levels in sea water, sediments, estuaries and harbours posing threat to non-target species (Hodge *et al.*, 1979; Margaillan *et al.*, 1985; Langston *et al.*, 1987). Even though organotins are found to be neurotoxic to rats (Desaiah and Ho, 1979; Ali *et al.*, 1986), the molecular mechanisms of toxic action are not fully understood. These compounds have been shown to inhibit oxidative phosphorylation (Aldridge *et al.*, 1977), depress thymus dependent immunity (Seinen *et al.*, 1979), and inhibit basal as well as calmodulin activated Ca^{2+} ATPase in rat synaptosomes (Rao *et al.*, 1987).

In decapod Crustaceans, hemolymph sugar level is regulated by a hyperglycemic hormone (HGH). Abramowitz *et al.* (1944) were the first to determine the presence of hemolymph sugar regulating principle in *Callinectes* and called as diabetogenic factor. Since then, a good amount of work has been carried out on the chemical nature, mode and site of action of the HGH (Keller *et al.*, 1985; Sedlmeier, 1985; Liu *et al.*, 1998; Sithigorngul *et al.*, 1999). The amino acid composition of HGH from several crustaceans has been reported (Kleinholz, 1975; Keller and Wunderer, 1978; Keller, 1981; Newcomb, 1983; Huberman and Aguilar, 1986; Soye *et al.*, 1990; Yasuda *et al.*, 1994; Ohira *et al.*, 1997; Marco *et al.*, 1998). Recently, the gene for

HGH was also cloned from several crustaceans (Reddy and Ramamurthi, 1999). We have determined the expression of HGH gene at different molt and reproductive stages (Reddy *et al.*, 1997). HGH, produced by neurosecretary cells belonging to the X-organ, stored in and released from the sinus gland located in the eyestalks, is the major hormonal regulator of carbohydrate metabolism in crustacea.

Although there has been considerable work on HGH in a number of species, data are limited for the factors regulating the release of HGH. The release of HGH is influenced by external factors such as photoperiod, temperature, nutrition and stress (Kleinholz and Keller, 1979; Van Herp and Payen, 1991) as well as internal factors (Fingerman, 1995; 1997). We have demonstrated earlier that xenobiotics such as sumithion and hexachlorocyclohexane induces hyperglycemia in the crab, *Oziotelphusa* and hypothesised that these compounds produce this effect indirectly by stimulating release of HGH. The objectives of the present study are (a) to test our hypothesis generated earlier that xenobiotics induce hyperglycemia in decapod crustaceans by using organotin compounds and (b) to determine for the first time what effect, if any, organotin compounds have on the hemolymph sugar level in *Oziotelphusa*, *in vivo*.

Adult, inter molt (stage C₄) crabs of *Oziotelphusa senex senex* were collected from paddy fields and irrigation canals (Tirupati). They were acclimatized to the laboratory conditions (temperature $27 \pm 1^\circ\text{C}$; RH 75% and a light period of 12 hours) for one week before being used for experimentation. Only male crabs weighing 30–32 g were used in the present study. They were fed daily once *ad libitum* sheep meat and medium was changed 2 h after feeding. Feeding was stopped 24 h before commencing the experiment to overcome differences, if any, due to differential feeding.

Crabs were divided into 10 batches of 9 animals each. The first batch served as control and received no treatment. The second batch of crabs received 10 μl of 10% ethanol and served as controls. The 3rd, 4th and 5th batches were injected with 10 μmoles of tripalmitin, fentin and fenbutatin, respectively, through the arthrodial mem-

brane of the coxa of 3rd pair of walking legs in 10 µl volume. The organotin compounds obtained from Environmental Protection Agency, Research Triangle Park, USA through a generous gift by Dr. K. S. P.Rao were dissolved in small amount of ethanol and diluted in crustacean saline.

The eyestalks were removed (ESX) by cutting off the organs at the base without prior ligation but with cautery of wound after operation from the remaining batches. Crabs in batch 6 were not

Table I. Hemolymph sugar level in the crab after different treatments.

	Hemolymph sugar level (mg/100 ml)
Intact crabs	33.29 ± 1.24
Ethanol injected intact crabs	32.63 ± 3.88 (- 1.98)
Tripalmitin injected intact crabs	53.83 ± 2.49 ^a (61.71)
Fentin injected intact crabs	51.16 ± 2.14 ^a (53.68)
Fenbutatin injected intact crabs	39.40 ± 4.16 ^a (18.35)
Eyestalkless crabs (ESX)	22.15 ± 1.83 ^a (- 33.46)
Ethanol injected ESX crabs	21.11 ± 1.94 ^{a,b} (- 4.70)
Tripalmitin injected ESX crabs	21.07 ± 1.36 ^{a,b} (- 4.87)
Fentin injected ESX crabs	20.83 ± 2.17 ^{a,b} (- 5.95)
Fenbutatin injected ESX crabs	21.21 ± 2.56 ^{a,b} (- 4.24)
F ratio	227.37
p value	< 0.0001

Values are Mean ± S. D. of 9 individual observations. Values in paranthesis are percent change from intact crabs. For calculation of % change for intact-injected and ESX crabs, intact crabs served as controls; For ESX-injected crabs, ESX crabs served as control.

^a Values are significant at $p < 0.0001$ from intact crabs.

^b Values are not significant from ESX crabs.

given any treatment and used after 24 h post-ablation. Crabs in batch 7 were injected with 10 µl 10% ethanol and served as ESX control. The crabs in batches 8–10 were injected with 10 µmoles of tripalmitin, fentin and fenbutatin, respectively, in 10 µl volume, after 24 h of ESX. The hemolymph samples were collected 2 h after the injection at the same time of the day to avoid influence of circadian rhythm. The hemolymph sugar level was determined following the method of Carroll *et al.* (1956) using anthrone reagent. The data were analysed by using ANOVA and Student Newman-Keul (SNK) test (Steel and Torrie, 1960).

Bilateral ESX significantly ($P < 0.0001$) decreased (-33.46%) hemolymph sugar level (Table I). The hemolymph sugar levels increased significantly ($p < 0.0001$) after injection of organotin compounds into intact animals. Among the three organotins, tripalmitin produced most elevation (61.71%) in hemolymph sugar level. On the contrary, injection of organotin compounds into ESX crabs did not affect the hemolymph sugar level.

Organotin compounds might have induced hyperglycemia in the intact crabs in several different ways such as (a) by triggering release of HGH, (b) by mimicking the action of this hormone or (c) even by directly stimulating glycogenolysis by activating the enzyme phosphorylase. However, given the fact that organotin compounds were not able to produce hyperglycemia in ESX crabs, we favour the hypothesis that organotins exerted its effect by stimulating release of HGH from the sinus glands of eyestalks. This also supports the hypothesis, that the sinus glands in the eyestalks of *Oziotelphusa* are the main release site for HGH.

Acknowledgements

This research work was funded by grants from DST (SP/SO/CO4/96) and ICAR (4(65)97-ASR-1) to PSR.

- Abramowitz A. A., Hisaw F. L. and Papendrea D. V. (1944), The occurrence of a diabetogenic factor in the eyestalks of crustaceans. *Biol. Bull.* **86**, 1–5.
- Aldridge W. N., Street B. W. and Skilleter D. N. (1977), Oxidative phosphorylation, halide dependent and halide independent effects of triorganotin and triorganolead compounds on mitochondrial functions. *Biochem. J.* **168**, 353–364.
- Ali S. T., William S. Jr., Glen D. N. and Philip T. G. (1986), Cholinergic and dopaminergic alterations in the mouse central nervous system following acute trimethyl tin exposure. *Acta Pharmacol. Toxicol.* **59**, 179–188.
- Carroll N. V., Longley R. W. and Roe J. H. (1956), Glycogen determination in liver and muscle by use of anthrone reagent. *J. Biol. Chem.* **220**, 583–593.
- Desaiah D. and Ho I. K. (1979), Effect of acute and continuous morphine administration on catecholamine-sensitive adenosine triphosphatase in mouse brain. *J. Pharmacol. Exp. Ther.* **208**, 80–85.
- Fingerman M. (1995), Endocrine mechanisms in crayfish, with emphasis on reproduction and neurotransmitter regulation of hormone release. *Am. Zool.* **35**, 68–78.
- Fingerman M. (1997), Crustacean endocrinology: A retrospective, prospective, and introspective analysis. *Physiol. Zool.* **70**, 257–269.
- Hodge V., Seidel S. L. and Goldberg E. D. (1979), Determination of tin IV and organotin compounds in natural waters, coastal sediments and macroalgae by atomic absorption spectrometry. *Annal. Chem.* **51**, 1256–1259.
- Huberman A. and Aguilar M. B. (1986), A neurosecretory hyperglycemic hormone from the sinus gland of the Mexican crayfish *Procambarus bouvieri* (Ortmann). I. Purification and biochemical characterization of the most abundant form of the hormone. *Comp. Biochem. Physiol.* **85 B**, 197–203.
- Keller R. and Wunderer G. (1978), Purification and amino acid composition of the neurosecretory hyperglycemic hormone from the sinus gland of the shore crab *Carcinus maenas*. *Gen. Comp. Endocrinol.* **34**, 328–355.
- Keller R. (1981), Purification and amino acid composition of the hyperglycemic neurohormone from the sinus gland of *Orconectes limosus* and comparison with the hormone from *Carcinus maenas*. *J. Comp. Physiol.* **141**, 445–450.
- Keller R., Jaros P. P. and Kegel G. (1985), Crustacean hyperglycemic neuropeptides. *Am. Zool.* **25**, 207–221.
- Kleinholz L. H. (1975), Purified hormones from the crustacean eyestalk and their physiological specificity. *Nature*. **258**, 256–257.
- Kleinholz L. H. and Keller R. (1979), Endocrine regulation in Crustacea In: *Hormones and Evolution* (E. J. W. Barrington, ed.). Academic Press, New York. pp. 160–213.
- Langston W. J., Burt G. R. and Zkou M. J. (1987), Tin and organotin in water, sediment and benthic organisms of poole harbour. *Mar. Poll. Bull.* **18**, 634–639.
- Liu C. Y., Chen S. K., Kou G. M. and Kuo C. M. (1998), Identification and characterization of a hyperglycemic hormone from freshwater giant prawn, *Macrobrachium rosenbergii*. *Comp. Biochem. Physiol.* **121**, 315–321.
- Marco K. G., Brandt W. and Gade G. (1998), Elucidation of the amino acid sequence of a crustacean hyperglycemic hormone from the spiny lobster, *Jasus lalandii*. *Biochem. Biophys. Res. Commun.* **248**, 578–583.
- Margaillan A., Milano J. C. and Vernet J. C. (1985), Continuation by tin in some areas of the mediterranean coast. *J. Res. Oceanogr.* **10**, 96–99.
- Newcomb R. W. (1983), Peptides in the sinus gland of *Cardisoma carnifex*: Isolation and amino acid analysis. *J. Comp. Physiol.* **153 B**, 207–211.
- Ohira, T., Watanabe, T., Nagasawa, H. and Aida, K. (1997), Cloning and sequence analysis of a cDNA encoding a crustacean hyperglycemic hormone from the kumura prawn *Penaeus japonicus*. *Mol. Mar. Biol. Biotech.* **6**, 59–63.
- Rao K. S. P., Chetty C. S. and Desaiash D. (1987), Effects of tricyclohexylhydroxytin on the kinetics of adenosine triphosphatase system and protection by thiol reagents. *J. Biochem. Toxicol.* **2**, 125–140.
- Reddy P. S. and Ramamurthi R. (1999), Recent trends in crustacean endocrine research. *PINSA.* **65 B**, 15–32.
- Reddy P. S., Prestwich G. D. and Chang E. S. (1997), Crustacean hyperglycemic hormone gene expression in the lobster *Homarus americanus*. *XIII Int. Congress Comp. Endocrinology*. Monduzzi editore, Bologna, Italy, pp. 51–55.
- Sedlmeier D. (1985), Mode of action of crustacean hyperglycemic hormone. *Am. Zool.* **25**, 223–232.
- Seinen W., Vos J. G., Brands R. and Hoojkaas H. (1979), Lymphocytotoxicity and immunosuppression by organotin compounds: Suppression of graft-versus-host reactivity, blast transformation, E-rosetta formation by di-n-butyltin dichloride and di-n-octyltin dichloride. *Immunopharmacology.* **1**, 343–355.
- Sithigorngul W., Jaideechoey S., Saraithongkuv W., Longyaut S. and Sithigorngul P. (1999), Purification and characterization of an isoform of crustacean hyperglycemic hormone from the eyestalk of *Macrobrachium rosenbergii*. *J. Exp. Zool.* **284**, 217–224.
- Soyez D., Noel P. Y., Van Deijnen J. E., Martin M., Mortel A. and Payen G. G. (1990), Neuropeptides from the sinus gland of the lobster *Homarus americanus*: Characterization of hyperglycemic peptides. *Gen. Comp. Endocrinol.* **79**, 261–264.
- Steel R. G. D. and Torrie J. H. (1960), In: *Principles and Procedures of Statistics, a Biomedical Approach*. McGraw-Hill Book Company, New York.
- Van Herp F. and Payen G. G. (1991), Crustacean neuroendocrinology: perspectives for the control of reproduction in aquacultural systems. *Bull. Inst. Zool. Acad. Sin. Monogr.* **16**, 513–539.
- Yasuda A., Yasuda Y., Fujita T. and Naya Y. (1994), Characterization of crustacean hyperglycemic hormone from the crayfish (*Procambarus clarkii*): multiplicity of molecular forms by stereoinversion and diverse functions. *Gen. Comp. Endocrinol.* **95**, 387–393.