

## Antimicrobial Activity of Isopteropodine

Rubén García<sup>a,\*</sup>, Cesia Cayunao<sup>a</sup>, Ronny Bocic<sup>a</sup>, Nadine Backhouse<sup>a</sup>,  
Carla Delporte<sup>a</sup>, Mercedes Zaldivar<sup>b</sup>, and Silvia Erazo<sup>a</sup>

<sup>a</sup> Department of Pharmacological and Toxicological Chemistry, School of Chemical and Pharmaceutical Sciences, University of Chile, P. O. Box 233, Santiago 1, Chile.  
Fax: 56-2-2227900. E-mail: rgarciam@uchile.cl

<sup>b</sup> Department of Biochemistry and Molecular Biology, School of Chemical and Pharmaceutical Sciences, University of Chile, Santiago, Chile

\* Author for correspondence and reprint requests

Z. Naturforsch. **60c**, 385–388 (2005); received October 18/November 29, 2004

Bioassay-directed fractionation for the determination of antimicrobial activity of *Uncaria tomentosa*, has led to the isolation of isopteropodine (0.3%), a known *Uncaria* pentacyclic oxindol alkaloid that exhibited antibacterial activity against Gram positive bacteria.

*Key words:* *Uncaria tomentosa*, Isopteropodine, Antibacterial Activity

### Introduction

*Uncaria tomentosa* (Willd. ex Roemer and Schultes DC.), commonly known as “cat’s claw”, is a large climbing shrub, belonging to the Rubiaceae family. Cat’s claw is indigenous to the Amazon rainforest and other tropical areas of South and Central America, including Peru, Colombia, Ecuador, Guyana, Trinidad, Venezuela, Suriname, Costa Rica, Guatemala, and Panama (Keplinger *et al.*, 1999; Lock de Ugaz, 1995).

The bark of cat’s claw is used in either folk medicine or in procuring phytotherapeutic drugs. This species contain several active compounds which were tested widely for possible medicinal value (rheumatism, tumors, anti-inflammatory effect and breast cancer) (Williams, 2001; Riva *et al.*, 2001; Aguilar *et al.*, 2002; Gattuso *et al.*, 2004).

Chemical studies on *Uncaria tomentosa* revealed the presence of diverse compounds as quinovic glycosides, polyoxygenated triterpenoids, catechins, sterols and alkaloids, with diverse pharmacological actions (Aquino *et al.*, 1989; Wagner *et al.*, 1985; Keplinger *et al.*, 1999; Kang *et al.*, 2002; Sandoval *et al.*, 2002; Mur *et al.*, 2002; Montoro *et al.*, 2004).

However, of all the extensive biological activities investigated in this plant, specially for pentacyclic oxindol alkaloids (Kang *et al.*, 2002; Lee *et al.*, 1999; Falkiewicz and Lukasiak, 2001) – only the quinovic glycosides were reported to have antimicrobial (antiviral) (Aquino *et al.*, 1989; Williams, 2001) activity – reputed in folk medicine for *Uncaria tomentosa*. These instances reinforced our

interest for evaluating the antimicrobial activity of “cat’s claw” bark against bacteria and fungi.

### Materials and Methods

#### *General experimental procedures*

The solvent used for NMR studies was CDCl<sub>3</sub>. The measurements of the NMR spectra were carried out on a Bruker AMX-300 [<sup>1</sup>H NMR (300 MHz), <sup>13</sup>C NMR (75 MHz)] spectrometer. Column chromatography (CC) was carried out using silica gel 60 G (Merck, 7734). TLC was performed on silica gel GF 254 plates (Merck, 5554); the spots were detected by UV light (254, 366 nm), Liebermann Burchard test and/or *p*-anisaldehyde and Dragendorff reagents.

GC-MS analyses were made on a Fisons MB 800 mass spectrometer (MS) coupled with a Hewlett-Packard mod. 5890 series II gas chromatograph (GC), equipped with a 25 m × 0.2 mm i.d. HP U-2 column, with 0.25 μm film thickness. The initial oven temperature was held at 40 °C for 6 min, it was then increased at 7 °/min up to 200 °C, at 110 KPa with helium as carrier gas.

HPLC was performed on a Shimadzu LC-10 AD, with a Shimadzu SPD-10 AV UV-VIS detector. Column: C<sub>18</sub> (Phenomenex); solvent system: acetonitrile/buffer K<sub>2</sub>HPO<sub>4</sub> (pH 7) 50:50 with a flow of 0.75 ml/min to 27 °C.

#### *Plant material*

*U. tomentosa* bark was kindly provided by Hochstetter Laboratory (collected from Peruvian

rainforest), and a voucher is kept at the botany faculty laboratory.

#### Extraction and isolation

Ground dried bark (720 g) was sequentially extracted at room temperature with *n*-hexane, dichloromethane and methanol, yielding after removal of the solvents *in vacuo*, 2.10 g (0.3%), 2.86 g (0.4%) and 82.5 g (11.5%), respectively.

Dichloromethane and methanol extracts were active in the preliminary bio-assay, showing the presence of a single bioactive alkaloid substance. The extracts were separately worked up dissolving the dried extracts in acetic acid (2 N), filtering, basifying with concentrated ammonia solution, and extracting with chloroform (5 × 25 ml). The chloroform extracts were joined and subjected to CC over silica gel, eluting an unpurified alkaloid with dichloromethane/ethyl acetate 9.5:0.5 to dichloromethane/ethyl acetate 8:2, in fractions 49–72, respectively. Preparative TLC on silica gel led to the isolation of the purified alkaloid compound. The purity of this substance was evaluated by gas chromatography-mass spectroscopy, showing a single peak at the retention time (Rt) of 21.5 min. Crystallization from MeOH gave colorless needles.

*Isopteropodine*: M.p. 207–209 °C. –  $[\alpha]_D^{24} - 85^\circ$  ( $c = 0.764$ , CHCl<sub>3</sub>). – UV(MeOH):  $\lambda_{\max} = 208, 243, 283$  nm (sh). – MS:  $m/z$  (% rel. int.) = 368 (4.5) [M<sup>+</sup>], 223 (13), 208 (11), 180 (13), 146 (55), 130 (61), 117 (58), 103 (25), 77 (72), 69 (100), 55 (74); C<sub>21</sub>H<sub>2</sub>N<sub>2</sub>O<sub>4</sub>. – <sup>13</sup>C NMR (ppm):  $\delta = 181.43, 167.64, 154.98, 140.23, 133.76, 127.67, 124.53, 122.52, 109.82, 109.70, 72.14, 71.25, 56.93, 54.14, 53.51, 51.02, 37.83, 34.84, 30.17, 30.42, 18.65$ . – <sup>1</sup>H NMR (ppm):  $\delta = 8.56$  (1H, brs), 7.42 (1H, s), 7.28 (1H, d,  $J = 7.5$  Hz), 7.19 (1H, t,  $J = 7.6$  Hz), 7.02 (1H, t,  $J = 7.5$  Hz), 6.93 (1H, d,  $J = 7.6$  Hz), 4.35 (1H, m), 3.60 (3H, s), 3.29 (2H, m), 2.46 (5H, m), 2.01 (1H, m), 1.59 (2H, m), 1.41 (3H, d,  $J = 6.2$  Hz), 0.87 (1H, appq,  $J = 12.8$  Hz) (\* brs: broad singlet; appq: apparent quartet).

#### Antimicrobial assays

The antimicrobial activity of the extracts was determined against *Escherichia coli* (ATCC 8739), *Klebsiella pneumoniae* (isolated from a patient), *Salmonella aviatum* (ATCC 2228), *Pseudomonas aeruginosa* (ATCC 14207), *Staphylococcus aureus* (ATCC 6538P), *Micrococcus flavus* (ATCC 10290)

and *Bacillus subtilis* (ATCC 6633), *Candida albicans* and *Saccharomyces cerevisiae*.

Dilutions of 100 and 200 µg/ml of DMSO extract's solution were added to a fixed volume of Plate Count Agar (PCA) and Tryptic Soy Broth (TSB). They were then superficially inoculated with a single line of an overnight culture of the different microorganisms and incubated at 37 °C for 24 h for bacteria and 28 °C for fungi. Results were recorded as growth or growth inhibition at each extract concentration.

The active extracts were submitted to a bioautography agar overlay bioassay in order to determine the active compounds (Rahalison *et al.*, 1995).

The turbidimetric method (Balows *et al.*, 1991) was used for determining MIC values of the isolated substance against *S. aureus* and *B. subtilis*. It consisted in preparing serial dilutions of the compound (Table I) in TSB culture medium. Each tube (carrying the same volume) was inoculated with a fixed volume of the culture medium to obtain 2 ml of final volume. Each dilution of the sample in study was assayed in triplicate.

#### Results and Discussion

*U. tomentosa* active extracts (dichloromethane and methanol extracts) showed antimicrobial activity only against *M. flavus* and *B. subtilis* and was inactive against *C. albicans* and *S. cerevisiae*.

The methanol extract presented the major antibacterial activity at concentrations of 100 µg/ml and 200 µg/ml. A bioautographic agar overlay in TLC of the extracts and the isolated alkaloid was carried out on a silica gel 60 F<sub>254</sub> plate developed with dichloromethane/ethyl acetate 1:1. The bioautograms were sprayed with an aqueous solution of thiazolyl blue (MTT). A positive antimicrobial reaction was observed as a clear inhibition zone against a purple background (Rahalison *et al.*, 1995).

The isolated alkaloid was identified unambiguously as isopteropodine (Uncarine E), by its mass fragmentation, <sup>1</sup>H NMR, <sup>13</sup>C NMR and literature data (Chan *et al.*, 1966; Shamma and Foley, 1967; Wagner *et al.*, 1985; Seki *et al.*, 1993).

The isopteropodine bark quantification (0.3%) was assayed by HPLC using the isolated alkaloid as internal reference (Rt: 12.1; Ganzera *et al.*, 2001).

Isopteropodine was inactive, in the same way as *Uncaria* extracts, against *C. albicans* and *S. cerevisiae*, and was active only against the Gram positive bacteria *S. aureus* and *B. subtilis*, with a MIC value

Table I. Isopteropodine MIC value determinations.

Bacteria	Isopteropodine concentration [ $\mu\text{M}$ ]												
	353	380	408	435	462	489	516	543	571	598	625	652	679
<i>S. aureus</i>	+	+	-	-	-	-	-	-	-	-	-	-	-
<i>B. subtilis</i>	+	+	+	+	+	+	+	+	+	+	+	+	-

+, Bacterial growth.

-, No bacterial growth.

of 150  $\mu\text{g/ml}$  (408  $\mu\text{M}$ ) and 250  $\mu\text{g/ml}$  (679  $\mu\text{M}$ ), respectively (Table I) [ampicillin: MIC 5  $\mu\text{g/ml}$  (14  $\mu\text{M}$ ) for *S. aureus* and 10  $\mu\text{g/ml}$  (29  $\mu\text{M}$ ) for *B. subtilis*].

In conclusion, these results support scientifically the use of this species in popular medicine as antimicrobial, despite its weaker activity compared with the standard antibiotic.

#### Acknowledgements

This research has been performed under the auspices of “Programa de Ciencias y Tecnología para el Desarrollo, CYTED, Subprograma X” and Laboratorio Hochstetter S. A.

- Aguilar J., Rojas P., Marcelo A., Plaza A., Bauer R., Reiningger E., Klaas C., and Merfort I. (2002), Anti-inflammatory activity of two different extracts of *Uncaria tomentosa* (Rubiaceae). *J. Ethnopharm.* **81**, 271–276.
- Aquino R., De Simone F., Pizza C., Conti C., and Stein M. L. (1989), Plant metabolites, structure and *in vitro* antiviral activity of quinovic glycosides from *Uncaria tomentosa* and *Guettarda platypoda*. *J. Nat. Prod.* **52**, 679–685.
- Balows A., Hausler W., Hermann K., and Isenberg H. (1991), *Manual of Clinical Microbiology*, 5<sup>th</sup> ed. (Shadomy H. J., ed.). Amer. Soc. Microbiol. Washington DC., pp. 1105–1112.
- Chan K., Morsingh F., and Yeoh G. (1966), Alkaloids of *Uncaria pteropoda*. Isolation and structures of pteropodine and isopteropodine. *J. Chem. Soc. Perkin Trans.* **24**, 2245–2249.
- Falkiewicz B. and Lukasiak J. (2001), Vilcacora [*Uncaria tomentosa* (Willd.) DC. and *Uncaria guianensis* (Aubl.) Gmel]. A review of published scientific literature. *Case Rep. Clin. Pract. Rev.* **2**, 305–316.
- Ganzera M., Muhammad I., Khan R., and Khan I. (2001), Improved method for the determination of oxindole alkaloids in *Uncaria tomentosa* by high performance liquid chromatography. *Planta Med.* **67**, 447–450.
- Gattuso M., Di Sapio O., Gattuso S., and Pereyra E. (2004), Morphoanatomical studies of *Uncaria tomentosa* and *Uncaria guianensis* bark and leaves. *Phytochem.* **11**, 213–223.
- Kang T., Matsumoto K., Tohda M., Murakami Y., Takayama H., Kitajima M., Aimi N., and Watanabe H. (2002), Pteropodine and isopteropodine positively modulate the function of rat muscarinic M (1) and 5-HT (2) receptors expressed in *Xenopus oocyte*. *Eur. J. Pharmacol.* **24**, 39–45.
- Keplinger K., Laus G., Wurm M., Dierich M., and Teppner H. (1999), *Uncaria tomentosa* (Willd.) DC. Ethnomedicinal use and new pharmacological, toxicological and botanical results. *J. Ethnopharm.* **64**, 23–34.
- Lee K., Zhou B., Kingston D., Vaisberg A., and Hammond G. (1999), Bioactive indole alkaloids from the bark of *Uncaria guianensis*. *Planta Med.* **65**, 759–760.
- Lock de Ugaz O. (1995), Revisión del Género *Uncaria*. *Uncaria tomentosa* y *Uncaria guianensis*: Las “Uñas de Gato”. *Revista de Química* **9**, 49–61.
- Montoro P., Carbone V., Quiroz J., De Simone F., and Pizza C. (2004), Identification and quantification of components in extracts of *Uncaria tomentosa* by HPLC-ES/MS. *Phytochem. Anal.* **15**, 55–64.
- Mur E., Hartig F., Eibl G., and Schirmer M. (2002), Randomized double blind trial of an extract from the pentacyclic alkaloid-chemotype of *Uncaria tomentosa* for the treatment of rheumatoid arthritis. *J. Rheumatol.* **29**, 678–681.
- Rahalison L., Hamburger M., Hostettmann K., and Monod F. (1995), A bioautographic agar overlay method for detection of antifungal compounds from higher plant. *Phytochem. Anal.* **2**, 129–203.

- Riva L., Coradini D., Di Fronzo G., De Feo V., De Tommasi N., De Simone F., and Pizza C. (2001), The anti-proliferative effects of *Uncaria tomentosa* extracts and fractions on the growth of breast cancer cell line. *Anticancer Res.* **21**, 2457–2461.
- Sandoval M., Okuhama N., Zhang X., Condezo L., Lao J., Angeles F., Musah R., Bobrowski P., and Miller M. (2002), Anti-inflammatory and antioxidant activities of cat's claw (*Uncaria tomentosa* and *Uncaria guianensis*) are independent of their alkaloid content. *Phytomed.* **9**, 325–337.
- Seki H., Takayama H., Aimi N., Skai S., and Ponglux D. (1993), A nuclear magnetic resonance study on the eleven stereoisomers of heteroyohimbine-type oxindole alkaloids. *Chem. Pharm. Bull.* **41**, 2077–2086.
- Shamma M. and Foley K. (1967), Mass spectrometry and the stereochemistry of the pentacyclic oxindole alkaloids. *J. Org. Chem.* **32**, 4141–4143.
- Wagner H., Kreutzkamp B., and Jurcic K. (1985), The alkaloids of *Uncaria tomentosa* and their phagocytosis-stimulating action. *Planta Med.* **51**, 419–423.
- Williams J. (2001), Review of antiviral and immunomodulating properties of plants of the Peruvian rainforest with a particular emphasis on Uña de Gato and Sangre de Grado. *Altern. Med. Rev.* **6**, 567–579.