Structure Proposal for a New Pyoverdinin from *Pseudomonas* sp. PS 6.10

Herbert Budzikiewicz^{a,*}, Mathias Schäfer^a, Diana Uría Fernández^{a,b},
and Jean-Marie Meyer^c

^a Institut für Organische Chemie der Universität zu Köln, Greinstr. 4, D-50939 Köln, Germany. Fax +49-221-470-5057. E-mail: aco88@uni-koeln.de

^b Present address: Waters MS Technology Centre, Manchester, UK

^c Laboratoire de Microbiologie et de Génétique, UPRES-A 7010 du CNRS, Université Louis Pasteur, 28 rue Goethe, F-67000 Strasbourg, France

* Author for correspondence and reprint requests

Z. Naturforsch. **61c**, 815–820 (2006); received August 22, 2006

From *Pseudomonas* sp. PS 6.10, when grown in a casamino acid medium, a pyoverdinin was isolated whose primary structure could be deduced from its mass spectrometric fragmentation pattern and amino acid analysis. It belongs to the smallest representatives of this group of siderophores comprising only six amino acids in its peptide chain. When grown in a succinate minimal medium the corresponding ferri-bactin considered to be the biogenetic precursor of the pyoverdinin was obtained as the major component.

Key words: *Pseudomonas*, Pyoverdinin, Siderophore

Introduction

Pseudomonas sp. PS 6.10 is a member of the fluorescent species in the rRNA homology group I of the family Pseudomonadaceae which produces siderophores (“pyoverdins”) with high Fe³⁺ complexing constants. The strain was isolated from the cultivated mushroom *Agaricus bisporus* as a member of the “*Pseudomonas reactans*” group since it was non-pathogenic and developed a positive white line test against *Pseudomonas tolaasii* (Munsch *et al.*, 2000). Furthermore, siderotyping analysis of the strain showed that its pyoverdinin-iso-electrofocusing pattern was original and that the pyoverdinin was highly strain-specific in its iron transport capacity.

Today about 50 complete or fairly complete pyoverdinin structures have been elucidated (Budzikiewicz, 2004; Fuchs *et al.*, 2001), and from preliminary siderotyping studies it appears that many more are to be expected. Pyoverdins consist of three distinct structural parts, viz. a dihydroxy-quinoline chromophore responsible for their fluorescence, a peptide chain comprising 6 to 12 amino acids bound to the chromophore carboxy group,

and a small dicarboxylic acid (or its monoamide) connected amidically to its NH₂ group (cf. **1**). The peptide chain has a twofold function. It provides two of the ligand sites for Fe³⁺, and it is responsible for the recognition of the ferri-pyoverdins by specific receptors located at the surface of the producing cell (Meyer *et al.*, 2002). The variability of the peptide chain is closely connected with the second function: It safeguards that a given ferri-pyoverdinin is available only to the producing strain because of the usually highly specific interaction between the ferri-pyoverdinin and its receptor outer membrane protein (Hohnadel and Meyer, 1988).

In the bacterial culture medium occasionally together with pyoverdins compounds are encountered which are considered to be their biogenetic precursors. They have the same peptide chain as the corresponding pyoverdins, but differ in the *N*-terminal part. Thus in ferri-bactins the pyoverdinin chromophore is replaced by a condensation product of D-Tyr and L-2,4-diaminobutanoic acid (Böckmann *et al.*, 1997; Hohlneicher *et al.*, 1992, 2001). The strain presently studied, *Pseudomonas* sp. PS 6.10, when grown in a casamino acid medium, produced large amounts of pyoverdinin, but in succinate minimal medium the corresponding ferri-bactin was produced as the major component.

Over the years much knowledge has been accumulated on the mass spectrometric fragmentation behavior of pyoverdins after electrospray ioniza-

Abbreviations: Common amino acids, 3-letter code; Dab, 2,4-diaminobutanoic acid; AcOHOrn, *N*⁴-acetyl-*N*⁴-hydroxy Orn; OHAsp, *threo*-β-hydroxy Asp; MS, mass spectrometry; ESI, electrospray ionization; CA, collision activation.

tion (ESI) and collisional activation (CA) (Budzikiewicz, 2004; Fuchs and Budzikiewicz, 2001a) which in many cases allows the determination of the amino acid sequence of the peptide chain, even from crude bacterial culture extracts without previous isolation of the siderophores. We wish to demonstrate this for a pyoverdins and the corresponding ferri-bactin obtained from the *Pseudomonas* sp. PS 6.10 strain.

Materials and Methods

Mass spectrometry was done using a Finnigan-MAT 900 ST with an ESI source, solvent CH₃OH/H₂O 1:1. Mass selected fragmentation by CA was conducted either in the quadrupole region in front of or in the ion trap. GC-MS analysis of the trifluoroacetyl isopropyl (TAP) derivatives of the amino acids obtained by hydrolysis was performed with an Agilent Technologies GC 6890 apparatus. Chiral amino acid analysis of the TAP derivatives was effected on a Chirasil-L-Val column (Chrompack, Frankfurt) using a Varian 3400CX GC apparatus.

The bacterial strain was grown either in a cas-amino acid or in a succinate minimal medium (Meyer *et al.*, 1997). For the work-up of the culture after addition of ferric citrate and the isolation of the ferri-pyoverdins by chromatography on XAD-4 resin see Georgias *et al.* (1999). Siderotyping controls including isoelectrophoresis analysis of the pyoverdins and pyoverdin-mediated iron uptake studies were done as described previously (Munsch *et al.*, 2000).

Results

Pyoverdins

After total hydrolysis and TAP derivatization by GC/MS analysis and GC analysis using a chiral column (Dallakian *et al.*, 1999) the presence of D-Ala, OHAsp, L-Dab, L-Lys, L-Orn and/or OHOrn was established.

The molecular mass of the main component was determined by ESI-MS as 1090 u. As will be shown below it corresponds to a species with a succinamide side chain (Fig. 1, **1a**). For structure elucidation the precursor ions $[M+H]^+$ or $[M+2H]^{2+}$ of **1a** were excited by CA both in the quadrupole region and in the ion trap of the mass spectrometer. In the quadrupole several collisions may occur and ions resulting from consecutive fragmentation processes will be observed. In the ion trap a single ion species is selected and only single-step fragmentation processes are possible, but the relatively long lived ions in the trap may

Table I. Amino acid sequence characteristic fragment ions found in the MS-CA spectra of $[M+H]^+$ and/or of $[M+2H]^{2+}$ of **1a** (see text).

<i>n</i>	Amino acid	A	B	Y''	<i>n</i>
1	Ala	400	428		6
2	Orn		542		5
3	OHAsp		673	550	4
4	Dab		773	419	3
5	AcOHOrn		945	319	2
6	Lys			147	1

Fig. 1. Structure proposed for pyoverdins **1** (**1a**: R = NH₂; **1b**: R = OH; **1c**: R = OCH₃).

Fig. 2. Quadrupole ion CA mass spectrum (lower part) of $[M+2H]^{2+}$ of **1a**. Marked with Chr are the fragments formed by *retro*-Diels-Alder decomposition of the chromophore (m/z 303) and its further cleavage products; m/z 70–129 comprise the fragments characteristic for amino acids (see text).

partially undergo rearrangement processes. CA of $[M+H]^+$ mainly yields fragments containing the protonated chromophore, while in the CA spectra of $[M+2H]^{2+}$ also ions comprising parts of the *C*-terminus of the peptide chain (Y-ions). [The *N*-terminal fragments obtained by cleavage before, in or after peptide bond $-\text{CHR}-\text{CO}-\text{NH}-\text{CHR}'-$ are designated by A, B, and C, the *C*-terminal ones by X, Y, and Z. The number of transferred hydrogen atoms is indicated by hyphens. Thus, $-\text{CHR}\text{CO}^+$ is a B ion, $\text{H}_3\text{N}^+-\text{CHR}'-$ an Y'' ion (Roepstorff and Fohlman, 1984).] are observed, since the second proton can induce fragmentation anywhere in the chain (Budzikiewicz, 2004; Fuchs and Budzikiewicz, 2001a).

Both in the quadrupole and the ion trap CA spectra of $[M+H]^+$ (m/z 1091) loss of 1 to 4 H_2O molecules is observed. Cleavage of the various peptide bonds yields A_1 and the complete series of B ions (see Table I). All these ions show loss of H_2O , A_1 in addition loss of NH_3 and of the entire succinamide residue with back transfer of one H atom (m/z 301). $[\text{B}_3-\text{H}_2\text{O}]^+$ (m/z 655) loses $\text{HOOC}-\text{CHO}$ from OHAsp by a McLafferty rearrangement (transfer of H to the peptide car-

bonyl group) (m/z 581). B_5 (m/z 945) is accompanied by $[\text{B}_5+\text{H}_2\text{O}]^+$ (m/z 963) formed by transfer of the *C*-terminal hydroxy group (Fuchs and Budzikiewicz, 2001b).

In the quadrupole CA spectrum (Fig. 2) of $[M+2H]^{2+}$ (m/z 546) the *retro*-Diels-Alder fragmentation of the chromophore can be observed. Loss of C-2 and C-1 of the protonated chromophore together with the peptide chain leads to m/z 303 of low abundance, because this ion decomposes further by the loss of NH_3 and subsequently CH_2O (m/z 286 and 256), of $[\text{CH}_2-\text{CH}_2-\text{CONH}_2+\text{H}]$ (m/z 230) and of the entire side chain with back-transfer of a H atom (m/z 204). This fragmentation is characteristic for a pyoverdins chromophore (in contrast to isopyoverdins) (Fuchs and Budzikiewicz, 2001a). Ions at m/z 129/84, 115/70 and 101 (the expected m/z 56 is beyond the recording range of the instrument) are in agreement with the presence of Lys, Orn and Dab $[\text{H}_2\text{N}-(\text{CH}_2)_n-\text{CHNH}_2-\text{CO}^+$ and $\text{H}_2\text{N}-(\text{CH}_2)_{n-1}-\text{CH}=\text{CH}^+]$, respectively, those at m/z 114 and 86 with an N^4 -acyl- N^4 -hydroxy-ornithine residue ($\text{NHOH}-\text{CH}_2-\text{CH}_2-\text{CH}=\text{CH}-\text{CO}^+$ and subsequent loss of CO).

Most pronounced is the signal at m/z 400 representing the A_1 ion. Abundant A_1 and weak B_1 ions are typically observed with a small neutral amino acid bound to the chromophore, in the present case Ala. A_1 can lose NH_3 (m/z 383), H_2O (m/z 382), and the entire side chain (m/z 301). B_2 can be found at m/z 542. The mass difference corresponds to Orn.

In the ion trap CA spectrum of $[M+2H]^{2+}$ again the whole series of B ions can be seen as well as the loss of H_2O , $HOOC-CHO$ from m/z 655 and formation of $[B_5+H_2O]^+$ as discussed above, but in addition the Y'' ions up to Y_4'' (m/z 550) are present (see Table I).

These results allow to propose the structure **1** (Fig. 1) for the pyoverdins of *Pseudomonas* sp. PS 6.10. The conclusions derived from the mass spectrometric fragmentation pattern are confirmed by the amino acid analysis and by NMR data.

In some bacterial cultures additional components with a succinic acid side (the hydrolysis product of succinamide) and with a methyl succinate side chain (**1b** and **1c**) could be found. The latter one may be an artifact (Demange *et al.*, 1990).

Ferribactins

After total hydrolysis and TAP derivatization by GC/MS and GC analysis the presence of Ala, OHAsp, Dab, Glu, Lys, Orn and/or OHOrn and Tyr was established.

In its CA spectrum (Fig. 3) $[M+H]^+$ (m/z 1109) loses up to 4 H_2O and also the Glu side chain (loss of 129 u, followed by that of 1 and 2 H_2O). From the sequence specific ions (see Table II) B_1 is of high abundance showing the loss of H_2O and of 129 u. The same degradation is also observed for A_1 . B_3 loses H_2O and $HOOC-CHO$ (indicated by McL) as described above. The ions characteristic for Lys (m/z 129 and 84; those for Orn and Dab disappear in the general noise) and for AcOHOrn (m/z 86) can be seen in the quadrupole spectrum of $[M+2H]^{2+}$ (cf. above).

From the ions characteristic for the *N*-terminal part of ferribactins only the cleavage product $HO-C_6H_4-CH_2-CH=NH_2^+$ (m/z 136) has been mentioned previously (Fuchs and Budzikiewicz, 2001a; Fuchs *et al.*, 2001). In a recent publication (Budzikiewicz *et al.*, 2006) it is shown that the ions m/z 154, 182 (present in the quadrupole spectra of both $[M+H]^+$ and of $[M+2H]^{2+}$), 400 and 355

Fig. 3. Quadrupole ion CA mass spectrum of $[M+H]^+$ of the ferribactin with a Glu side chain. Chr designates the fragments formed by further decomposition of A_1 ; McL is the McLafferty fragment of B_3 (see text).

<i>n</i>	Amino acid	A		B		Y''		<i>n</i>
		Glu	Suca	Glu	Suca	Glu	Suca	
1	Ala	418	388		416			6
2	Orn			560	530			5
3	OHAsp			691	661			4
4	Dab			791	761	419	419	3
5	AcOHOrn					319	319	2
6	Lys					147	147	1

Table II. Amino acid sequence characteristic fragment ions found in the MS-CA spectra of $[M+H]^+$ and/or of $[M+2H]^{2+}$ of ferribactin with Glu *N*-terminus (left columns) and with a succinamide *N*-terminus (right columns).

Fig. 4. Fragment ion A_1 from the ferribactin **2a** (**2b** has a $\text{CO}-\text{CH}_2-\text{CH}_2-\text{CONH}_2$ terminus).

(only from $[M+2H]^{2+}$) are degradation products of the ion A_1 (**2a**, Fig. 4) which still contain the tetrahydropyrimidine ring with the Ala residue.

A second compound exhibits a $[M+H]^+$ ion at m/z 1079 in accordance with a succinamide instead of Glu *N*-terminus. Accordingly no losses of 129 u are observed. A_1 and the B ions and their further degradation products (see above) occur at masses 30 u lower than those observed for the ferribactin with a Glu substituent (Table II). A_1 , B_1 and B_2 show in addition loss of NH_3 and A_1 and B_1 loss of the succinamide side chain (-99 u). The ions 84, 129 and 86 characteristic for Lys and AcOHOrn as well as m/z 136 and 154 (see above) are present in the CA spectrum of $[M+2H]^{2+}$.

Discussion

The type of siderophores (pyoverdinin and ferribactin), the nature of the chromophore side chain, and the sequence of the amino acids in the peptide portion could be deduced from the CA spectra of their $[M+H]^+$ and $[M+2H]^{2+}$ ions. This is certainly not possible for every pyoverdinin, but partial information at least can be obtained in most cases. Subsequent confirmation was obtained by identification of the amino acids after total hydrolysis (which was also necessary for determining their chiralities).

The siderophores of *Pseudomonas* sp. PS 6.10 are remarkable in several ways. A peptide chain with six amino acids is so far the lower limit observed for pyoverdins. Pyoverdins with a free *C*-terminal amino acid have been encountered, but in many cases it could be shown that they are the hydrolysis products of cyclodepsipeptidic precursors where the *C*-terminal carboxy group forms an ester bond with an in-chain hydroxy-amino acid (Ser or Thr). This safeguards the pyoverdinin from an attack by enzymes able to degrade peptides starting from the *C*-terminus (Amann *et al.*, 2000; Budzikiewicz, 2004). In the present case OHAsp is the only hydroxy-amino acid in the peptide chain, but it is one of the ligand sites for Fe^{3+} and thus can not be used for ester formation. Thus the *C*-terminal carboxy group must be free.

The identification of a ferribactin with a succinamide *N*-terminus is remarkable also. With only one recent exception (see below) only Glu [and in one case derivatives of Glu (Uría Fernández *et al.*, 2003)] was found which is considered to be the starting compound for the other pyoverdinin side chains derived from the citric acid cycle. Succinamide as a side chain has been encountered at the beginning of the bacterial production of pyoverdins and in their immediate biogenetic precursors, the dihydropyoverdins (Budzikiewicz, 2004). In the meantime a second example was encountered. From the culture of a pyoverdinin-negative transposon mutant of *Pseudomonas fluorescens* ATCC 17400 ferribactins with a Glu and with a succinamide *N*-terminus could be isolated (Budzikiewicz *et al.*, 2006). It seems that a side chain modification of ferribactins can occur when a transformation into pyoverdinin for some reason is hampered.

So far it was not quite obvious under which experimental circumstances precursors of the respective pyoverdins could be found in culture medium (Budzikiewicz, 2004; cf. also Uría Fernández *et al.*,

2003). This is the first case where different media lead to the production of variable amounts of pyoverdins and ferribactins, respectively.

Acknowledgement

Patricia Munsch is acknowledged for the gift of the *Pseudomonas* sp. PS 6.10 strain, Christel Gruffaz and Annie Hoeft for technical assistance and GC analyses.

- Amann C., Taraz K., Budzikiewicz H., and Meyer J.-M. (2000), The siderophores of *Pseudomonas fluorescens* 18.1 and the importance of cyclopeptidic substructures for the recognition at the cell surface. *Z. Naturforsch.* **55c**, 671–680.
- Böckmann M., Taraz K., and Budzikiewicz H. (1997), Biogenesis of the pyoverdins chromophore. *Z. Naturforsch.* **52c**, 319–324.
- Budzikiewicz H. (2004), Siderophores of Pseudomonadaceae *sensu stricto* (fluorescent and non-fluorescent *Pseudomonas* spp.). *Progr. Chem. Org. Nat. Prod.* **87**, 81–237.
- Budzikiewicz H., Schäfer M., Uría Fernández D., Matthijs S., and Cornelis P. (2006), Characterization of the chromophores of pyoverdins and related siderophores by electrospray tandem mass spectrometry. *BioMetals* (in press).
- Dallakian P., Voss J., and Budzikiewicz H. (1999), Assignment of the absolute configuration of the amino acids of pyoverdins by GC/MS. *Chirality* **11**, 381–386.
- Demange P., Wendenbaum S., Linget C., Mertz C., Cung M. T., Dell A., and Abdallah M. A. (1990), Bacterial siderophores: structure and NMR assignment of pyoverdins Pa, siderophores of *Pseudomonas aeruginosa* ATCC 15692. *Biol. Metals* **3**, 55–170.
- Fuchs R. and Budzikiewicz H. (2001a), Structural studies of pyoverdins by mass spectrometry. *Curr. Org. Chem.* **5**, 265–288.
- Fuchs R. and Budzikiewicz H. (2001b), Rearrangement reactions in the electrospray ionization mass spectra of pyoverdins. *Int. J. Mass Spectrom.* **210/211**, 603–612.
- Fuchs R., Schäfer M., Geoffroy V., and Meyer J.-M. (2001), Siderotyping – a powerful tool for the characterization of pyoverdins. *Curr. Top. Med. Chem.* **1**, 31–57.
- Georgias H., Taraz K., Budzikiewicz H., Geoffroy V., and Meyer J.-M. (1999), The structure of the pyoverdins from *Pseudomonas fluorescens* 1. 3. Structural and biological relationships of pyoverdins from different strains. *Z. Naturforsch.* **54c**, 301–308.
- Hohlneicher U., Hartmann R., Taraz K., and Budzikiewicz H. (1992), Struktur von Ferribactin aus *Pseudomonas fluorescens* ATCC 13525. *Z. Naturforsch.* **47b**, 1633–1638.
- Hohlneicher U., Schäfer M., Fuchs R., and Budzikiewicz H. (2001), Ferribactins as the biosynthetic precursors of the *Pseudomonas* siderophores pyoverdins. *Z. Naturforsch.* **56c**, 308–310.
- Hohnadel G. and Meyer J.-M. (1988), Specificity of pyoverdine-mediated iron uptake among fluorescent *Pseudomonas* strains. *J. Bacteriol.* **170**, 4865–4873.
- Meyer J.-M., Stintzi A., De Vos D., Cornelis P., Tappe R., Taraz K., and Budzikiewicz H. (1997), Use of siderophores to type pseudomonads: the three *Pseudomonas aeruginosa* pyoverdins systems. *Microbiology* **143**, 35–43.
- Meyer J.-M., Geoffroy V. A., Bayse C., Cornelis P., Barelmann I., Taraz K., and Budzikiewicz H. (2002), Siderophore-mediated iron uptake in fluorescent *Pseudomonas*: characterization of the pyoverdine-receptor binding site of three cross-reacting pyoverdins. *Arch. Biochem. Biophys.* **397**, 179–183.
- Munsch P., Geoffroy V., Alatosava T., and Meyer J.-M. (2000), Application of siderotyping for characterization of *Pseudomonas tolaasii* and '*Pseudomonas reactans*' isolates associated with brown blotch disease of cultivated mushrooms. *Appl. Environ. Microbiol.* **66**, 4834–4841.
- Roepstorff P. and Fohlman J. (1984), Proposal of a common nomenclature for sequence ions in mass spectra of peptides. *Biomed. Mass Spectrom.* **11**, 601.
- Uría Fernández D., Fuchs R., Schäfer M., Budzikiewicz H., Meyer J.-M. (2003), The pyoverdins of *Pseudomonas fluorescens* G173, a novel structural type accompanied by unexpected natural derivatives of the corresponding ferribactin. *Z. Naturforsch.* **58c**, 1–10.