Liquid Chromatography-Mass Spectroscopy and Liquid Chromatography-Ultraviolet/Visible Photodiode Array Analysis of Selected *Colchicum* Species

Ahmad A. Gharaibeh^{a,*}, Ala'a Al-Serini^a, Rana M. Qasaymeh^b,
Amani S. Ma'aya'h^b, Khaled Tawaha^c, Tamam El-Elimat^{b,d}, and Feras Q. Alali^{b,*}

^a Department of Applied Chemistry, Faculty of Science and Arts,
Jordan University of Science and Technology, P. O. Box 3030, Irbid 22110, Jordan.
E-mail: agharaib@just.edu.jo

^b Department of Medicinal Chemistry and Pharmacognosy, Faculty of Pharmacy,
Jordan University of Science and Technology, P. O. Box 3030, Irbid 22110, Jordan.
E-mail: falali@just.edu.jo

^c Department of Pharmaceutical Sciences, Faculty of Pharmacy, The University of Jordan,
Amman 11942, Jordan

^d Current address: Department of Chemistry and Biochemistry,
University of North Carolina at Greensboro, Greensboro, NC 27402, USA

* Authors for correspondence and reprint requests

Z. Naturforsch. **67c**, 451–460 (2012); received September 21, 2011/July 22, 2012

An in-house strategy to dereplicate colchicinoid alkaloids was recently developed by our team. It aimed at quickly identifying *Colchicum* constituents using LC-MS (liquid chromatography-mass spectroscopy) and LC-UV/Vis PDA (liquid chromatography-ultraviolet/visible photodiode array) techniques. In this project, our goal was to validate the developed method through analysing the alkaloid-rich fractions of three *Colchicum* species that had been previously studied phytochemically using the traditional bioactivity-guided fractionation methodology. The analysed species were *Colchicum tauri* Siehe ex Stefanoff, *Colchicum stevenii* Kunth, and *Colchicum tunicatum* Feinbr., all belonging to the family Colchicaceae. In addition to identifying the compounds previously isolated and characterized by the traditional methodology, the new strategy succeeded in tentatively identifying a set of known compounds, but new to the species.

Key words: Dereplication, Colchicinoids, LC-APCI-MS

Introduction

Most natural products drug discovery programs rely on a traditional bioactivity-guided fractionation approach. Although this approach succeeds in isolating most of the well-known drugs of natural origin, it is tedious, costly, time-consuming, and may end up in isolating previously known compounds (Ghisalberti, 1993; Kingston, 1996). In the field of natural products, interest has been growing in the last twenty years in developing fast and efficient strategies to enable the discrimination between previously isolated known compounds and new compounds at the level of the crude extract. This strategy which is termed dereplication is very important for the efficient use of human and financial resources. By adopting a suitable dereplication method, tedious and time-consuming isolation of previously known compounds can be avoided, and hence efforts are focused on the

targeted isolation of compounds showing novel or uncommon spectroscopic features (Hostettmann *et al.*, 2001; Wolfender *et al.*, 2003). In a pioneering work, Corley and Durley (1994) developed strategies to search through scientific and technical network files using molecular weight, carbon count, structure fragments, and taxonomy for rapid identification of known compounds.

Hyphenated techniques, which can be defined as the coupling of high performance liquid chromatography (HPLC) separation technologies with on-line spectroscopic/spectrometric detection technologies, are the most commonly used techniques in dereplication strategies. Of these, liquid chromatography-mass spectroscopy (LC-MS) has a wide range of applications in the development of dereplication tools for the tentative identification of plant and microbial secondary metabolites (Ackermann *et al.*, 1996; Alali and El-Alali, 2005;

Carmona *et al.*, 2006; Jin *et al.*, 2006; Liao *et al.*, 2007; Liu *et al.*, 2007; Petsalo *et al.*, 2006; Sarker and Nahar, 2012).

A dereplication strategy was recently introduced by our research group for the analysis of colchicinoid alkaloids using LC-MS and liquid chromatography-ultraviolet/visible photodiode array (LC-UV/Vis PDA) techniques (Alali *et al.*, 2008a). Briefly, this strategy used a three-step approach. First, LC-MS was used to acquire the molecular ion and, hence, the molecular weight of the compound. These data were searched across natural product databases, particularly the Dictionary of Natural Products (Chapman & Hall, London, UK; Version 15:2) and against a self-compiled library of colchicinoid alkaloids. Next, LC-UV/Vis PDA analysis was used to acquire the UV/Vis spectrum of a compound that serves as a fingerprint grouping each compound into one of the different structural classes that are typical for colchicine analogues. Finally, the mass fragmentation data and the chromatographic retention times (and hence, relative polarity) were used to identify the structures of the compounds. The developed method was used to analyse *Colchicum brachyphyllum* Boiss. & Haussk. ex Boiss. (Colchicaceae), a *Colchicum* species that has been previously studied phytochemically in our laboratory (Alali *et al.*, 2005, 2008b). The interesting results obtained by the dereplication method encouraged us to investigate three *Colchicum* species, previously studied with the bioactivity-guided fractionation approach, applying the above described method to further validate our method and to detect the presence of any interesting compounds that were not separable at the preparative level.

The alkaloid-rich fractions of *Colchicum tauri* Siehe ex Stefanoff, *Colchicum stevenii* Knuth, and *Colchicum tunicatum* Feinbr. were analysed using our dereplication strategy (Alali *et al.*, 2008a). Using the bioactivity-guided fractionation approach, the following compounds had previously been isolated: from *C. tauri*: (-)-colchicine, 2-demethyl-(-)-colchicine, 3-demethyl-(-)-colchicine, (-)-demecolcine, *N*-methyl-(-)-demecolcine, 3-demethyl-*N*-methyl-(-)-demecolcine, β -lumicolchicine, 3-demethyl- β -lumicolchicine, and apigenin (Alali *et al.*, 2006a); from *C. stevenii*: (-)-colchicine, 2-demethyl-(-)-colchicine, β -lumicolchicine, cornigerine, isoandrocymbine, and *O*-methyl androcymbine (Al-Mahmoud *et al.*, 2006); from *C. tunicatum*: (-)-colchicine, 3-demethyl-(-)-colchicine, cornige-

rine, β -lumicolchicine, and (-)-androbiphenylene (Alali *et al.*, 2006b). In the present study, we report compounds tentatively identified using the dereplication method and compare them with those isolated by the traditional bioactivity-guided fractionation methodology.

Results and Discussion

The alkaloid-rich fractions of corms, leaves, and flowers of *C. tauri*, and corms and leaves of *C. stevenii* and *C. tunicatum*, respectively, were analysed analytically as outlined in the Experimental Section using LC-MS and LC-UV/Vis PDA techniques. Fig. 1 shows typical LC-MS total ion chromatograms (TICs) of the analysed parts of *C. tauri*, *C. stevenii*, and *C. tunicatum*. The molecular ion for each peak of each part is shown in Fig. 2. Table I summarizes the UV_{max} and mass spectral data of each peak as well. The identified compounds were given consecutive numbers corresponding to the order of their elution during reversed phase (RP)-HPLC as well as the order of the investigated species and their parts as arranged in Fig. 1.

The (+)-APCI mass spectrum of peak 1 at $t_R = 9.8$ min of the corms of *C. tauri* showed a parent molecular ion at m/z 372 for $[M+H]^+$, 28 Da less than the analogous peak in (-)-colchicine (Figs. 1 and 2), and the compound was eluted earlier (*i.e.* it is more polar) than (-)-colchicine. The UV/Vis PDA spectrum had two absorption maxima at 245 and 354 nm, implying structural similarities to (-)-colchicine. These data suggested that this compound was (-)-demecolcine (Alali *et al.*, 2006a; Freyer *et al.*, 1987). The (+)-APCI mass fragmentation spectrum of the molecular ion peak showed a peak at m/z 341 which was attributed to the ion $[MH-NH_2CH_3]^+$, which was followed by loss of CO tropolonic to m/z 313.

In a way analogous to the identification of peak 1, peaks 2, 3, 4, 5, 6, 7, 8, 10, 11, 12, 13, 14, 15, and 18 were identified as *N*-methyl-(-)-demecolcine, 3-demethyl-*N*-methyl-(-)-demecolcine, 2-demethyl-(-)-colchicine, (-)-colchicine or (-)-colchifoline, (-)-colchicine, β -lumi-(-)-colchicine, 2-demethyl-(-)-demecolcine or 3-demethyl-(-)-demecolcine, 3-demethyl-(-)-colchicine, apigenin, (-)-isoandrocymbine, crociflorinone, *O*-methyl-(-)-androcymbine, (-)-cornigerine, and (-)-colchicine, respectively (Figs. 1 and 2 and Table I) (Alali *et al.*, 2005, 2006a; Alali and El-Alali, 2005;

A) *Colchicum tauri*: **B) *Colchicum stevenii*** **C) *Colchicum tunicatum*:**

Fig. 1. (+)-APCI total ion chromatograms of the alkaloid-rich fractions of (A) *C. tauri*, (B) *C. stevenii*, and (C) *C. tunicatum*.

Al-Mahmoud *et al.*, 2006; Chommadov *et al.*, 1990; Freyer *et al.*, 1987; O'Neil *et al.*, 2001; Potesilova *et al.*, 1977; Rosso and Zuccaro, 1998; Roth and Rupp, 1995; Turdikulov *et al.*, 1974).

For *C. tauri*, we were unable to suggest a chemical structure for peak 9 at $t_R = 11.7$ min in the flowers and leaves that showed a parent molecular ion at m/z 370 for $[M+H]^+$. The retention

Fig. 2. (+)-APCI mass spectra of the TIC chromatographic peaks of *C. tauri*, *C. stevenii*, and *C. tunicatum* arranged based on the appearance of the corresponding chromatographic peaks in Fig. 1.

Table I. Retention times, UV_{max} , and mass spectral data of the (+)-APCI TIC chromatographic peaks.

Peak number	<i>Colchicum</i> species (plant part)	t_R [min]	UV_{max} [nm]	LC-MS and LC-MS/MS- (+)-APCI mass fragments (m/z)	Compound
1	<i>C. tauri</i> (corms)	9.8	245, 354	372 $[M+H]^+$ (base peak), 358, 341, 334, 313, 304, 298, 235, 227, 163	(-)-Demecolcine (1)
2	<i>C. tauri</i> (corms, flowers)	11.4	243, 350	386 $[M+H]^+$ (base peak), 370, 358, 355, 343, 327, 315, 283, 240, 227, 177	<i>N</i> -Methyl-(-)-demecolcine (2)

Table I continued.

Peak number	<i>Colchicum</i> species (plant part)	t_R [min]	UV_{max} [nm]	LC-MS and LC-MS/MS-(+)-APCI mass fragments (m/z)	Compound
3	<i>C. tauri</i> (corms)	11.7	244, 356	372 $[M+H]^+$ (base peak), 340, 329, 302, 299, 240, 227, 177	3-Demethyl- <i>N</i> -methyl-(-)-demecolcine (3)
4	<i>C. tauri</i> (corms, leaves) <i>C. stevenii</i> (corms, leaves)	13.2 13.4	246, 354	386 $[M+H]^+$ (base peak), 372, 327, 314, 299, 240, 227, 177	2-Demethyl-(-)-colchicine (4)
5	<i>C. tauri</i> (corms, leaves, flowers) <i>C. stevenii</i> (corms, leaves)	15.2	245, 353	416 $[M+H]^+$ (base peak), 327, 235, 227, 163	(-)-Colchicine (5)
					(-)-Colchifoline (5)
6	<i>C. tauri</i> (corms, leaves, flowers) <i>C. stevenii</i> (corms, leaves) <i>C. tunicatum</i> (corms, leaves)	15.7	247, 354	400 $[M+H]^+$ (base peak), 341, 313, 299, 282, 235, 227, 177, 163, 126	(-)-Colchicine (6)
7	<i>C. tauri</i> (corms, leaves, flowers) <i>C. stevenii</i> (corms, leaves) <i>C. tunicatum</i> (corms)	15.8	225, 273	400 $[M+H]^+$ (base peak), 368, 341, 327, 313, 235, 227, 177, 163, 127, 113	β -Lumi-(-)-colchicine (7)

Table I continued.

Peak number	<i>Colchicum</i> species (plant part)	t_R [min]	UV _{max} [nm]	LC-MS and LC-MS/MS- (+)-APCI mass fragments (m/z)	Compound
8	<i>C. tauri</i> (leaves)	9.7	242, 352	358 [M+H] ⁺ , 329, 327, 264, 235, 227, 187, 177
					3-Demethyl-(-)-demecolcine (8) or
9	<i>C. tauri</i> (leaves, flowers)	11.7	ND	370 [M+H] ⁺ (base peak), 356, 327, 297, 240, 227, 163	Unknown
10	<i>C. tunicatum</i> (corms, leaves) <i>C. tauri</i> (leaves, flowers)	13.2	243, 355	386 [M+H] ⁺ (base peak), 327, 253, 235, 227, 163
		13.3			3-Demethyl-(-)-colchicine (10)
11	<i>C. tauri</i> (flowers)	18.9	239, 259, 342	271 [M+H] ⁺ (base peak), 240, 217, 203, 189	Apigenin (11)
12	<i>C. stevenii</i> (corms, leaves)	7.8	225, 254, 288	372 [M+H] ⁺ , 327, 314, 251, 235, 227, 191, 177	(-)-Isoandrocymbine (12)

Table I continued.

Peak number	<i>Colchicum</i> species (plant part)	t_R [min]	UV _{max} [nm]	LC-MS and LC-MS/MS-(+)-APCI mass fragments (m/z)	Compound
13	<i>C. stevenii</i> (corms, leaves)	10.9	ND	328 [M+H] ⁺ , 301, 235, 227, 177	Crociflorinone (13)
14	<i>C. stevenii</i> (corms, leaves)	11.8	227, 256, 292	386 [M+H] ⁺ (base peak), 329, 315, 301, 235, 227, 177	O-Methyl-(-)-androcymbine (14)
15	<i>C. stevenii</i> (corms, leaves) <i>C. tunicatum</i> (leaves)	17.0	244, 353	384 [M+H] ⁺ (base peak), 341, 329, 325, 299, 297, 283, 235, 227, 177, 163	(-)-Cornigerine (15)
16	<i>C. stevenii</i> (leaves)	9.9	ND	370 [M+H] ⁺ (base peak), 356, 327, 313, 235, 227, 177	Unknown
17	<i>C. tunicatum</i> (corms)	11.9	ND	386 [M+H] ⁺ (base peak), 355, 327, 280, 251, 235, 227, 163	Unknown
18	<i>C. tunicatum</i> (corms, leaves)	15.4	244, 356	386 [M+H] ⁺ (base peak), 370, 327, 299, 283, 267, 226, 216, 174	(-)-Colchiceine (18)

ND, not detected.

time and UV/Vis PDA spectra were not conclusive. For the same reasons, we were unable to suggest chemical structures for peak 16 (t_R = 9.9 min) in the leaves of *C. stevenii* and peak 17 (t_R = 11.9 min) in the corms of *C. tunicatum* that showed molecular ion peaks at m/z 370 and 386, respectively.

Of the nine compounds previously isolated from *C. tauri* by bioactivity-guided fractionation (see Introduction) (Alali *et al.*, 2006a), we were able to identify all except 3-demethyl- β -lumicolchicine. Moreover, we were able to tentatively identify two compounds new for the species, namely: 3-demethyl-(-)-demecolcine [or 2-demethyl-(-)-demecolcine],

(-)-colchicine [or (-)-colchifoline], in addition to one compound, which was not identified and thus considered unknown.

All six compounds previously isolated from *C. stevenii* (Al-Mahmoud *et al.*, 2006) were identified here, and we were able to tentatively identify two new compounds, namely: (-)-colchicine [or (-)-colchifoline] and crociflorinone, in addition to one compound which was not identified and thus considered unknown.

Of the five colchicinoids previously isolated from *C. tunicatum* (Alali *et al.* 2006b), four were identified here except for (-)-androbiphenylene. On the other hand, one compound new to the species was tentatively identified namely (-)-colchicine, in addition to one compound which was not identified and thus considered unknown.

These findings, from the three studied species, clearly demonstrate the advantage of using LC-MS and LC-PDA dereplication strategies as powerful and economical tools for dereplication of natural products in crude plant extracts. Although it took around a year of hard and costly work to isolate and identify the active components from *Colchicum* species phytochemically, we were able to tentatively identify the majority of the constituents from the crude alkaloid fraction in a single linear analytical run. Moreover, the strategy was able to discriminate between previously isolated, known compounds and new compounds hitherto not described. Hence, the application of this strategy before setting out a large-scale isolation project is of great importance to avoid the tedious isolation of known constituents, and thus focus on the targeted isolation of constituents presenting novel or unusual spectroscopic features. In comparison with our already published paper (Alali *et al.*, 2008a), in the current work, we used a different LC-MS instrument (triple quadrupole mass analyser rather than ion trap) in a different laboratory and operated by a different technician, and the results thus demonstrate the robustness of the developed method.

Experimental

General

LC-MS data were collected using an Applied Biosystems (Foster City, CA, USA) triple quadrupole LC-MS instrument (API 3200) operated in the APCI positive ionization mode and an Agilent 1200 series HPLC instrument (Santa Clara,

CA, USA). The separation was achieved using a Hypersil BDS (125 mm × 4 mm; 5 µm) column (Thermo Electron, Auchtermuchty, UK). The mobile phase used was: (A) H₂O acidified with 0.1% formic acid; (B) MeOH. The flow rate was 1 mL/min in the following gradient combinations: 0–2 min, 90% (v/v) A/10% B; 2–27 min, 10% A/90% B; 27–30 min 90% A/10% B. The injection volume was 20 µL, and the total run time was 30 min. The mass detector conditions were as follows: APCI positive ionization mode; full scan mode from *m/z* 50 to 800; corona discharge voltage, NC, 5 kV; APCI temperature, 500 °C; gas flow rate, 5 L/min.

UV/Vis PDA spectra were obtained on an Agilent 1200 series HPLC instrument, equipped with an Agilent 1200 quaternary pump, a 1200 photodiode array detector in the range between 200 and 400 nm, and a 1200 series auto-sampler. Mobile phase, flow rate, analytical column, injection volume, and run times were the same as those used for LC-MS.

Formic acid (extra pure) and methanol (HPLC grade) were obtained from Scharlau Chemie S.A. (Barcelona, Spain). (-)-Colchicine standard was purchased from Fluka Chemie (Buchs, Switzerland).

Sample preparation and analysis

The alkaloid-rich fractions of the different parts of *C. tauri*, *C. stevenii*, and *C. tunicatum* came from our earlier work (Al-Mahmoud *et al.*, 2006; Alali *et al.*, 2006a, b) and were stored in glass vials wrapped with aluminum foil, sealed with Parafilm® and maintained at –2 °C. For the dereplication studies, an aliquot of the alkaloid-rich fraction of each plant part (2 mg) was dissolved in the mobile phase, filtered through a 0.45-µm Teflon filter, and then transferred to 2-mL amber HPLC vials. A 20-µL aliquot was injected into the LC-UV/Vis PDA and LC-MS systems. (-)-Colchicine standard was used for retention time matching.

Acknowledgements

The authors acknowledge the financial support from the Deanship of Scientific Research, JUST, Irbid, Jordan, and the technical help of Mr. Ashraf Al-Matalq, Pharmaceutical Research Center, JUST, Irbid, Jordan.

- Ackermann B. L., Regg B. T., Colombo L., Stella S., and Coutant J. E. (1996), Rapid analysis of antibiotic-containing mixtures from fermentation broths by using liquid chromatography-electrospray ionization-mass spectrometry and matrix-assisted laser desorption ionization-time-of-flight-mass spectrometry. *J. Am. Soc. Mass Spectrom.* **7**, 1227–1237.
- Al-Mahmoud M. S., Alali F. Q., Tawaha K., and Qasaymeh R. M. (2006), Phytochemical study and cytotoxicity evaluation of *Colchicum stevenii* Kunth (Colchicaceae): A Jordanian meadow saffron. *Nat. Prod. Res.* **20**, 153–160.
- Alali F. and El-Alali A. (2005), LC-MS analysis of corms of *Colchicum hierosolymitanum* (Colchicaceae). *Acta Pharm. Turc.* **47**, 143–150.
- Alali F. Q., El-Elimat T., Li C., Qandil A., Alkofahi A., Tawaha K., Burgess J. P., Nakanishi Y., Kroll D. J., Navarro H. A., III J. O. F., Wani M. C., and Oberlies N. H. (2005), New colchicinoids from a native Jordanian meadow saffron, *Colchicum brachyphyllum*: Isolation of the first naturally occurring dextrorotatory colchicinoid. *J. Nat. Prod.* **68**, 173–178.
- Alali F., Ma'aya'h A. S., Alkofahi A., Qandil A., Li C., Burgess J. P., Wani M. C., and Oberlies N. H. (2006a), A new colchicinoid from *Colchicum tauri*, an unexplored meadow saffron native to Jordan. *Nat. Prod. Commun.* **1**, 95–99.
- Alali F., Tawaha K., El-Elimat T., Qasaymeh R., Li C., Burgess J. P., Nakanishi Y., Kroll D. J., Wani M., and Oberlies N. H. (2006b), Phytochemical studies and cytotoxicity evaluations of *Colchicum tunicaicum* Feinbr and *Colchicum hierosolymitanum* Feinbr (Colchicaceae): Two native Jordanian meadow saffrons. *Nat. Prod. Res.* **20**, 558–566.
- Alali F., Gharaibeh A., Ghawanmeh A., Tawaha K., and Oberlies N. H. (2008a), Colchicinoids from *Colchicum crocifolium* Boiss.: A case study in dereplication strategies for (-)-colchicine and related analogs using LC-MS and LC-PDA techniques. *Phytochem. Anal.* **19**, 385–394.
- Alali F. Q., Tahboub Y. R., Al-Daraysih I. S., and El-Elimat T. (2008b), LC-MS and LC-PDA vs. phytochemical analysis of *Colchicum brachyphyllum*. *Pharmazie* **63**, 860–865.
- Carmona M., Zalacain A., Sanchez A. M., Novella J. L., and Alonso G. L. (2006), Crocetin esters, picrocrocin and its related compounds present in *Crocus sativus* stigmas and *Gardenia jasminoides* fruits. Tentative identification of seven new compounds by LC-ESI-MS. *J. Agric. Food Chem.* **54**, 973–979.
- Chommadov B., Yusupov M., and Sadykov A. (1990), Photochemical isomers of tropolone alkaloids. *Chem. Nat. Compd.* **26**, 113–128.
- Corley D. G. and Durlay R. C. (1994), Strategies for database dereplication of natural products. *J. Nat. Prod.* **57**, 1484–1490.
- Freyer A., Zarga M. A., Firdous S., Guinaudeau H., and Shamma M. (1987), Five new alkaloids from *Colchicum ritchii*. *J. Nat. Prod.* **50**, 684–689.
- Ghisalberti E. L. (1993), Detection and isolation of bioactive natural products. In: *Bioactive Natural products: Detection, Isolation and Structural Determination* (Colegate S. M. and Molyneux R. J., eds.). CRC Press, Inc., Boca Raton, FL, USA, pp. 9–57.
- Hostettmann K., Wolfender J., and Terreaux C. (2001), Modern screening techniques for plant extracts. *Pharm. Biol.* **39**, 18–32.
- Jin M., Yang Y., Su B., and Ren Q. (2006), Rapid quantification and characterization of soya saponins by high-performance liquid chromatography coupled with electrospray mass spectrometry. *J. Chromatogr. A* **1108**, 31–37.
- Kingston D. G. I. (1996), Natural products as pharmaceuticals and sources for lead structures. In: *The Practice of Medicinal Chemistry* (Meyer P., ed.). Academic Press, London, pp. 101–116.
- Liau B., Jong T., Lee M., and Chen S. (2007), LC-APCI-MS method for detection and analysis of tryptanthrin, indigo, and indirubin in Daqingye and Banlangen. *J. Pharm. Biomed. Anal.* **43**, 346–353.
- Liu A., Guo H., Ye M., Lin Y., Sun J., Xu M., and Guo D. (2007), Detection, characterization and identification of phenolic acids in Danshen using high-performance liquid chromatography with diode array detection and electrospray ionization mass spectrometry. *J. Chromatogr. A* **1161**, 170–182.
- O'Neil M. J., Smith A., and Heckelman P. E. (2001), *The Merck Index*, 13th ed. Merck Research Laboratories, Division of Merck and Co., Inc., White House Station, NJ.
- Petsalo A., Jalonen J., and Tolonen A. (2006), Identification of flavonoids of *Rhodiola rosea* by liquid chromatography-tandem mass spectrometry. *J. Chromatogr. A* **1112**, 224–231.
- Potesilova H., Dolejs L., Sedmera P., and Santavy F. (1977), Substances from plants of subfamily Wurmbeaeoideae and their derivatives 83. Constitution of 3 alkaloids (10,11-epoxycolchicine, colchicine and 3-demethyl-N-deacetyl-N-formylcolchicine) isolated from seeds of *Colchicum latifolium*. *Collect. Czech. Chem. Commun.* **42**, 1571–1580.
- Rosso A. and Zuccaro S. (1998), Determination of alkaloids from the colchicine family by reversed-phase high-performance liquid chromatography. *J. Chromatogr. A* **825**, 96–101.
- Roth L., and Rupp G. (1995), *Roth Collection of Natural Products Data*. VCH, Weinheim, Germany.
- Sarker S. D. and Nahar L. (2012), Hyphenated techniques and their applications in natural products analysis. *Methods Mol. Biol.* **864**, 301–340.
- Turdikulov Kh., Yusupov M., Aslanov Kh., and Sadykov A. (1974), Structure of crociflorinone. *Chem. Nat. Compd.* **10**, 844–845.
- Wolfender J., Ndjoko K., and Hostettmann K. (2003), Review: Liquid chromatography with ultraviolet absorbance-mass spectrometric detection and with nuclear magnetic resonance spectroscopy: a powerful combination for the on-line structural investigation of plant metabolites. *J. Chromatogr. A* **1000**, 437–455.