

Louise Ralley, Wolfgang Schuch<sup>a</sup>, Paul D. Fraser and Peter M. Bramley\*

# Genetic modification of tomato with the tobacco lycopene $\beta$ -cyclase gene produces high $\beta$ -carotene and lycopene fruit

DOI 10.1515/znc-2016-0102

Received May 18, 2016; revised July 4, 2016; accepted July 6, 2016

**Abstract:** Transgenic *Solanum lycopersicum* plants expressing an additional copy of the lycopene  $\beta$ -cyclase gene (*LCYB*) from *Nicotiana tabacum*, under the control of the *Arabidopsis* polyubiquitin promoter (*UBQ3*), have been generated. Expression of *LCYB* was increased some 10-fold in ripening fruit compared to vegetative tissues. The ripe fruit showed an orange pigmentation, due to increased levels (up to 5-fold) of  $\beta$ -carotene, with negligible changes to other carotenoids, including lycopene. Phenotypic changes in carotenoids were found in vegetative tissues, but levels of biosynthetically related isoprenoids such as tocopherols, ubiquinone and plastoquinone were barely altered. Transformants showed tolerance to the bleaching herbicide  $\beta$ -cyclase inhibitor, 2-(4-chlorophenylthio) triethylamine. The phenotype was inherited for at least three generations.

**Keywords:**  $\beta$ -carotene; genetic modification; lycopene; lycopene  $\beta$ -cyclase; *Solanum lycopersicum*.

**Dedication:** This work is dedicated to the memory of Professor Peter Böger, who kindly provided the opportunity for PDF and PMB to work in his laboratory on the mode of action of bleaching herbicides.

## 1 Introduction

Carotenoids are a major group of isoprenoids, widely distributed in nature [1]. Several of them are reported to have health benefits [2], most notably reduction in the incidence of prostate cancer from dietary lycopene [3]

<sup>a</sup>Present address: Fraunhofer Chile Research Foundation, Las Condes, Santiago, Chile.


\*Corresponding author: Peter M. Bramley, School of Biological Sciences, Royal Holloway University of London, Egham, Surrey TW20 0EX, UK, E-mail: p.bramley@rhul.ac.uk

Louise Ralley, Wolfgang Schuch and Paul D. Fraser: School of Biological Sciences, Royal Holloway University of London, Egham, Surrey TW20 0EX, UK

and alleviation of vitamin A deficiency by  $\beta$ -carotene, which is pro-vitamin A [4]. Deficiency of vitamin A causes xerophthalmia, blindness and premature death, especially in children aged 1–4 [5]. Since humans cannot synthesise carotenoids *de novo*, these health-promoting compounds must be taken in sufficient quantities in the diet. Consequently, increasing their levels in fruit and vegetables is beneficial to health. Tomato products are the most common source of dietary lycopene. Although ripe tomato fruit contains  $\beta$ -carotene, the amount is relatively low [1]. Therefore, approaches to elevate  $\beta$ -carotene levels, with no reduction in lycopene, are a goal of plant breeders. One strategy that has been employed to increase levels of health promoting carotenoids in fruits and vegetables for human and animal consumption is genetic modification [6].

Lycopene, an acyclic carotenoid and precursor of  $\beta$ -carotene, is formed from the sequential desaturation of phytoene. These reactions are catalysed by the enzymes phytoene desaturase (PDS) and  $\zeta$ -carotene desaturase (ZDS). The desaturation sequence occurs in the *cis* geometric isomer configuration and the action of a carotene isomerase (CRTISO) converts either *cis*-neurosporene or poly-*cis* lycopene to all-*trans* lycopene prior to cyclisation [7]. In green tissues, the action of light and chlorophyll are believed to overcome the necessity for CRTISO activity [8]. Cyclisation of lycopene yields  $\beta$ -carotene, via the action of a  $\beta$ -cyclase. In tomato, two lycopene cyclase are present: *LCYB* is believed to predominate in the formation of vegetative carotenoids, whilst the *CYCB* gene shows ripening specific expression and *CYCB* is thus associated with  $\beta$ -carotene production during ripening [9]. In vegetative tissues,  $\epsilon$ -carotene is formed via the action of a  $\epsilon$ -ring cyclase and  $\beta$ -ring cyclase, leading to the synthesis of lutein, via hydroxylation, which is the predominant vegetative carotenoid [10]. The complete pathway is shown in Figure 1.

Since the herbicide 2-(4-chlorophenylthio) triethylamine (CPTA) is known to inhibit  $\beta$ -carotene formation in tomato fruit [11], the transgenic lines have been grown in its presence to determine if overexpression of *LCYB* can overcome this inhibition.



**Figure 1:** Carotenoid biosynthesis in tomato.

GGPP, geranylgeranyl diphosphate; PSY-1 and -2, phytoene synthase 1 and 2; PDS, phytoene desaturase; ZDS,  $\zeta$ -carotene desaturase; CRTISO, carotene isomerase; LCYE, lycopene  $\epsilon$ -cyclase; *LCYB*, lycopene  $\beta$ -cyclase; *CYCB*, fruit specific lycopene  $\beta$ -cyclase; CRTRB1,  $\beta$ -carotene hydroxylase; CYP97A, P450 carotenoid  $\beta$ -ring hydroxylase; CYP97C, P450 carotenoid  $\epsilon$ -hydroxylase; ZEP-1, zeaxanthin epoxidase; NXS, neoxanthin synthase.

## 2 Materials and methods

### 2.1 Plant material

Tomato (*Solanum lycopersicum* cv Ailsa Craig) plants were grown in the glasshouse with supplementary lighting. Three plants per genotype were grown in a randomised manner concurrently with their respective backgrounds and fruit harvested at mature red ripe (7 days post breaker, dpb). Two or more fruits per plant were pooled to provide one replicate per plant and three per genotype.


### 2.2 Construction of the pVBLCYB vector

Digestion of pUC19 with *Sma*I released the *LCYB* cDNA as a 1.6-kb fragment, including the sequence encoding the transit peptide, which was cloned into the *Sma*I site of

the pSP vector (Promega, Southampton, UK), containing the polyubiquitin promoter (*UBQ3*; At5g03240; UniProt KB-Q1EC66) from *Arabidopsis thaliana*, the nopaline synthase terminator, the *nptII* gene for kanamycin resistance, controlled by the *AoPR1* promoter from *Asparagus officinalis*. The *Age*I fragment was sub-cloned into the pVB6 plant binary vector and designated pVBLCYB (Figure 2). The vector/insert was sequenced to confirm the manipulations. *E. coli* strain XL1-Blue (Promega) was used in these experiments. Sequences of the tobacco and tomato *LCYB* are shown in Supplementary Material, Figure 1.

### 2.3 Transformation of tomato plants

*Agrobacterium tumefaciens* LBA 4404 was transformed with pVBLCYB by triparental mating, using the helper plasmid pRK2103 [12]. Stem explants of tomato were transformed and plants regenerated as described previously



**Figure 2:** Schematic diagram of the *LCYB* construct pVBLCY.

*UBQ3*, ubiquitin three promoter; *LCYB*, *N. tabacum* lycopene  $\beta$ -cyclase; *nptII*, nopaline synthase terminator; *AscI*, *BamHI*, *AgeI* restriction sites; LB, left border; RB, right border.

[13]. Kanamycin (50 mg/ml) and carbenicillin (250 mg/ml) were used to select resistant transformants.

## 2.4 DNA and RNA analyses

DNA was extracted for PCR analysis from leaf material (ca. 200 mg), as described previously [13]. The presence of the transgene in the primary transformants ( $T_0$ ) was confirmed by PCR using primers designed to amplify a 500 bp internal fragment of *LCYB*. Southern blot analysis of the  $T_0$  and first-generation ( $T_1$ ) plants was carried out using the cetyltrimethylammonium bromide (CTAB) method [14]. DNA was digested with the appropriate restriction enzymes, fragments separated in 1% (w/v) agarose gels and transferred to Hybond N<sup>+</sup> (Amersham, Little Chalfont, UK) overnight in 20% saline sodium citrate (SSC) solution. <sup>32</sup>P-labelled fragments (using the Stratagene Random primer labelling kit, Prime It II), corresponding to the *LCYB* and *nptII* regions of the transgene, were used to probe the filters. Hybridisation, washing and detection were performed as described in the Hybond N<sup>+</sup> instructions. Northern blot analysis was carried out on tomato fruit of the  $T_0$  and  $T_1$  generations, harvested at 7 dpb [15]. Total RNA (20  $\mu$ g) from the fruit was electrophoresed on 1.4% agarose/formaldehyde gels and then transferred to Hybond-N<sup>+</sup> in 20x SSC overnight. The 1.6-kb cDNA fragment was radiolabelled with <sup>32</sup>P, as described above. Following pre-hybridisation, the filter was hybridised at 65°C in 7.5% SDS, 100 mM K phosphate buffer, pH 7.5. Each filter was washed for 15 min in 2x SSC, 0.5% SDS and then twice for 15 min with 1x SSC, 0.5% SDS and finally monitored for radioactivity for 24 h with a phosphorimager (MACBAS V2.2, Fuji Film Ltd, Bedford, UK).

## 2.5 Treatment with CPTA

Sterilised seeds (ca. 30) from the homozygous  $T_1$  line C4S1-15 and azygous control were germinated on Murashige and Skoog (MS) agar containing the herbicide

2-(4-chlorophenylthio) triethylamine (CPTA, a gift from Prof. P. Böger) at 30 and 40  $\mu$ M. The seedlings were grown in a controlled environment with a 16-h photoperiod and day and night temperatures of 23 and 19°C, respectively. They were assessed for germination and bleaching after 2 weeks and samples taken for isoprenoid analysis.

## 2.6 Analysis of isoprenoids

The isoprenoids of  $T_0$ ,  $T_1$  and  $T_2$  progeny of leaf and 7 dpb fruit were extracted, separated, identified and quantified by HPLC analyses, as described in [16].

## 3 Results

### 3.1 Plants expressing the *LCYB* gene from tobacco showed increased levels of $\beta$ -carotene in ripe fruit and changes to leaf carotenoids

Following transformation, progeny were selected on the basis of kanamycin resistance (*nptII*) and PCR analysis. During regeneration, growth and development of vegetative tissues from all transgenic lines appeared phenotypically normal. However, as fruit ripening commenced, virtually all transgenic lines showed a greater degree of orange colouration in their fruit compared to the control. Analysis of the  $T_0$  generation showed that increases in  $\beta$ -carotene content of the fruit were responsible for the altered colouration (Table 1). Of the  $T_0$  transformants, 85% exhibited a lycopene-to- $\beta$ -carotene ratio within a range of 2:1 to 6:1, due to increases in  $\beta$ -carotene, with the wild-type (Ailsa Craig) ratio being 7.46:1. Analysis of the  $T_1$  and  $T_2$  generations established that the phenotype was inherited (Table 1).

Insertions at a single locus were determined by Southern blotting using a *nptII* probe. Of the 20 transformants,

**Table 1:** Carotenoid contents of fruit of lines of transgenic tomato plants expressing the tobacco  $\beta$ -lycopene cyclase gene, *LCYB*.

Plant line	Generation/Zygoty	Carotenoid content ( $\mu\text{g/g DW}$ )			
		Lycopene	$\beta$ -Carotene	Total <sup>a</sup>	Ratio L: $\beta$ -C
Wild type	–	2133 $\pm$ 479	290 $\pm$ 29	2560 $\pm$ 508	7.46
603 C4S1	T <sub>0</sub>	5546 $\pm$ 1397	1711 $\pm$ 103	7394 $\pm$ 1490	3.24
603 C1S1	T <sub>0</sub>	1192 $\pm$ 370	783 $\pm$ 46	2112 $\pm$ 416	1.52
603 C2S1	T <sub>0</sub>	2560 $\pm$ 297	1247 $\pm$ 37	3944 $\pm$ 334	2.05
603 C1S2	T <sub>0</sub>	2773 $\pm$ 471	928 $\pm$ 81	3838 $\pm$ 552	2.99
Azygous control <sup>b</sup>	T <sub>1</sub>	2009 $\pm$ 134	329 $\pm$ 53	2669 $\pm$ 204	6.11
603 C4S1-1	T <sub>1</sub> , He	2240 $\pm$ 99	609 $\pm$ 82	2991 $\pm$ 181	3.68
603 C4S1-15	T <sub>1</sub> , Ho	2069 $\pm$ 40	725 $\pm$ 28	2936 $\pm$ 32	2.85
603 C4S1-5	T <sub>1</sub> , Ho	1493 $\pm$ 43	783 $\pm$ 53	2418 $\pm$ 96	1.91
603 C1S1-4	T <sub>1</sub> , He	1493 $\pm$ 79	406 $\pm$ 6	2041 $\pm$ 85	3.68
603 C1S1-18	T <sub>1</sub> , Ho	1770 $\pm$ 104	667 $\pm$ 59	2579 $\pm$ 163	2.65
603 C1S1-19	T <sub>1</sub> , Ho	2560 $\pm$ 110	580 $\pm$ 16	3282 $\pm$ 126	4.41
603 C2S1-1	T <sub>1</sub> , Ho	2133 $\pm$ 85	725 $\pm$ 36	3000 $\pm$ 122	2.94
603 C4S1-15	T <sub>2</sub> , Ho	3250 $\pm$ 1455	623 $\pm$ 175	4034 $\pm$ 163	5.22

Fruit were harvested at 7 dpb. <sup>a</sup>Includes lutein, phytofluene,  $\zeta$ -carotene and phytoene at 1.8, 0.3, 0.14 and 1.3% of the total carotenoids, respectively. <sup>b</sup>Represents the mean of azygous plants from 603 C4S1, C1S1 and C2S1 lines. He, hemizygous; Ho, homozygous; T<sub>0</sub>, primary transformants; T<sub>1</sub>, second generation; T<sub>2</sub>, third generation. All values are the means of three replicates  $\pm$  S.E.

five lines were single copy, while the others had copy numbers ranging from 2 to 4 for the *nptII* gene (data not shown). Homozygous lines were generated from single-copy primary transformants, showing a range of  $\beta$ -carotene contents (603 C1S1, C1S2, C2S1 and C4S1, with lycopene-to- $\beta$ -carotene ratios of 1.5, 2.99, 2.05 and 3.24, respectively; Table 1). Zygoty was assigned by Southern analysis using the *nptII* probe, co-segregation was assessed by expression of the transgene and inheritance of the phenotype by HPLC analysis. Co-segregation occurred in all cases and the high  $\beta$ -carotene phenotype was retained in the third generation (T<sub>2</sub>). As shown in Supplementary Material, Figure 2, the ratio of lycopene-to- $\beta$ -carotene ratio differed between azygous, hemizygous and homozygous fruit. Although significant elevations in  $\beta$ -carotene were found in fruit of the transgenic lines, the content of biosynthetically related carotenoids such as phytoene,  $\gamma$ -carotene,  $\delta$ -carotene,  $\alpha$ -carotene, lutein, zeaxanthin, violaxanthin and neoxanthin was not altered.

The pattern of  $\beta$ -carotene and lycopene geometric isomers found in the transgenic fruit were identical to the wild type (i.e.  $\beta$ -carotene 90–95% all-*trans*, 10–5% *cis*; all-*trans* lycopene 96–98%, 15-*cis*- lycopene 1–4% and 9-*cis* lycopene 0.6%). Alterations in other fruit isoprenoids were minor, with increases of 1.35-, 1.50- and 1.01-fold for tocopherols, plastoquinone and ubiquinone, respectively, in line 603 C4S1-15 (data not shown).

There were also changes to the carotenoid levels in transgenic leaves. For example, in the leaves of line 603 C4S1, all the carotenoids were increased, with lutein exhibiting the largest elevation (Table 2).

### 3.2 *LCYB* gene expression correlated with increased $\beta$ -carotene levels in fruit

Northern blot analysis showed a positive correlation between the level of transgene expression and increased

**Table 2:** Carotenoid content of leaves of Ailsa Craig and a *LCYB* transgenic line.

Genotype	Carotenoid ( $\mu\text{g/g DW}$ )					
	Total	$\beta$ -Carotene	Lutein	Anthera	Viola	Neox
Ailsa Craig	83.7 $\pm$ 1.5	22.0 $\pm$ 0.36	33.7 $\pm$ 4.1	6.3 $\pm$ 1.5	13.4 $\pm$ 1.5	8.3 $\pm$ 0.04
% Total		26.3	40.3	7.53	16.0	9.90
603 C4S1-15	275.6 $\pm$ 0.73	33.7 $\pm$ 1.7	147.3 $\pm$ 0.40	21.4 $\pm$ 0.50	42.2 $\pm$ 0.13	31.0 $\pm$ 1.5
% Total		12.2	53.4	7.80	15.3	11.2

All values are the means of three replicates  $\pm$  S.E. Anthera, antheraxanthin; Viola, violaxanthin, Neox, neoxanthin. Lycopene and  $\alpha$ -carotene were not detected.

$\beta$ -carotene formation (Figure 3). A comparison of expression between chloroplast-containing (leaf and green fruit) tissues and chromoplast-containing ripe fruit showed that although the *UBQ3* promoter has been generally found to be constitutive in plants, *LCYB* transcript abundance was approximately 10-fold greater in ripe fruit (data not shown).

### 3.3 Transgenic lines showed tolerance to CPTA


A comparison of Ailsa Craig seedling and those of the transgenic line 603 C4S1-15, in the presence of CPTA, revealed dramatic differences in plant vigour. The control seedlings showed significant chlorosis, whilst those of

the transformed line remained green (Supplementary Figure 3). Pigment analysis of the leaves revealed lower levels of carotenoids in the control (55% reduction at 40  $\mu$ M CPTA), and qualitatively,  $\beta$ -carotene levels were some 4.3-fold less in the control leaves (Table 3). Increases in lycopene in these leaves were also observed. There was also an increase in lutein levels in 603 C4S1-15 in comparison to Ailsa Craig at both CPTA concentrations.

## 4 Discussion

There have been several reports of using genetic modification for targeted and untargeted increases in carotenoid levels in crop plants, albeit with varying degrees of success [17]. With respect to  $\beta$ -carotene, a modest, 1.5-fold, increase was achieved using the *PSY-1* gene with a constitutive promoter (CaMV 35S) [18], whilst transformation with the *Arabidopsis LCYB* and *PDS* promoter elevated  $\beta$ -carotene some 7-fold [19]. Alternatively, a bacterial *PDS*, *CrtI* resulted in a 3-fold increase, but a decrease in lycopene content [20]. This was surprising, as *CRTI* catalyses lycopene, not  $\beta$ -carotene biosynthesis. Such an effect may be caused by a feed-forward mechanism for the upregulation of *LCY* or *CYCB*, as described previously in tomato fruit [21]. Therefore, although increases in  $\beta$ -carotene were found, unintended perturbations to other carotenoids and related isoprenoids occurred, or else the total carotenoid content was reduced.

In contrast, in the present study, levels of  $\beta$ -carotene have been increased up to 6-fold, without a decrease in lycopene, perturbations to other fruit carotenoids, nor to the detriment of plant vigour. The phenotype was stable for at least three generations (Table 1). Therefore, the transgenic fruit contain an improved carotenoid profile compared with non-transgenic cultivars and introgression lines [22]. This is the first report of using the *UBQ3* promoter in tomato, although it has been used with other dicot plants [23]. Although constitutive, the *UBQ* promoter



**Figure 3:** Northern blot analysis showing transcript abundance of the tobacco *LCYB* gene in transgenic lines. (A) Primary ( $T_0$ ) transformants: 1, control Ailsa Craig (1:7); 2, C1S1 (1:1.5); 3, C1S2 (1:3.0), 4, t C3S1 (not determined); 5, C4S1 (1:3.2); 6, C4S2 (1:5), 7, C5S1 (1:6.5). (B)  $T_1$  progeny: 1, Ailsa Craig control (1:7); 2, C4S1-1, hemizygous (1:3.7); 3, C4S1-5, homozygous (1:1.9); 4, C4S1-8, homozygous (1:2); 5, C4S1-12, azygous (1:6); 6, C4S1-15,  $T_2$  homozygous (1:5.2), 7, C4S1-18, azygous (1:8); 8, Control Ailsa Craig (1:7). The lycopene-to- $\beta$ -carotene ratios are provided in parenthesis. Uniform RNA loading was verified by probing with actin.

**Table 3:** Carotenoid content of leaves from plants treated with CPTA.

Plant type and treatment	Carotenoid content ( $\mu$ g/g FW)						
	Total	Lycopene	$\beta$ -Carotene	$\alpha$ -Carotene	Lutein	Neox	Viola
AC + 30 $\mu$ M CPTA	62.1	2.9 $\pm$ 0.1 (4.6)	4.7 $\pm$ 0.99 (7.5)	7.1 $\pm$ 0.5 (11.3)	44.7 $\pm$ 7.6 (72.0)	1.8 $\pm$ 0.3 (2.4)	0.9 $\pm$ 0.4 (1.4)
AC + 40 $\mu$ M CPTA	47.4	1.4 $\pm$ 0.1 (3.0)	1.8 $\pm$ 0.2 (3.8)	7.3 $\pm$ 0.7 (15.6)	34.7 $\pm$ 0.1 (73.0)	0.4 $\pm$ 0.04 (2.1)	1.8 $\pm$ 0.2 (3.8)
603 C4S1-15 + 30 $\mu$ M CPTA	76.9	0.6 $\pm$ 0.13 (0.8)	7.2 $\pm$ 1.5 (9.5)	6.1 $\pm$ 1.2 (8.0)	58.0 $\pm$ 6.1 (76.2)	2.2 $\pm$ 0.2 (2.8)	2.8 $\pm$ 0.2 (3.6)
603 C4S1-15 + 40 $\mu$ M CPTA	106.2	0.6 $\pm$ 0.07 (0.6)	7.8 $\pm$ 0.7 (7.5)	3.3 $\pm$ 0.73 (3.1)	90.6 $\pm$ 6.9 (85)	0.6 $\pm$ 0.06 (0.6)	3.3 $\pm$ 0.13 (3.1)

Percentage of total carotenoids shown in brackets. Values are the means of three replicates  $\pm$  s.e. Viola, violaxanthin, Neox, neoxanthin. AC, Ailsa Craig; 603 C4S1-15, *LCYB* line. Refer to Table 2 for control without CPTA treatment.

gave differential expression levels of the transgene in vegetative and fruit tissues, with lower expression found in chloroplast-containing tissues that probably prevented the occurrence of gene silencing and/or co-suppression of the endogenous gene [24], since the homology of the tobacco and tomato  $\beta$ -cyclases is 87.2% (Supplemental Material, Figure 1). Since the endogenous *LCYB* gene is down-regulated during fruit ripening [9], co-suppression is not an issue. Despite the low expression in vegetative tissue, changes to carotenoids do occur (Table 2), most specifically a 4.4-fold increase in lutein, suggesting that *LCYB* catalyses the rate-limiting step in the formation of this xanthophyll.

Growth of the transgenic lines with CPTA did not cause changes to plant vigour, unlike the chlorosis shown with the wild type (Supplementary Material, Figure 3). Lycopene did not accumulate in the *LCYB* line treated with CPTA (Table 3). Tolerance to CPTA in the transgenic line may be due to the increase in *LCYB* protein levels, and/or because the tobacco *LCYB* is intrinsically more tolerant to CPTA than that of tomato. Therefore, such an approach is feasible for the production of a crop resistant to this herbicide and for screening putative bleaching herbicides that have different modes of action to CPTA [25]. The increase in total carotenoids in the transgenic line is consistent with the report that CPTA modulates mRNA levels of carotenoid genes [26].

Nutritionally, an average ripe *LCYB* transgenic fruit contains virtually all the recommended daily allowance (RDA) for vitamin A compared to the wild type, which has about 39% of the RDA [27]. For comparison, the *crtI* transgenic tomato contains similar levels of  $\beta$ -carotene (ca. 70% RDA), but only 50% of the wild type lycopene content. Thus, this new genotype not only has a significantly elevated provitamin A content, but maintains its level of lycopene, compared to the wild type.

In summary, expression of *LCYB* under the control of the *UBQ3* promoter has resulted in the specific elevation of  $\beta$ -carotene in ripe tomato fruit, but no changes to other fruit carotenoids. There is much current debate concerning the acceptance of genetically modified plants. The present study has produced a tomato with the potential to reduce vitamin A deficiency and contribute towards a high antioxidant diet. Such germplasm will provide a valuable resource for further detailed analysis [28] and add to the genetic resources for nutritional benefit.

**Acknowledgments:** We are grateful to Dr Susanne Römer for providing the *NtLCYB* cDNA clone. The work was supported by the European Community FAIR Programme (# CT9616333). Syngenta Agrochemicals is gratefully

acknowledge for the provision of glasshouse space and some laboratory facilities. We thank Karen Bacon and Carol Fisher for maintenance of the tomato plants and Mark Harker for some of the Southern analyses.

## References

1. Bramley PM. Carotenoid biosynthesis and chlorophyll degradation. In: Seymour GB, Tucker GA, Poole M, Giovannoni J, editors. Molecular biology and biochemistry of fruit ripening. New Jersey: Wiley, 2013:75–116.
2. Perveen R, Suleria HA, Anjum FM, Butt MS, Pasha I, Ahmad S. Tomato (*Solanum lycopersicum*) carotenoid and lycopenes chemistry; metabolism, absorption, nutrition and allied health claims – a comprehensive review. Crit Rev Fd Sci Nutr 2105;55:919–29.
3. Holzapfel NP, Holzapfel BM, Champ S, Feldthausen J, Clements J, Huttmacher DW. The potential role of lycopene for the prevention and therapy of prostate cancer: from molecular mechanisms to clinical evidence. Int J Mol Sci 2013;14:14620–46.
4. Sommer A, Vyas KS. A global clinical view on vitamin A and carotenoids. Am J Clin Nutr 2012;96:12045–65.
5. Stevens GA, Bennett JE, Hennocq Q, Lu Y, De-Regil LM, Rogers L, et al. Trends and mortality effects of vitamin A deficiency in children in 138 low-income and middle-income countries between 1991 and 2013: a pooled analysis of population-based surveys. Lancet Glob Health 2015;3:e528–36.
6. Giuliano G, Tavazza R, Diretto G, Beyer P, Taylor MA. Metabolic engineering of carotenoid biosynthesis in plants. Trends Biotechnol 2008;26:139–45.
7. Isaacson T, Ronen G, Zamir D, Hirschberg J. Cloning *tangerine* from tomato reveals a carotenoid isomerase essential for the production of  $\beta$ -carotene and xanthophylls in plants. Plant Cell 2002;14:333–42.
8. Park H, Kreunen SS, Cuttriss AJ, DellaPenna D, Pogson BJ. Identification of a carotenoid isomerase provides insight in carotenoid biosynthesis, prolamellar body formation and photomorphogenesis. Plant Cell 2002;14:321–32.
9. Pecker I, Gabbay R, Cunningham FX, Hirschberg J. Cloning and characterisation of the cDNA for lycopene  $\beta$ -cyclase from tomato reveals a decrease in its expression during fruit ripening. Plant Mol Biol 1990;30:807–19.
10. Hirschberg J. Carotenoid biosynthesis in flowering plants. J Curr Opin Plant Biol 2001;4:210–8.
11. Simpson DJ, Baqar MR, Lee TH. Chemical regulation of plastid development III. Effect of light and CPTA on chromoplast ultrastructure and carotenoids in *Capsicum annum*. Z Pflanzenphysiol 1977;82:189–209.
12. Hoekema A, Hirsch PR, Hooykas PJ, Schilperoort RA. A binary vector strategy based on separation of *vir* and T-region of the *Agrobacterium tumefaciens* Ti-plasmid. Nature 1983;303:179–80.
13. Bird CR, Smith CJ, Ray JA, Moureau P, Bevan MW, Bird AS, et al. the tomato polygalacturonase gene and ripening-specific expression in transgenic plants. Plant Mol Biol 1988;11:651–62.
14. Murray MG, Thompson WF. Rapid isolation of high molecular weight plant DNA. Nucl Acid Res 1980;8:4321–25.

15. Sambrook J, Fritsch FF, Maniatis T. Molecular cloning: a laboratory manual, 1st ed. New York: Cold Spring Harbor Laboratory Press, 1989.
16. Fraser PD, Pinto ME, Holloway DE, Bramley PM. Application of HPLC with photodiode array detection to the metabolic profiling of plant isoprenoids. *Plant J* 2000;24:551–8.
17. Fraser PD, Enfissi EM, Bramley PM. Genetic engineering of carotenoid formation in tomato fruit and the potential application of systems and synthetic biology approaches. *Arch Biochem Biophys* 2009;m483:196–204.
18. Fraser, PD, Römer S, Shipton CA, Mills PB, Kiano JW, Misawa N, et al. Evaluation of transgenic tomatoes expressing an additional phytoene synthase in a fruit-specific manner. *Proc Natl Acad Sci* 2002;99:1092–7.
19. Rosati C, Aquiliani R, Dharmapuri S, Pallara P, Marusic C, Tavazza R, et al. Metabolic engineering of  $\beta$ -carotene and lycopene content in tomato fruit. *Plant J* 2000;24:413–9.
20. Römer S, Fraser PD, Kiano JW, Shipton CA, Misawa N, Schuch W, et al. Elevation of the provitamin A content of transgenic tomato plants. *Nature Biotech* 2000;18:666–9.
21. Fraser PD, Enfissi EM, Halket JM, Truesdale MR, Yu D, Gerrish CR, et al. Manipulation of phytoene levels in tomato fruit: Effects on isoprenoids, plastids, and intermediary metabolism. *Plant Cell* 2007;19:3194–211.
22. Fernie AR, Tadmor Y, Zamir D. Natural genetic variation for improving crop quality. *Curr Opin Plant Biol* 2006;9:196–202.
23. Norris SR, Meyer SE, Callis J. The intron of *Arabidopsis thaliana* polyubiquitin genes is conserved in location and is a quantitative determinant of chimeric gene expression. *Plant Mol Biol* 1993;21:985–06.
24. Fray RG, Grierson D. Identification and genetic analysis of normal and mutant phytoene synthase genes of tomato by sequencing, complementation and co-suppression. *Plant Mol Biol* 1993;22:589–602.
25. Windhovel U, Sandmann G, Böger P. Genetic engineering of resistance to bleaching herbicides affecting phytoene desaturase and lycopene cyclase in cyanobacterial carotenogenesis. *Pest Biochem Physiol* 1997;57:68–78.
26. Al-Babili S, Hartung W, Kleinig H, Beyer P. CPTA modulates levels of carotenogenic proteins and their mRNAs and affects carotenoid and ABA content as well as chloroplasts structure in *Narcissus pseudonarcissus* flowers. *Plant Biol* 1999;1:607–12.
27. NIH Fact Sheet <https://ods.od.nih.gov>.
28. Giuliano G. Plant carotenoids: genomics meets multi-gene engineering. *Curr Opin Plant Biol* 2014;19:111–17.

---

**Supplemental Material:** The online version of this article (DOI: 10.1515/znc-2016-0102) offers supplementary material, available to authorized users.