Crystal structure of (1'*R**,2*R**,3*R**,4*S**,5*S**,6*S**)-2,3,4,5-tetrahydro-6-hydroxy-2-(1'-hydroxyethyl)-3-hydroxymethyl-5-methoxy-2,4-heptanofuran, C₁₁H₁₆O[CH(OH)CH₃](OH)(CH₂OH)(OCH₃)

K. Peters, E.-M. Peters

Max-Planck-Institut für Festkörperforschung, Heisenbergstraße 1, D-70506 Stuttgart, Germany

M. Kasch and W. Tochtermann

Christian-Albrechts-Universität Kiel, Institut für Organische Chemie, Olshausenstraße 40, D-24098 Kiel, Germany

Received August 14, 1997, CSD-No. 409074

Table 1. Parameters used for the X-ray data collection

Crystal:	colorless plate, size 0.15 x 0.5 x 0.75 mm
Wavelength:	Mo K α radiation (0.71073 Å)
μ :	0.90 cm ⁻¹
Diffractometer:	Siemens P4
Scan mode:	ω
$T_{\text{measurement}}$:	293 K
$2\theta_{\text{max}}$:	55°
$N(hkl)_{\text{unique}}$:	3561
Criterion for F_o :	$F_o > 3 \sigma(F_o)$
$N(\text{param})_{\text{refined}}$:	194
Program:	SHELXTL-plus

Source of material: The title compound was prepared as described in ref. 1.

C₁₅H₂₈O₅, monoclinic, $P12_1/n1$ (No. 14), $a = 8.021(1)$ Å, $b = 21.679(3)$ Å, $c = 9.418(1)$ Å, $\beta = 108.89(1)^\circ$, $V = 1549.5$ Å³, $Z = 4$, $R(F) = 0.059$, $R_w(F) = 0.061$.

Table 2. Final atomic coordinates and displacement parameters (in Å²)

Atom	Site	x	y	z	U_{iso}
H(1)	4e	-0.0999(3)	0.7789(1)	0.4978(2)	0.08
H(3)	4e	0.1426(3)	0.82005(8)	0.8566(2)	0.08
H(4)	4e	-0.1685(3)	0.87625(9)	0.8466(2)	0.08
H(5)	4e	0.0011(3)	0.97315(9)	0.7768(2)	0.08
H(6A)	4e	0.2346(3)	0.85773(9)	0.5424(2)	0.08
H(6)	4e	0.288(4)	0.758(1)	0.525(3)	0.072(9)
H(7A)	4e	0.3711(3)	0.8104(1)	0.8277(3)	0.08
H(7B)	4e	0.4808(3)	0.8220(1)	0.7194(3)	0.08
H(8A)	4e	0.3244(3)	0.9251(1)	0.7889(3)	0.08
H(8B)	4e	0.5122(3)	0.9190(1)	0.7727(3)	0.08
H(9A)	4e	0.5606(3)	0.9420(1)	1.0160(3)	0.08
H(9B)	4e	0.6050(3)	0.8717(1)	1.0145(3)	0.08
H(10A)	4e	0.3379(3)	0.8467(1)	1.0425(3)	0.08
H(10B)	4e	0.4497(3)	0.8874(1)	1.1771(3)	0.08
H(10)	4e	-0.287(5)	0.873(2)	0.545(4)	0.11(1)
H(11A)	4e	0.2308(4)	0.9587(1)	0.9601(3)	0.08
H(11B)	4e	0.2529(4)	0.9574(1)	1.1317(3)	0.08
H(12A)	4e	0.0524(3)	0.8705(1)	1.0694(3)	0.08
H(12B)	4e	-0.0268(3)	0.9370(1)	1.0318(3)	0.08
H(13A)	4e	-0.0006(4)	0.8368(1)	0.3366(3)	0.08
H(13B)	4e	-0.1270(4)	0.8908(1)	0.3453(3)	0.08
H(13C)	4e	-0.2053(4)	0.8281(1)	0.2701(3)	0.08
H(14A)	4e	-0.3672(4)	1.0126(1)	0.4952(4)	0.08
H(14B)	4e	-0.1849(4)	0.9951(1)	0.4783(4)	0.08
H(14C)	4e	-0.1926(4)	1.0392(1)	0.6079(4)	0.08
H(30A)	4e	-0.0209(3)	0.73666(9)	0.7481(2)	0.08
H(30B)	4e	-0.1900(3)	0.77308(9)	0.7474(2)	0.08
H(30)	4e	0.051(4)	0.724(1)	0.982(3)	0.071(9)

Table 3. Final atomic coordinates and displacement parameters (in Å²)

Atom	Site	x	y	z	U_{11}	U_{22}	U_{33}	U_{12}	U_{13}	U_{23}
O(1)	4e	0.0330(2)	0.91399(6)	0.6330(1)	0.0370(7)	0.0297(6)	0.0392(8)	-0.0014(5)	0.0177(6)	0.0059(5)
C(1)	4e	-0.1152(3)	0.8228(1)	0.4992(2)	0.042(1)	0.047(1)	0.035(1)	-0.0066(9)	0.0123(9)	-0.0035(9)
C(2)	4e	0.0387(2)	0.84660(8)	0.6343(2)	0.035(1)	0.0296(9)	0.0322(9)	-0.0025(7)	0.0140(8)	0.0010(7)
C(3)	4e	0.0268(3)	0.82919(8)	0.7894(2)	0.0343(9)	0.0297(9)	0.0327(9)	-0.0019(7)	0.0164(7)	0.0017(7)

Table 3. (Continued)

Atom	Site	x	y	z	U ₁₁	U ₂₂	U ₃₃	U ₁₂	U ₁₃	U ₂₃
C(4)	4e	−0.0531(3)	0.88589(9)	0.8430(2)	0.045(1)	0.033(1)	0.039(1)	0.0014(8)	0.0234(9)	0.0032(8)
C(5)	4e	−0.0563(3)	0.93639(9)	0.7280(2)	0.039(1)	0.032(1)	0.043(1)	0.0022(8)	0.0175(9)	0.0042(8)
O(5)	4e	−0.2338(2)	0.95152(7)	0.6461(2)	0.0400(8)	0.0426(8)	0.064(1)	0.0084(7)	0.0229(7)	0.0187(7)
C(6)	4e	0.2161(3)	0.82909(9)	0.6136(2)	0.041(1)	0.036(1)	0.040(1)	−0.0017(8)	0.0228(9)	−0.0017(8)
O(6)	4e	0.1988(2)	0.76574(7)	0.5684(2)	0.057(1)	0.0423(9)	0.067(1)	−0.0031(7)	0.0368(9)	−0.0137(8)
C(7)	4e	0.3837(3)	0.8363(1)	0.7492(3)	0.036(1)	0.047(1)	0.048(1)	0.0054(9)	0.0200(9)	0.002(1)
C(8)	4e	0.4303(3)	0.9009(1)	0.8157(3)	0.042(1)	0.051(1)	0.053(1)	−0.004(1)	0.014(1)	0.006(1)
C(9)	4e	0.5122(3)	0.9017(1)	0.9865(3)	0.049(1)	0.055(1)	0.052(1)	−0.001(1)	−0.001(1)	−0.001(1)
C(10)	4e	0.3850(3)	0.8871(1)	1.0717(3)	0.067(2)	0.051(1)	0.039(1)	0.004(1)	0.002(1)	−0.002(1)
O(10)	4e	−0.2809(2)	0.83655(8)	0.5175(2)	0.0359(8)	0.054(1)	0.056(1)	−0.0081(7)	0.0089(7)	−0.0051(8)
C(11)	4e	0.2308(4)	0.9328(1)	1.0428(3)	0.070(2)	0.049(1)	0.048(1)	0.000(1)	0.012(1)	−0.012(1)
C(12)	4e	0.0444(3)	0.9060(1)	1.0065(3)	0.070(2)	0.043(1)	0.041(1)	0.006(1)	0.029(1)	−0.0041(9)
C(13)	4e	−0.1117(4)	0.8468(1)	0.3491(3)	0.067(2)	0.081(2)	0.034(1)	−0.012(1)	0.007(1)	0.001(1)
C(14)	4e	−0.2456(4)	1.0040(1)	0.5486(4)	0.057(2)	0.066(2)	0.093(2)	0.015(1)	0.025(1)	0.045(2)
C(30)	4e	−0.0657(3)	0.76841(9)	0.7970(2)	0.049(1)	0.033(1)	0.049(1)	−0.0040(9)	0.025(1)	0.0043(9)
O(30)	4e	−0.0351(2)	0.75171(7)	0.9505(2)	0.062(1)	0.0452(9)	0.063(1)	0.0134(8)	0.0417(9)	0.0234(8)

References

1. Kasch, M.: Racemische und optisch aktive überbrückte Furanoside als Bausteine zur Synthese modifizierter Nucleoside. Dissertation, Universität Kiel, Germany 1996.
2. Sheldrick, G. M.: Program Package SHELXTL-plus. Release 4.1. Siemens Analytical X-Ray Instruments Inc., Madison (WI 53719), USA 1990.