Crystal structure of (3*aR*,6*aR*)-3-[(1*R*,2*S*,3*S*)-1,3:2,4-di-*O*-ethylidene-1,2,3,4-tetrahydroxy-1-butyl]-3*a*,6*a*-dihydrofuro[2,3-*d*]isoxazole, C₁₃H₁₇NO₆

S. Henkel, T. Leibold and V. Jäger

Universität Stuttgart, Institut für Organische Chemie, Pfaffenwaldring 55, D-70569 Stuttgart, Germany

Received January 13, 1997, CSD-No. 402796

Table 3. (Continued)

Atom	Site	x	y	z	U ₁₁	U ₂₂	U ₃₃	U ₁₂	U ₁₃	U ₂₃
C(81)	4a	0.7099(7)	0.5435(6)	0.0491(3)	0.071(4)	0.096(4)	0.055(3)	−0.001(4)	0.026(3)	0.000(3)
O(9)	4a	0.6261(4)	0.6046(3)	0.1650(1)	0.052(2)	0.055(2)	0.042(2)	−0.001(2)	0.003(1)	0.005(1)
C(10)	4a	0.5287(6)	0.6954(4)	0.2090(2)	0.059(3)	0.034(2)	0.038(2)	−0.006(2)	0.002(2)	0.004(2)
C(11)	4a	0.5627(6)	0.6619(4)	0.2853(2)	0.062(3)	0.030(2)	0.040(2)	−0.011(2)	−0.003(2)	−0.001(2)
C(12)	4a	0.4858(7)	0.7485(5)	0.3436(2)	0.084(4)	0.041(2)	0.038(2)	−0.001(3)	−0.005(3)	−0.001(2)
O(13)	4a	0.3130(5)	0.7153(4)	0.3517(2)	0.074(2)	0.101(3)	0.054(2)	−0.005(2)	0.007(2)	−0.008(2)
C(14)	4a	0.304(1)	0.6422(7)	0.4146(3)	0.128(6)	0.090(5)	0.070(4)	−0.025(5)	0.035(4)	−0.008(4)
C(15)	4a	0.440(1)	0.6285(7)	0.4496(3)	0.159(7)	0.079(4)	0.044(3)	0.003(5)	0.020(4)	0.013(3)
C(16)	4a	0.5749(8)	0.6864(6)	0.4076(2)	0.105(5)	0.058(3)	0.046(2)	0.014(3)	−0.018(3)	−0.002(2)
O(17)	4a	0.6826(6)	0.5729(4)	0.3792(2)	0.122(4)	0.076(2)	0.051(2)	0.025(3)	−0.027(2)	0.005(2)
N(18)	4a	0.6653(6)	0.5670(4)	0.3045(2)	0.088(4)	0.051(2)	0.050(2)	0.014(3)	−0.010(2)	0.001(2)

Acknowledgments. We are grateful to the Fonds der Chemischen Industrie, Bayer AG and Hoechst AG for financial support. We also thank Ms. S. Bucovarova for experimental assistance and Dr. Wolfgang Frey for help with the preparation of the files.

References

- Leibold, T.: Furan-Nitriloxid-Cycloaddukte - diastereomerenreine Vorläufer für die Synthese von biologisch aktiven Desoxynojirimycin- und Castanospermin-Analoga. Dissertation, Universität Stuttgart, Germany 1995.
- Müller, I.; Jäger, V.: Synthesis of amino sugars via isoxazolines. Nitrile oxide-furan adducts as key intermediates in the isoxazoline route towards novel amino sugar derivatives. *Tetrahedron* **41** (1985) 3519-3528.
- Jäger, V.; Müller, R.; Leibold, T.; Hein, M.; Schwarz, M.; Fengler, M.; Jaroskova, L.; Pätzelt, M.; LeRoy, P.-Y.: Synthesis of glycosidase-inhibiting iminopolyols via isoxazolines. *Bull. Soc. Chim. Belg.* **103** (1994) 491-507.
- Liu, K.-C.; Shelton, B. R.; Howe, R. K.: A particularly convenient preparation of benzohydroximinoyl chlorides (nitrile oxide precursors). *J. Org. Chem.* **45** (1980) 3916-3918.
- Henkel, S.; Leibold, T.; Jäger, V.: Crystal structure of 1,3:2,4-di-*O*-ethylidene-L-xylose oxime, C₉H₁₅NO₅. *Z. Kristallogr. NCS* **213** (1998) 65-66.
- Henkel, S.; Leibold, T.; Jäger, V.: Crystal structure of (3a*S*,6a*S*)-3-[(1*R*,2*S*,3*S*)-1,3:2,4-di-*O*-ethylidene-1,2,3,4-tetrahydroxy-1-butyl]-3a,6a-dihydrofuro[2,3-*d'*]isoxazole, C₁₃H₁₇NO₆. *Z. Kristallogr. NCS* **213** (1998) 69-70.
- Sheldrick, G. M.: Phase Annealing in SHELX-90: Direct Methods for Large Structures. *Acta Crystallogr. A* **46** (1990) 467-473.
- Sheldrick, G. M.: SHELXL-93. Program for refining crystal structures. University of Göttingen, Germany 1993.