Crystal structure of (3*aS*,6*aS*)-3-[(1*R*,2*S*,3*S*)-1,3:2,4-di-*O*-ethylidene-1,2,3,4-tetrahydroxy-1-butyl]-3*a*,6*a*-dihydrofuro[2,3-*d*]isoxazole, C₁₃H₁₇NO₆

S. Henkel, T. Leibold and V. Jäger

Universität Stuttgart, Institut für Organische Chemie, Pfaffenwaldring 55, D-70569 Stuttgart, Germany

Received January 13, 1997, CSD-No. 402797

Source of material: The title compound was prepared (see ref. 1) by 1,3-dipolar cycloaddition of furan (see ref. 2 and 3) with the corresponding hydroximoyl chloride which was derived (see ref. 4) from the corresponding L-xylose oxime (see ref. 5). The mixture of diastereomeric cycloadducts [(3*aR*,6*aR*) / (3*aS*,6*aS*) = 70 : 30] was separated by MPLC and both isomers were recrystallized from petroleum ether / ethyl acetate (see refs. 1 and 6).

C₁₃H₁₇NO₆, orthorhombic, *P*2₁2₁2₁ (No. 19), *a* = 6.541(1) Å, *b* = 8.938(2) Å, *c* = 23.161(4) Å, *V* = 1354.1 Å³, *Z* = 4, *R*(*F*) = 0.046, *R*_w(*F*²) = 0.112.

Table 1. Parameters used for the X-ray data collection

Crystal:	colorless block, size 0.4 x 0.5 x 0.75 mm
Wavelength:	Mo <i>K</i> _α radiation (0.71073 Å)
μ:	1.11 cm ⁻¹
Diffractometer:	Nicolet P3
Scan mode:	Wyckoff
<i>T</i> _{measurement} :	293 K
2θ _{max} :	60°
<i>N</i> (<i>hkl</i>) _{unique} :	2226
Criterion for <i>I</i> _o :	<i>I</i> _o > 2 σ(<i>I</i> _o)
<i>N</i> (<i>param</i>) _{refined} :	181
Programs:	SHELXS-86, SHELXL-93

Table 2. Final atomic coordinates and displacement parameters (in Å²)

Atom	Site	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> _{iso}
H(1)	4 <i>a</i>	-0.1310(3)	0.9820(2)	0.86685(8)	0.036
H(3)	4 <i>a</i>	-0.2503(3)	0.8097(3)	0.9328(1)	0.050
H(31A)	4 <i>a</i>	-0.0322(5)	0.7413(4)	1.0086(1)	0.095
H(31B)	4 <i>a</i>	0.0390(5)	0.6000(4)	0.9740(1)	0.095
H(31C)	4 <i>a</i>	-0.1866(5)	0.6110(4)	0.9964(1)	0.095
H(5A)	4 <i>a</i>	-0.3983(3)	0.7546(3)	0.8437(1)	0.053
H(5B)	4 <i>a</i>	-0.3076(3)	0.6241(3)	0.8064(1)	0.053
H(6)	4 <i>a</i>	-0.1930(3)	0.8709(2)	0.77980(9)	0.042
H(8)	4 <i>a</i>	0.1165(4)	0.8948(3)	0.72995(9)	0.049
H(81A)	4 <i>a</i>	0.4342(5)	0.7955(4)	0.7045(1)	0.090
H(81B)	4 <i>a</i>	0.2625(5)	0.6786(4)	0.6905(1)	0.090
H(81C)	4 <i>a</i>	0.3981(5)	0.6590(4)	0.7458(1)	0.090
H(10)	4 <i>a</i>	0.1071(3)	1.0683(2)	0.80348(8)	0.036
H(12)	4 <i>a</i>	0.0695(4)	1.1514(3)	0.9385(1)	0.061
H(14)	4 <i>a</i>	0.2314(7)	1.5140(4)	0.8653(2)	0.101
H(15)	4 <i>a</i>	0.4939(6)	1.4641(3)	0.9313(2)	0.087
H(16)	4 <i>a</i>	0.3355(4)	1.2485(3)	0.9886(1)	0.069

Table 3. Final atomic coordinates and displacement parameters (in Å²)

Atom	Site	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> ₁₁	<i>U</i> ₂₂	<i>U</i> ₃₃	<i>U</i> ₁₂	<i>U</i> ₁₃	<i>U</i> ₂₃
C(1)	4 <i>a</i>	-0.0283(3)	0.9086(2)	0.85465(8)	0.0283(8)	0.0276(8)	0.0339(8)	0.0033(8)	-0.0018(7)	0.0007(7)
O(2)	4 <i>a</i>	0.0341(2)	0.8241(2)	0.90368(6)	0.0297(6)	0.0393(7)	0.0339(7)	-0.0017(6)	-0.0026(6)	0.0089(6)
C(3)	4 <i>a</i>	-0.1359(3)	0.7415(3)	0.9255(1)	0.033(1)	0.047(1)	0.046(1)	-0.001(1)	0.0028(9)	0.013(1)
C(31)	4 <i>a</i>	-0.0732(5)	0.6666(4)	0.9812(1)	0.052(2)	0.079(2)	0.060(2)	-0.003(2)	0.000(1)	0.037(2)
O(4)	4 <i>a</i>	-0.1957(3)	0.6319(2)	0.88547(8)	0.0417(8)	0.0376(8)	0.066(1)	-0.0076(7)	-0.0035(8)	0.0122(8)
C(5)	4 <i>a</i>	-0.2737(3)	0.7009(3)	0.8344(1)	0.0304(9)	0.043(1)	0.060(1)	-0.0037(9)	-0.008(1)	-0.004(1)
C(6)	4 <i>a</i>	-0.1220(3)	0.8080(2)	0.80806(9)	0.0329(9)	0.0336(9)	0.0386(9)	0.0003(9)	-0.0083(8)	-0.0026(8)
O(7)	4 <i>a</i>	0.0341(3)	0.7242(2)	0.77893(7)	0.0408(8)	0.0371(7)	0.0419(7)	-0.0036(7)	-0.0003(7)	-0.0106(6)
C(8)	4 <i>a</i>	0.1827(4)	0.8210(3)	0.75503(9)	0.046(1)	0.044(1)	0.0314(9)	-0.004(1)	0.0005(9)	-0.0082(8)

Table 3. (Continued)

Atom	Site	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> ₁₁	<i>U</i> ₂₂	<i>U</i> ₃₃	<i>U</i> ₁₂	<i>U</i> ₁₃	<i>U</i> ₂₃
C(81)	4a	0.3329(5)	0.7304(4)	0.7209(1)	0.056(2)	0.069(2)	0.055(1)	−0.010(1)	0.012(1)	−0.033(1)
O(9)	4a	0.2911(2)	0.8949(2)	0.79984(6)	0.0345(7)	0.0383(7)	0.0344(7)	−0.0020(7)	0.0037(6)	−0.0109(6)
C(10)	4a	0.1570(3)	0.9923(2)	0.83054(8)	0.0332(9)	0.0280(8)	0.0287(8)	−0.0001(7)	0.0006(7)	−0.0018(7)
C(11)	4a	0.2731(3)	1.0702(2)	0.87725(8)	0.0366(9)	0.0287(8)	0.0293(8)	−0.0050(8)	0.0042(8)	−0.0010(7)
C(12)	4a	0.1766(4)	1.1909(3)	0.9131(1)	0.045(1)	0.054(1)	0.054(1)	−0.008(1)	0.011(1)	−0.027(1)
O(13)	4a	0.1040(4)	1.3125(3)	0.8781(1)	0.078(2)	0.055(1)	0.123(2)	0.026(1)	−0.034(2)	−0.041(1)
C(14)	4a	0.2413(7)	1.4241(4)	0.8853(2)	0.095(3)	0.048(2)	0.109(3)	0.017(2)	−0.001(3)	−0.023(2)
C(15)	4a	0.3890(6)	1.3980(3)	0.9221(2)	0.070(2)	0.049(1)	0.099(2)	−0.017(2)	0.004(2)	−0.027(2)
C(16)	4a	0.3620(4)	1.2453(3)	0.9470(1)	0.061(2)	0.065(2)	0.047(1)	−0.019(2)	0.001(1)	−0.024(1)
O(17)	4a	0.5268(3)	1.1426(2)	0.93263(8)	0.054(1)	0.057(1)	0.0514(9)	−0.0115(9)	−0.0170(9)	−0.0053(8)
N(18)	4a	0.4607(3)	1.0469(2)	0.88762(8)	0.0418(9)	0.0371(9)	0.0407(9)	−0.0045(8)	−0.0059(8)	0.0005(8)

Acknowledgments. We are grateful to the Fonds der Chemischen Industrie, Bayer AG and Hoechst AG for financial support. We also thank Ms. S. Bucovarova for experimental assistance and Dr. Wolfgang Frey for help with the preparation of the files.

References

- Leibold, T.: Furan-Nitriloxid-Cycloaddukte - diastereomerenreine Vorläufer für die Synthese von biologisch aktiven Desoxynojirimycin- und Castanospermin-Analoga. Dissertation, Universität Stuttgart, Germany 1995.
- Müller, I.; Jäger, V.: Synthesis of amino sugars via isoxazolines. Nitrile oxide-furan adducts as key intermediates in the isoxazoline route towards novel amino sugar derivatives. *Tetrahedron* **41** (1985) 3519-3528.
- Jäger, V.; Müller, R.; Leibold, T.; Hein, M.; Schwarz, M.; Fengler, M.; Jaroskova, L.; Pätz, M.; LeRoy, P.-Y.: Synthesis of glycosidase-inhibiting iminopolyols via isoxazolines. *Bull. Soc. Chim. Belg.* **103** (1994) 491-507.
- Liu, K.-C.; Shelton, B. R.; Howe, R. K.: A particularly convenient preparation of benzohydroximinoyl chlorides (nitrile oxide precursors). *J. Org. Chem.* **45** (1980) 3916-3918.
- Henkel, S.; Leibold, T.; Jäger, V.: Crystal structure of 1,3:2,4-di-*O*-ethylidene-L-xylose oxime, C₉H₁₅NO₅. *Z. Kristallogr. NCS* **213** (1998) 65-66.
- Henkel, S.; Leibold, T.; Jäger, V.: Crystal structure of (3a*R*,6a*R*)-3-[(1*R*,2*S*,3*S*)-1,3:2,4-di-*O*-ethylidene-1,2,3,4-tetrahydroxy-1-butyl]-3a,6a-dihydrofuro[2,3-*d*]isoxazole, C₁₃H₁₇NO₆. *Z. Kristallogr. NCS* **213** (1998) 67-68.
- Sheldrick, G. M.: Phase Annealing in SHELX-90: Direct Methods for Large Structures. *Acta Crystallogr. A* **46** (1990) 467-473.
- Sheldrick, G. M.: SHELXL-93. Program for refining crystal structures. University of Göttingen, Germany 1993.