Crystal structure of 2*RS*,5-diphenyl-8*RS*-3,5,7-triazatricyclo[6.5.1.0^{3,7}]-tetradeca-9,11,13-triene-4,6-dione, C₂₃H₁₉N₃O₂

G. E. Gream, P. K. Kirkbride and E. R. T. Tiekink*,¹

The University of Adelaide, Department of Chemistry, Australia 5005

Received April 20, 2002; accepted and available on-line July 16, 2002; CCDC-No. 1267/851

drance. The molecule comprises a sequence of fused 5-, 6- and 8-membered rings. The sequence of $d(\text{C}—\text{C})$ bond distances for C8, C9 through to C1 are 1.483(4) Å, 1.325(4) Å, 1.446(4) Å, 1.323(4) Å, 1.441(3) Å and 1.322(3) Å, respectively. The torsion angles C8–C9–C10–C11 (6.8(5)°, C9–C10–C11–C12 (45.7(7)°), C10–C11–C12–C13 (–5.4(4)°) and C1–C13–C12–C11 (–48.1(4)°) reveal significant buckling within the ring. The patterns of parameters just described are indicative of a lack of conjugation within the 8-membered ring.

Table 1. Data collection and handling.

Crystal:	colourless plate, size 0.08 × 0.40 × 0.40 mm
Wavelength:	Mo K α radiation (0.71069 Å)
μ :	0.90 cm ^{–1}
Diffractometer, scan mode:	Rigaku AFC6R, $\omega/2\theta$
$2\theta_{\text{max}}$:	50.6°
$N(hkl)_{\text{measured}}$, $N(hkl)_{\text{unique}}$:	3448, 3261
Criterion for I_{obs} , $N(hkl)_{\text{gt}}$:	$I_{\text{obs}} > 2\sigma(I_{\text{obs}})$, 2315
$N(\text{param})_{\text{refined}}$:	253
Programs:	teXsan [2], SHELXS-86 [3], SHELXL-97 [4], ORTEP [5]

Abstract

C₂₃H₁₉N₃O₂, monoclinic, $P12_1/n1$ (No. 14), $a = 10.247(2)$ Å, $b = 16.465(2)$ Å, $c = 10.572(4)$ Å, $\beta = 92.22(2)^\circ$, $V = 1782.3$ Å³, $Z = 4$, $R_{\text{gt}}(F) = 0.043$, $wR_{\text{ref}}(F^2) = 0.155$, $T = 173$ K.

Source of material

Treatment of a mixture of (*E*)- and (*Z*)-7-benzylidenecycloocta-1,3,5-triene in ethyl acetate with 4-phenyl-1,2,4-triazoline-3,5-dione gave a mixture of four compounds [1]. Chromatography on silica gel gave the title compound, mp 442 K – 443 K.

Experimental details

The C-bound H atoms were placed in their geometrically calculated positions and included in the final refinement in the riding model approximation.

Discussion

The central C₂N₃ plane is planar to 0.042 Å with the greatest deviation being 0.067(2) Å for the C6 atom. The N5-phenyl group is not coplanar as evidenced by the C4–N5–C51–C52 torsion angle of 35(2)°. Presumably this occurs to as to minimise steric hin-

Table 2. Atomic coordinates and displacement parameters (in Å²).

Atom	Site	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> _{iso}
H(2)	4e	0.1111	0.2154	–0.1627	0.032
H(8)	4e	0.4330	0.1629	0.1569	0.038
H(9)	4e	0.5568	0.2350	0.0019	0.043
H(10)	4e	0.6282	0.1611	–0.1398	0.051
H(11)	4e	0.5658	0.0298	–0.1873	0.050
H(12)	4e	0.3728	0.0028	–0.2683	0.047
H(13)	4e	0.2137	0.1132	–0.2740	0.038
H(14a)	4e	0.3597	0.0565	0.0176	0.037
H(14b)	4e	0.2441	0.0958	0.0957	0.037
H(22)	4e	–0.0098	0.0742	–0.0799	0.042
H(23)	4e	–0.1932	0.0453	0.0325	0.049
H(24)	4e	–0.2633	0.1356	0.1843	0.049
H(25)	4e	–0.1460	0.2531	0.2253	0.047
H(26)	4e	0.0383	0.2827	0.1127	0.040
H(52)	4e	0.1023	0.4871	0.0472	0.040
H(53)	4e	0.1146	0.6251	0.0858	0.049
H(54)	4e	0.3109	0.6834	0.1536	0.055
H(55)	4e	0.4950	0.6038	0.1802	0.062
H(56)	4e	0.4871	0.4656	0.1353	0.054

* Correspondence author (e-mail: chmtert@nus.edu.sg)

¹ Current address: National University of Singapore, Department of Chemistry, Singapore 117543

Table 3. Atomic coordinates and displacement parameters (in Å²).

Atom	Site	x	y	z	U ₁₁	U ₂₂	U ₃₃	U ₁₂	U ₁₃	U ₂₃
O(4)	4e	0.1284(2)	0.37786(9)	-0.1052(2)	0.0317(9)	0.0297(8)	0.0322(9)	0.0003(7)	-0.0032(7)	0.0035(7)
O(6)	4e	0.4445(2)	0.3287(1)	0.2021(2)	0.050(1)	0.047(1)	0.047(1)	0.0062(9)	-0.0235(9)	-0.0101(9)
N(3)	4e	0.2302(2)	0.2642(1)	-0.0281(2)	0.029(1)	0.025(1)	0.0252(9)	0.0003(8)	-0.0053(8)	0.0008(7)
N(5)	4e	0.2859(2)	0.3794(1)	0.0597(2)	0.029(1)	0.025(1)	0.031(1)	0.0001(8)	-0.0043(8)	-0.0026(8)
N(7)	4e	0.3127(2)	0.2476(1)	0.0774(2)	0.030(1)	0.029(1)	0.0276(9)	0.0030(8)	-0.0060(8)	-0.0008(8)
C(1)	4e	0.2387(2)	0.1296(1)	-0.0908(2)	0.028(1)	0.024(1)	0.032(1)	-0.0021(9)	0.005(1)	-0.0002(9)
C(2)	4e	0.1460(2)	0.1995(1)	-0.0766(2)	0.027(1)	0.026(1)	0.026(1)	-0.0029(9)	0.0000(9)	-0.0014(9)
C(4)	4e	0.2051(2)	0.3442(1)	-0.0337(2)	0.024(1)	0.026(1)	0.027(1)	-0.0017(9)	0.0024(9)	0.0016(9)
C(6)	4e	0.3587(2)	0.3197(2)	0.1227(2)	0.035(1)	0.035(1)	0.031(1)	0.002(1)	-0.002(1)	-0.003(1)
C(8)	4e	0.3965(2)	0.1751(1)	0.0699(2)	0.033(1)	0.032(1)	0.030(1)	0.007(1)	-0.000(1)	0.003(1)
C(9)	4e	0.5073(2)	0.1871(2)	-0.0142(2)	0.028(1)	0.037(1)	0.042(1)	0.003(1)	-0.003(1)	0.004(1)
C(10)	4e	0.5483(3)	0.1418(2)	-0.1083(3)	0.032(1)	0.053(2)	0.044(2)	0.006(1)	0.006(1)	0.008(1)
C(11)	4e	0.5013(3)	0.0701(2)	-0.1747(2)	0.046(2)	0.042(2)	0.037(1)	0.016(1)	0.012(1)	0.003(1)
C(12)	4e	0.3831(3)	0.0515(2)	-0.2206(2)	0.050(2)	0.034(1)	0.035(1)	0.005(1)	0.011(1)	-0.002(1)
C(13)	4e	0.2692(2)	0.1009(1)	-0.2026(2)	0.035(1)	0.031(1)	0.030(1)	-0.004(1)	0.005(1)	-0.003(1)
C(14)	4e	0.3080(2)	0.1065(1)	0.0297(2)	0.036(1)	0.028(1)	0.030(1)	0.003(1)	0.007(1)	0.0029(9)
C(21)	4e	0.0322(2)	0.1816(1)	0.0050(2)	0.025(1)	0.030(1)	0.025(1)	0.0009(9)	-0.0003(9)	0.0014(9)
C(22)	4e	-0.0375(2)	0.1113(2)	-0.0175(2)	0.034(1)	0.035(1)	0.037(1)	-0.003(1)	0.006(1)	-0.001(1)
C(23)	4e	-0.1461(3)	0.0941(2)	0.0489(3)	0.034(1)	0.042(2)	0.048(2)	-0.009(1)	0.005(1)	0.001(1)
C(24)	4e	-0.1873(2)	0.1471(2)	0.1388(2)	0.029(1)	0.052(2)	0.041(1)	0.002(1)	0.006(1)	0.009(1)
C(25)	4e	-0.1185(3)	0.2164(2)	0.1623(2)	0.038(1)	0.044(2)	0.037(1)	0.011(1)	0.010(1)	0.001(1)
C(26)	4e	-0.0089(2)	0.2340(1)	0.0956(2)	0.035(1)	0.030(1)	0.035(1)	0.001(1)	0.007(1)	0.000(1)
C(51)	4e	0.2931(2)	0.4637(1)	0.0856(2)	0.032(1)	0.028(1)	0.030(1)	-0.004(1)	0.005(1)	-0.003(1)
C(52)	4e	0.1834(2)	0.5110(2)	0.0728(2)	0.031(1)	0.035(1)	0.034(1)	-0.001(1)	0.006(1)	-0.002(1)
C(53)	4e	0.1905(3)	0.5923(2)	0.0968(3)	0.047(2)	0.032(1)	0.044(2)	0.004(1)	0.006(1)	-0.002(1)
C(54)	4e	0.3063(3)	0.6268(2)	0.1365(3)	0.059(2)	0.030(1)	0.050(2)	-0.006(1)	0.012(1)	-0.008(1)
C(55)	4e	0.4150(3)	0.5797(2)	0.1515(3)	0.048(2)	0.044(2)	0.065(2)	-0.016(1)	0.002(1)	-0.015(1)
C(56)	4e	0.4106(3)	0.4980(2)	0.1257(3)	0.036(1)	0.042(2)	0.058(2)	-0.005(1)	-0.000(1)	-0.012(1)

Acknowledgment. The Australian Research Council is thanked for support.

References

1. Kirkbride, P. K.: Some aspects of the chemistry of cyclooctatetraene and its derivatives. Ph.D. Thesis. The University of Adelaide, Australia 1983.
2. teXsan: Single Crystal Structure Analysis Software. Version 1.04. Molecular Structure Corporation. The Woodlands, TX, USA 1997.
3. Sheldrick, G. M.: SHELX-86. Program for the automatic solution of crystal structures. University of Göttingen, Germany 1986.
4. Sheldrick, G. M.: SHELXL-97. Program for crystal structure refinement. University of Göttingen, Germany, 1997.
5. Johnson, C. K.: ORTEPII. Report ORNL-5138, Oak Ridge National Laboratory, Tennessee, USA 1976.