Crystal structure of dimethyl (1 α ,2 β ,3 α ,4 β ,7 β ,8 α ,9 β ,10 α)-13-isopropylidene-pentacyclo[8.2.1.1^{4,7}.0^{2,9}.0^{3,8}]tetradeca-5,11-diene-2,9-dicarboxylate, C₂₁H₂₄O₄

R. A. Russell^I, A. C. Schultz^{II}, R. N. Warrener^{II} and E. R. T. Tiekink^{*,III,1}^I Deakin University, Centre for Chiral and Molecular Technologies, Geelong, Victoria 3217, Australia^{II} Central Queensland University, Centre for Molecular Architecture, Rockhampton, Queensland 4702, Australia^{III} The University of Adelaide, Department of Chemistry, Australia 5005

Received January 21, 2002, accepted and available on-line April 30, 2002; CCDC-No. 1267/790

Abstract

C₂₁H₂₄O₄, monoclinic, *P*12₁/*n*1 (No. 14), *a* = 9.547(2) Å, *b* = 19.014(3) Å, *c* = 9.877(2) Å, β = 91.42(2)°, *V* = 1792.4 Å³, *Z* = 4, *R*_{gt}(*F*) = 0.062, *wR*_{ref}(*F*²) = 0.194, *T* = 293 K.

Source of material

LiClO₄ (11 mg, 5 mol%) was added to a solution of Smith's diene [(1 α ,2 β ,5 β ,6 α)dimethyl tricyclo[4.2.1.0^{2,5}]nona-3,7-diene-3,4-dicarboxylate] (500 mg, 2.14 mmol) and 6,6-dimethylfulvene (1.1 g, 10.4 mmol) in dichloromethane (3 ml). The solution was stirred at room temperature for two days under nitrogen. Further dichloromethane was added (10 ml) and the solution washed with water. The organic fraction was dried (Na₂SO₄) and the solvent removed under reduced pressure. The resultant residue was separated by column chromatography; elution with petroleum spirit removed excess dimethylfulvene before the crude product was eluted with 1:1 EtOAc/petroleum spirit. Combined fractions were taken to dryness to give an oil which was triturated with MeOH to precipitate polymeric impurities. The MeOH solution

was taken to dryness to give adduct which was recrystallised from petroleum spirit; yield: 456 mg (64.5%); mp 378 K – 379 K. ¹H and ¹³C NMR as well as EIMS results are included in the deposited CIF-file.

Experimental details

The C-bound H atoms were placed in their geometrically calculated positions and included in the final refinement in the riding model approximation.

Discussion

The molecule adopts a staircase motif with successive risers being defined by C11–C12, C1–C13–C10, C2–C9, C3–C8, C4–C14–C7 and C5–C6 connectivities. The ester groups lie to the opposite side of the molecule to the C=CMe₂ group.

Table 1. Data collection and handling.

Crystal:	colourless, multifaceted, size 0.29 × 0.32 × 0.40 mm
Wavelength:	Mo <i>K</i> α radiation (0.7107 Å)
μ :	0.86 cm ⁻¹
Diffractometer, scan mode:	Rigaku AFC6R, $\omega/2\theta$
2 θ _{max} :	50.2°
<i>N</i> (<i>hkl</i>) _{measured} , <i>N</i> (<i>hkl</i>) _{unique} :	3370, 3175
Criterion for <i>I</i> _{obs} , <i>N</i> (<i>hkl</i>) _{gt} :	<i>I</i> _{obs} > 2 σ (<i>I</i> _{obs}), 2196
<i>N</i> (<i>param</i>) _{refined} :	227
Programs:	SHELXS-86 [1], SHELXL-97 [2], teXsan [3], DIBABS [4]

Table 2. Atomic coordinates and displacement parameters (in Å²).

Atom	Site	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> _{iso}
H(1)	4e	−0.1983	0.2420	−0.2055	0.091(3)
H(2a)	4e	0.0739	0.4138	−0.2227	0.109(4)
H(2b)	4e	0.1930	0.3721	−0.2939	0.109
H(2c)	4e	0.1829	0.3704	−0.1358	0.109
H(3)	4e	−0.1232	0.1875	−0.4430	0.091
H(4)	4e	0.1057	0.2488	−0.5248	0.091
H(5)	4e	−0.0126	0.1840	−0.7191	0.073(2)
H(6)	4e	0.0263	0.0583	−0.6903	0.073
H(7)	4e	0.1728	0.0380	−0.4775	0.091
H(8)	4e	−0.0815	0.0679	−0.4092	0.091
H(9a)	4e	0.3542	0.1214	0.0786	0.109
H(9b)	4e	0.4121	0.0997	−0.0626	0.109

* Correspondence author (e-mail: chmert@nus.edu.sg)

¹ Current address: National University of Singapore, Department of Chemistry, Singapore 117543

Table 2. Continued.

Atom	Site	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> _{iso}
H(9c)	4e	0.3310	0.0447	0.0240	0.109
H(10)	4e	−0.1131	0.0345	−0.1467	0.091
H(11)	4e	−0.0578	0.1006	0.0683	0.073
H(12)	4e	−0.1087	0.2238	0.0336	0.073
H(13a)	4e	−0.4670	0.0390	−0.3486	0.109
H(13b)	4e	−0.3870	0.0155	−0.2156	0.109

Table 2. Continued.

Atom	Site	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> _{iso}
H(13c)	4e	−0.3103	0.0148	−0.3542	0.109
H(13d)	4e	−0.5089	0.1469	−0.3746	0.109
H(13e)	4e	−0.3812	0.1968	−0.3983	0.109
H(13f)	4e	−0.4582	0.1987	−0.2600	0.109
H(14a)	4e	0.2820	0.1516	−0.5456	0.091
H(14b)	4e	0.2459	0.1580	−0.3897	0.091

Table 3. Atomic coordinates and displacement parameters (in Å²).

Atom	Site	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> ₁₁	<i>U</i> ₂₂	<i>U</i> ₃₃	<i>U</i> ₁₂	<i>U</i> ₁₃	<i>U</i> ₂₃
O(2)	4e	0.2278(2)	0.2427(1)	−0.2101(3)	0.036(1)	0.050(1)	0.100(2)	−0.005(1)	−0.011(1)	−0.013(1)
O(2a)	4e	0.0401(2)	0.3109(1)	−0.2297(3)	0.047(1)	0.030(1)	0.080(2)	−0.002(1)	−0.000(1)	−0.005(1)
O(9)	4e	0.2287(3)	0.0320(2)	−0.2207(3)	0.072(2)	0.075(2)	0.069(2)	0.041(2)	−0.014(1)	−0.012(1)
O(9a)	4e	0.2076(3)	0.1149(1)	−0.0653(2)	0.061(2)	0.075(2)	0.052(2)	0.016(1)	−0.024(1)	−0.011(1)
C(1)	4e	−0.1432(3)	0.2001(2)	−0.1820(3)	0.034(2)	0.036(2)	0.036(2)	0.004(1)	0.005(1)	−0.004(1)
C(2)	4e	−0.0005(3)	0.1911(1)	−0.2570(3)	0.031(1)	0.031(2)	0.034(2)	0.003(1)	0.001(1)	−0.004(1)
C(2a)	4e	0.1039(3)	0.2485(2)	−0.2270(3)	0.040(2)	0.035(2)	0.035(2)	−0.001(1)	0.001(1)	−0.005(1)
C(2b)	4e	0.1300(4)	0.3719(2)	−0.2197(5)	0.069(2)	0.035(2)	0.100(3)	−0.012(2)	0.007(2)	−0.008(2)
C(3)	4e	−0.0270(3)	0.1772(1)	−0.4124(3)	0.036(2)	0.029(1)	0.029(1)	−0.000(1)	0.002(1)	−0.001(1)
C(4)	4e	0.0846(3)	0.1985(2)	−0.5172(3)	0.040(2)	0.034(2)	0.037(2)	−0.003(1)	0.006(1)	0.004(1)
C(5)	4e	0.0290(3)	0.1623(2)	−0.6439(3)	0.052(2)	0.048(2)	0.028(2)	0.001(2)	0.006(1)	0.004(1)
C(6)	4e	0.0505(3)	0.0932(2)	−0.6282(3)	0.055(2)	0.045(2)	0.030(2)	0.002(2)	0.006(1)	−0.005(1)
C(7)	4e	0.1216(3)	0.0822(2)	−0.4906(3)	0.043(2)	0.033(2)	0.036(2)	0.004(1)	0.006(1)	−0.001(1)
C(8)	4e	0.0015(3)	0.0971(1)	−0.3916(3)	0.037(2)	0.028(1)	0.030(1)	−0.001(1)	0.001(1)	−0.000(1)
C(9)	4e	0.0320(3)	0.1100(2)	−0.2372(3)	0.034(1)	0.032(2)	0.030(2)	0.004(1)	0.001(1)	−0.000(1)
C(9a)	4e	0.1674(3)	0.0819(2)	−0.1781(3)	0.041(2)	0.042(2)	0.035(2)	0.004(1)	0.002(1)	0.004(1)
C(9b)	4e	0.3369(5)	0.0934(3)	−0.0010(4)	0.064(3)	0.122(4)	0.061(3)	0.012(3)	−0.027(2)	0.007(3)
C(10)	4e	−0.0958(3)	0.0852(2)	−0.1494(3)	0.041(2)	0.037(2)	0.033(2)	0.001(1)	0.005(1)	0.003(1)
C(11)	4e	−0.0818(3)	0.1218(2)	−0.0139(3)	0.050(2)	0.058(2)	0.030(2)	0.004(2)	0.007(1)	0.001(1)
C(12)	4e	−0.1096(3)	0.1892(2)	−0.0329(3)	0.044(2)	0.054(2)	0.034(2)	0.000(2)	0.006(1)	−0.009(1)
C(13)	4e	−0.2116(3)	0.1298(2)	−0.2133(3)	0.036(2)	0.036(2)	0.032(1)	−0.000(1)	0.008(1)	0.000(1)
C(13a)	4e	−0.3308(3)	0.1137(2)	−0.2773(3)	0.037(2)	0.046(2)	0.040(2)	−0.004(1)	0.008(1)	−0.004(1)
C(13b)	4e	−0.3781(4)	0.0390(2)	−0.3011(4)	0.053(2)	0.057(2)	0.067(2)	−0.016(2)	0.004(2)	−0.006(2)
C(13c)	4e	−0.4286(3)	0.1690(2)	−0.3325(4)	0.039(2)	0.064(2)	0.058(2)	0.005(2)	−0.006(2)	−0.005(2)
C(14)	4e	0.2079(3)	0.1497(2)	−0.4803(3)	0.039(2)	0.045(2)	0.040(2)	0.001(1)	0.008(1)	−0.001(1)

Acknowledgment. The Australian Research Council is thanked for support.

References

- Sheldrick, G. M.: SHELXS-86. Program for the solution of crystal structure. University of Göttingen, Germany 1986.
- Sheldrick, G. M.: SHELXL-97. Program for crystal structure refinement. University of Göttingen, Germany 1997.
- teXsan: Single Crystal Structure Analysis Software. Version 1.04. Molecular Structure Corporation. The Woodlands, TX, USA 1997.
- Walker, N.; Stuart, D.: An empirical method for correcting diffractometer data for absorption effects. *Acta Crystallogr. A* **39** (1983) 159–166.