Crystal structure of 4-(2-oxobenzothiazolin-3-yl)butanoic acid, $C_{11}H_{11}NO_3S$

F. Baysen^{*,I}, R. Yağbasan^I, E. Kendi^{II}, M. Aslantaş^{II}, B. Çakır^{III} and T. Önkol^{III}^I Gazi University, Gazi Education Faculty, Department of Physics Education, Beşevler 06500, Ankara, Turkey^{II} Hecettepe University, Department of Engineering Physics, Beytepe 06532, Ankara, Turkey^{III} Gazi University, Faculty of Pharmacy, Department of Pharmaceutical Chemistry, Hipodrom 06330, Ankara, Turkey

Received November 8, 2002, accepted in final form and available on-line April 15, 2003; CCDC-No. 1267/970

Abstract

$C_{11}H_{11}NO_3S$, triclinic, $P\bar{1}$ (No. 2), $a = 7.240(1)$ Å, $b = 7.495(2)$ Å, $c = 10.427(1)$ Å, $\alpha = 83.89(1)^\circ$, $\beta = 85.74(1)^\circ$, $\gamma = 71.16(1)^\circ$, $V = 532.0$ Å³, $Z = 2$, $R_{\text{gt}}(F) = 0.035$, $wR_{\text{ref}}(F^2) = 0.096$, $T = 293$ K.

Source of material

For the synthesis, 10.0 mmol ethyl 4-(2-oxobenzothiazolin-3-yl)-butanoate in concentrated hydrochloric acid (50 ml) was stirred at room temperature for 2 hours, then refluxed for 4 hours. The reaction mixture was cooled, poured into 100 g ice-water, and stirred for 1 hour. The precipitate was collected by filtration, washed with water, dried and crystallised from water.

Discussion

The benzene ring defined by C1–C2–C3–C4–C5–C6 atoms is planar. Torsion angles of C1–C2–C3–C4, C6–C5–C4–C3 and C1–C6–C5–C4 are $0.4(3)^\circ$, $0.1(3)^\circ$ and $0.3(3)^\circ$, respectively. The thiazolin ring defined by C1–C6–N–C7–S atoms is planar. Torsion angles of C7–S–C1–C6 and C6–N–C7–S are $0.6(1)^\circ$ and $1.2(2)^\circ$, respectively. The dihedral angle between these two planes is $0.99(8)^\circ$ implying that they are co-planar. The torsion angles of C7–N–C6–C5, N–C6–C1–C2, C7–S–C1–C2 and S2–C1–C2–C3 are $178.8(2)^\circ$, $179.2(2)^\circ$, $-178.6(2)^\circ$ and $179.1(2)^\circ$, respectively. O1 and C8 atoms are also at this plane. Torsion angles of C6–N–C7–O1, C1–S–C7–O1, C8–N–C6–C1 and C8–N–C7–O1 are $-179.0(2)^\circ$, $179.2(2)^\circ$, $-179.0(2)^\circ$ and $-0.7(3)^\circ$, respectively. C9 atom lies below $1.351(3)$ Å from thiazolin plane. Torsion angles of C7–N–C8–C9, C6–N–C8–C9 and N–C8–C9–C10 are $103.8(2)^\circ$, $-78.0(2)^\circ$ and $-68.5(2)^\circ$, re-

spectively. The bond lengths and angles in the 4-(2-oxobenzothiazolin-3-yl)butanoic acid are all in accord with similar structures in the literature [1–3]. The bond lengths of C–C are between $1.378(3)$ Å – $1.522(3)$ Å. The bond lengths of C7=O1, C11–O2 and C11=O3 are $1.214(2)$ Å, $1.262(2)$ Å and $1.263(2)$ Å, respectively. The bond lengths of C7–S and C1–S are $1.782(2)$ Å and $1.743(2)$ Å, respectively. The bond lengths of C6–N, C7–N and C8–N are $1.395(2)$ Å, $1.368(2)$ Å and $1.464(2)$ Å, respectively.

Table 1. Data collection and handling.

Crystal:	white needle, size $0.06 \times 0.27 \times 0.48$ mm
Wavelength:	Mo $K\alpha$ radiation (0.71073 Å)
μ :	2.94 cm^{-1}
Diffractometer, scan mode:	Enraf-Nonius CAD-4, $\omega/2\theta$
$2\theta_{\text{max}}$:	52.58°
$N(hkl)_{\text{measured}}$, $N(hkl)_{\text{unique}}$:	2300, 2130
Criterion for I_{obs} , $N(hkl)_{\text{gt}}$:	$I_{\text{obs}} > 2 \sigma(I_{\text{obs}})$, 1786
$N(\text{param})_{\text{refined}}$:	147
Programs:	SHELXS-97 [4] SHELXL-97 [5], ORTEP-III [6]

Table 2. Atomic coordinates and displacement parameters (in Å²).

Atom	Site	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> _{iso}
H(2)	2i	−0.4244	−0.0414	0.5929	0.070
H(8A)	2i	−0.1139	0.5419	0.6207	0.041
H(8B)	2i	−0.2521	0.5945	0.7436	0.041
H(5)	2i	−0.3345	0.4424	0.9429	0.046
H(4)	2i	−0.3352	0.3370	1.1597	0.053
H(9A)	2i	−0.4118	0.4711	0.6134	0.042
H(9B)	2i	−0.3442	0.3186	0.7301	0.042
H(10A)	2i	−0.1531	0.2997	0.4823	0.048
H(10B)	2i	−0.0717	0.1516	0.5994	0.048
H(2A)	2i	0.2404	0.0690	1.1354	0.048
H(3)	2i	−0.0519	0.1549	1.2552	0.054

* Correspondence author (e-mail: fatmac@gazi.edu.tr)

Table 3. Atomic coordinates and displacement parameters (in Å²).

Atom	Site	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> ₁₁	<i>U</i> ₂₂	<i>U</i> ₃₃	<i>U</i> ₁₂	<i>U</i> ₁₃	<i>U</i> ₂₃
S	2i	0.32516(6)	0.16590(7)	0.85628(5)	0.0272(2)	0.0507(3)	0.0421(3)	−0.0099(2)	−0.0009(2)	−0.0008(2)
O(2)	2i	−0.3456(2)	0.0043(2)	0.6171(1)	0.0589(9)	0.0509(9)	0.0424(8)	−0.0339(7)	−0.0150(6)	0.0080(6)
O(3)	2i	−0.3910(2)	0.1564(2)	0.4210(1)	0.0590(9)	0.0510(8)	0.0337(7)	−0.0311(7)	−0.0082(6)	0.0022(6)
O(1)	2i	0.2322(2)	0.3345(2)	0.6211(1)	0.0448(8)	0.069(1)	0.0399(8)	−0.0219(7)	0.0058(6)	0.0044(7)
N	2i	−0.0148(2)	0.3698(2)	0.7785(1)	0.0288(7)	0.0354(8)	0.0317(8)	−0.0138(6)	−0.0040(6)	−0.0008(6)
C(6)	2i	−0.0443(2)	0.3184(2)	0.9091(2)	0.0309(8)	0.0297(8)	0.0310(8)	−0.0143(7)	−0.0032(7)	−0.0042(7)
C(1)	2i	0.1269(2)	0.2062(2)	0.9673(2)	0.0297(8)	0.0310(9)	0.0354(9)	−0.0108(7)	−0.0014(7)	−0.0053(7)
C(11)	2i	−0.3165(3)	0.1242(3)	0.5303(2)	0.0350(9)	0.0347(9)	0.0354(9)	−0.0138(7)	0.0006(7)	−0.0047(7)
C(8)	2i	−0.1711(3)	0.4898(2)	0.6966(2)	0.0362(9)	0.0313(9)	0.0382(9)	−0.0133(7)	−0.0091(7)	0.0015(7)
C(5)	2i	−0.2193(3)	0.3681(3)	0.9809(2)	0.0314(9)	0.043(1)	0.039(1)	−0.0094(8)	0.0010(7)	−0.0052(8)
C(4)	2i	−0.2187(3)	0.3048(3)	1.1104(2)	0.043(1)	0.049(1)	0.042(1)	−0.0148(9)	0.0107(8)	−0.0088(9)
C(9)	2i	−0.2984(2)	0.3820(3)	0.6545(2)	0.0300(9)	0.036(1)	0.041(1)	−0.0119(7)	−0.0069(7)	−0.0038(8)
C(10)	2i	−0.1901(3)	0.2362(3)	0.5611(2)	0.039(1)	0.048(1)	0.042(1)	−0.0243(9)	0.0025(8)	−0.0088(9)
C(7)	2i	0.1740(3)	0.3046(3)	0.7310(2)	0.0339(9)	0.040(1)	0.0361(9)	−0.0184(8)	−0.0007(7)	−0.0032(7)
C(2)	2i	0.1258(3)	0.1438(3)	1.0969(2)	0.042(1)	0.040(1)	0.037(1)	−0.0109(8)	−0.0083(8)	0.0006(8)
C(3)	2i	−0.0487(3)	0.1949(3)	1.1678(2)	0.058(1)	0.047(1)	0.0299(9)	−0.018(1)	0.0018(8)	−0.0003(8)

Acknowledgment. The authors acknowledge the purchase of the CAD4 diffractometer under Grant DPT/TBAG1 of the Scientific and Technical Research Council of Turkey.

References

1. Van der Poorten, K.; Ucar, H.; Andrei, G.; Snoeck, R.; Balzarini, J.; De Clercq, E.; Pourpaert, J. H.: Synthesis and Antiviral Activity of 6-Benzoyl-Benzoxazolin-2-One and 6-Benzoyl-Benzothiazolin-2-One Derivatives. *Antiviral Chem. Chemother.* **10** (1999) 87-97.
2. Gokce, M.; Çakır, B.; Erol, K.; Sahin, M. F.: Synthesis and Antinociceptive Activity of [(2-Oxobenzothiazolin-3-yl)methyl]-4-Alkyl/Aryl-1,2,4-Triazoline-5-Thiones. *Arch. Pharm.* **334** (2001) 279-283.
3. Carato, P.; Yous, S.; Depreux, P.: Synthesis of 6-Chloromethyl Benzothiazolin-2-One and of 6-Chloromethyl Benzoxazolin-2-One. *Org. Prep. Proced. Int.* **32** (2000) 69-74.
4. Sheldrick, G. M.: SHELXS-97. Program for the Solution of Crystal Structures. University of Göttingen, Germany 1997.
5. Sheldrick, G. M.: SHELXL-97. Program for the Refinement of Crystal Structures. University of Göttingen, Germany 1997.
6. Farrugia, L. J.: ORTEP-III for Windows. Glasgow University, Department of Chemistry, Glasgow G12 8QQ, Scotland, UK 1998.