Crystal structure of $(\eta^4\text{-norborna-2,5-dien})\text{-(1,2-bis((R,R)-2,5-bis-(methoxymethyl)phospholanyl)benzene)rhodium(I) tetrafluoroborate, [Rh(C₂₂H₃₆O₄P₂)(C₇H₈)](BF₄)$

J. Holz, A. Börner, A. Spannenberg, C. Pribbenow, D. Heller and H.-J. Drexler*

Leibniz-Institut für Organische Katalyse an der Universität Rostock e.V., Buchbinderstr. 5-6, D-18055 Rostock, Germany

Received March 9, 2004, accepted and available on-line May 5, 2004; CCDC no. 1267/1271

the title compound. The ratio of the rate constants for the hydrogenation of the diolefin complexes is approximately 750. It is well known that the double bonds of the diolefines are not coordinated perpendicular to the P,Rh,P plane. The dihedral angle between the planes P,Rh,P and X,Rh,X (X = centroid of the double bond) is in the case of the (R,R)-COD-complex 21.8° (clockwise twist) and for the (R,R)-NBD-complex 22.2° (clockwise twist).

Table 1. Data collection and handling.

Crystal:	red prism, size 0.30 × 0.30 × 0.40 mm
Wavelength:	Mo K α radiation (0.71073 Å)
μ :	6.98 cm ⁻¹
Diffractometer, scan mode:	Stoe IPDS I, φ
2 θ_{max} :	45°
$N(hkl)_{\text{measured}}$, $N(hkl)_{\text{unique}}$:	7171, 4083
Criterion for I_{obs} , $N(hkl)_{\text{gt}}$:	$I_{\text{obs}} > 2 \sigma(I_{\text{obs}})$, 3876
$N(\text{param})_{\text{refined}}$:	361
Programs:	SHELXS-97 [4], SHELXL-97 [5]

Table 2. Atomic coordinates and displacement parameters (in Å²).

Atom	Site	Occ.	x	y	z	U_{iso}
H(1A)	2a		0.6959	0.0304	0.7967	0.08
H(2A)	2a		0.5513	-0.0512	0.6232	0.08
H(3A)	2a		0.4763	-0.2046	0.6878	0.08
H(4A)	2a		0.3333	-0.1648	0.8253	0.08
H(5A)	2a		0.4767	-0.0819	1.0022	0.08
H(6A)	2a		0.7170	-0.0678	0.9796	0.08
H(7A)	2a		0.7229	-0.1847	0.8194	0.08
H(7B)	2a		0.6364	-0.2277	0.9003	0.08
H(10A)	2a		0.0060	0.2031	0.8640	0.08
H(11A)	2a		-0.0192	0.3679	0.8389	0.08
H(12A)	2a		0.1401	0.4544	0.7722	0.08
H(13A)	2a		0.3290	0.3794	0.7341	0.08
H(14A)	2a		0.4774	0.1611	0.5665	0.08
H(15A)	2a		0.5300	0.3493	0.6318	0.08
H(15B)	2a		0.6093	0.2853	0.5642	0.08
H(16A)	2a		0.7345	0.3089	0.7764	0.08
H(16B)	2a		0.7118	0.2029	0.7389	0.08
H(17A)	2a		0.5522	0.3071	0.8539	0.08
H(18A)	2a		0.2854	0.3092	0.5334	0.08
H(18B)	2a		0.3295	0.2585	0.4307	0.08
H(19A)	2a		0.0118	0.1689	0.4387	0.08
H(19B)	2a		0.0609	0.2702	0.4885	0.08
H(19C)	2a		0.0735	0.2385	0.3620	0.08
H(20A)	2a		0.5959	0.2013	1.0101	0.08
H(20B)	2a		0.6748	0.1342	0.9460	0.08
H(21A)	2a		0.9488	0.2381	1.1498	0.08
H(21B)	2a		0.8167	0.1960	1.1754	0.08
H(21C)	2a		0.8894	0.1405	1.0934	0.08

Abstract

C₂₉H₄₄BF₄O₄P₂Rh, monoclinic, $P2_1$ (no. 4), $a = 10.119(2)$ Å, $b = 14.125(3)$ Å, $c = 11.542(2)$ Å, $\beta = 106.87(3)^\circ$, $V = 1578.7$ Å³, $Z = 2$, $R_{\text{gt}}(F) = 0.034$, $wR_{\text{ref}}(F^2) = 0.090$, $T = 200$ K.

Source of material

Standard preparation according to [1]. The synthesis of the ligand is described in [2].

Experimental details

The hydrogen atoms were placed in theoretical positions and refined using a riding model. The disorder of the BF₄⁻ anion was resolved in two tetrahedra, whereas all F and B atoms were split into two positions and treated as isotropic in the structure refinement resulting in an occupational ratio of 0.60(1)/0.40.

Discussion

Unexpected differences between the title compound and the related (Z,Z)-cycloocta-1,5-diene complex [2] in the catalytic hydrogenation of the diolefines norborna-2,5-diene and (Z,Z)-cycloocta-1,5-diene [3] motivated us to determine the crystal structure of

* Correspondence author

(e-mail: hans-joachim.drexler@ifok.uni-rostock.de)

Table 2. Continued.

Atom	Site	Occ.	x	y	z	U _{iso}
H(22A)	2a		−0.0136	0.0782	0.6767	0.08
H(23A)	2a		0.0074	−0.1050	0.7678	0.08
H(23B)	2a		−0.1257	−0.0432	0.7272	0.08
H(24A)	2a		−0.0273	−0.0536	0.9461	0.08
H(24B)	2a		−0.0478	0.0502	0.8977	0.08
H(25A)	2a		0.2009	−0.0508	0.9583	0.08
H(26A)	2a		−0.0214	−0.0272	0.5262	0.08
H(26B)	2a		0.1204	0.0245	0.5541	0.08
H(27A)	2a		0.2302	−0.1991	0.5455	0.08
H(27B)	2a		0.1076	−0.1488	0.4505	0.08
H(27C)	2a		0.2479	−0.0953	0.5041	0.08
H(28A)	2a		0.3044	0.0740	1.0833	0.08
H(28B)	2a		0.1595	0.1214	1.0547	0.08

Table 2. Continued.

Atom	Site	Occ.	x	y	z	U _{iso}
H(29A)	2a		0.2039	0.0110	1.3270	0.08
H(29B)	2a		0.3255	0.0578	1.2890	0.08
H(29C)	2a		0.1857	0.1122	1.2688	0.08
B(1)	2a	0.6	0.6775(7)	0.0762(5)	0.4334(6)	0.025(2)
F(1)	2a	0.6	0.584(1)	0.1444(6)	0.3742(9)	0.101(2)
F(2)	2a	0.6	0.6058(9)	−0.0063(6)	0.4336(9)	0.116(3)
F(3)	2a	0.6	0.7754(9)	0.0624(6)	0.3740(8)	0.110(3)
F(4)	2a	0.6	0.740(1)	0.1063(7)	0.5495(8)	0.123(3)
B(1')	2a	0.4	0.647(1)	0.0655(7)	0.3710(9)	0.035(4)
F(1')	2a	0.4	0.574(1)	0.138(1)	0.405(1)	0.101(2)
F(2')	2a	0.4	0.672(2)	−0.0060(9)	0.457(1)	0.116(3)
F(3')	2a	0.4	0.772(1)	0.099(1)	0.361(1)	0.110(3)
F(4')	2a	0.4	0.571(1)	0.031(1)	0.260(1)	0.123(3)

Table 3. Atomic coordinates and displacement parameters (in Å²).

Atom	Site	x	y	z	U ₁₁	U ₂₂	U ₃₃	U ₁₂	U ₁₃	U ₂₃
Rh(1)	2a	0.40941(3)	0.02366(3)	0.78636(3)	0.0198(2)	0.0257(2)	0.0298(2)	0.0033(2)	0.0123(1)	0.0054(2)
P(1)	2a	0.4244(1)	0.1800(1)	0.7476(1)	0.0198(7)	0.0253(7)	0.0275(8)	−0.0019(6)	0.0105(6)	0.0008(7)
P(2)	2a	0.2056(1)	0.0633(1)	0.8162(1)	0.0190(7)	0.0267(6)	0.0327(8)	−0.0004(5)	0.0108(6)	0.0037(5)
O(1)	2a	0.2119(4)	0.1800(3)	0.4994(4)	0.037(2)	0.041(2)	0.045(3)	−0.005(2)	0.002(2)	0.005(2)
O(2)	2a	0.7789(5)	0.2490(4)	1.0104(5)	0.046(3)	0.061(3)	0.070(3)	−0.007(2)	−0.008(3)	−0.014(3)
O(3)	2a	0.1363(5)	−0.1066(3)	0.6161(4)	0.054(3)	0.055(3)	0.045(3)	0.015(2)	0.022(2)	−0.008(2)
O(4)	2a	0.1760(5)	0.0099(4)	1.1531(4)	0.094(3)	0.077(4)	0.040(3)	−0.042(3)	0.024(2)	−0.002(3)
C(1)	2a	0.6276(5)	−0.0152(4)	0.8020(6)	0.020(3)	0.040(3)	0.050(4)	0.014(2)	0.016(3)	0.005(3)
C(2)	2a	0.5464(6)	−0.0617(4)	0.7041(6)	0.028(3)	0.049(4)	0.041(4)	0.011(3)	0.021(3)	−0.004(3)
C(3)	2a	0.5080(7)	−0.1565(4)	0.7483(7)	0.047(4)	0.029(3)	0.084(5)	−0.005(3)	0.028(4)	−0.011(3)
C(4)	2a	0.4144(7)	−0.1293(4)	0.8257(8)	0.054(4)	0.023(3)	0.083(5)	0.010(3)	0.039(4)	0.025(4)
C(5)	2a	0.4947(7)	−0.0813(5)	0.9250(6)	0.048(4)	0.062(4)	0.046(4)	0.025(3)	0.021(3)	0.028(4)
C(6)	2a	0.6417(6)	−0.0803(6)	0.9085(6)	0.032(4)	0.082(5)	0.042(4)	0.015(3)	0.012(3)	0.008(4)
C(7)	2a	0.6447(8)	−0.1756(5)	0.8494(9)	0.044(4)	0.048(5)	0.113(7)	0.025(4)	0.034(5)	0.033(5)
C(8)	2a	0.2807(5)	0.2447(3)	0.7768(4)	0.027(3)	0.025(3)	0.020(3)	0.001(2)	0.010(2)	−0.001(2)
C(9)	2a	0.1837(5)	0.1923(4)	0.8147(4)	0.016(3)	0.033(3)	0.023(3)	0.007(2)	0.003(2)	0.000(2)
C(10)	2a	0.0727(6)	0.2391(4)	0.8380(5)	0.028(3)	0.036(3)	0.042(3)	0.007(2)	0.012(3)	0.003(3)
C(11)	2a	0.0573(6)	0.3365(4)	0.8220(5)	0.027(3)	0.044(4)	0.037(4)	0.009(3)	−0.001(3)	−0.008(3)
C(12)	2a	0.1522(6)	0.3873(4)	0.7834(5)	0.035(3)	0.028(3)	0.037(3)	0.006(3)	0.006(3)	−0.002(2)
C(13)	2a	0.2640(6)	0.3429(4)	0.7617(5)	0.035(3)	0.027(3)	0.032(3)	0.005(2)	0.007(3)	0.000(2)
C(14)	2a	0.4467(6)	0.2160(4)	0.6002(5)	0.039(3)	0.040(3)	0.029(3)	−0.006(3)	0.016(3)	0.007(2)
C(15)	2a	0.5643(7)	0.2865(5)	0.6269(6)	0.051(4)	0.051(4)	0.057(4)	−0.021(3)	0.025(3)	0.007(3)
C(16)	2a	0.6659(6)	0.2607(5)	0.7479(7)	0.034(4)	0.052(4)	0.066(5)	−0.018(3)	0.022(3)	0.003(3)
C(17)	2a	0.5819(6)	0.2455(4)	0.8366(6)	0.022(3)	0.033(3)	0.038(3)	0.000(2)	0.001(3)	−0.002(2)
C(18)	2a	0.3141(6)	0.2498(4)	0.5083(6)	0.051(4)	0.039(3)	0.038(4)	0.004(3)	0.016(3)	0.016(3)
C(19)	2a	0.0790(8)	0.2177(7)	0.443(1)	0.051(5)	0.075(6)	0.128(8)	0.004(4)	−0.015(5)	0.030(6)
C(20)	2a	0.6543(6)	0.1992(4)	0.9579(6)	0.031(3)	0.041(4)	0.050(4)	0.001(3)	0.006(3)	−0.016(3)
C(21)	2a	0.8659(8)	0.2022(7)	1.1162(8)	0.050(5)	0.092(6)	0.080(6)	0.020(4)	−0.025(4)	−0.018(5)
C(22)	2a	0.0445(4)	0.0236(6)	0.7000(4)	0.014(2)	0.038(3)	0.044(3)	0.000(4)	0.005(2)	−0.006(4)
C(23)	2a	−0.0282(6)	−0.0419(5)	0.7671(6)	0.031(3)	0.051(4)	0.058(4)	−0.020(3)	0.027(3)	−0.015(3)
C(24)	2a	0.0011(6)	−0.0079(5)	0.8965(5)	0.029(3)	0.062(5)	0.050(4)	−0.014(3)	0.027(3)	−0.006(3)
C(25)	2a	0.1582(5)	0.0104(4)	0.9445(5)	0.032(3)	0.030(4)	0.044(3)	−0.006(3)	0.021(2)	0.007(3)
C(26)	2a	0.0659(6)	−0.0181(4)	0.5863(5)	0.037(3)	0.044(3)	0.034(3)	0.001(3)	0.009(3)	0.006(3)
C(27)	2a	0.1843(9)	−0.1397(7)	0.5215(8)	0.075(6)	0.079(6)	0.070(6)	0.009(5)	0.050(5)	−0.009(5)
C(28)	2a	0.2069(6)	0.0618(4)	1.0621(5)	0.036(3)	0.045(3)	0.037(4)	−0.009(2)	0.010(3)	0.002(3)
C(29)	2a	0.227(1)	0.0511(6)	1.2686(6)	0.132(8)	0.093(8)	0.035(4)	−0.030(6)	0.019(4)	−0.004(4)

Acknowledgments. We thank the Fonds der Chemischen Industrie and the DFG.

References

- Schrock, R. R.; Osborn, J. A.: Preparation and Properties of Some Cationic Complexes of Rhodium(I) and Rhodium(III). *J. Am. Chem. Soc.* **93** (1971) 2397–2407.
- Holz, J.; Stürmer, R.; Schmidt, U.; Drexler, H.-J.; Heller, D.; Krimmer, H.-P.; Börner, A.: Synthesis of chiral 2,5-Bis(oxymethyl)-functionalized bis(phospholanes) and their application in Rh- and Ru-catalyzed enantioselective hydrogenations. *Eur. J. Org. Chem.* (2001) 4615–4624.
- Drexler, H.-J.; Baumann, W.; Spannenberg, A.; Heller, D.: unpublished results.
- Sheldrick, G. M.: SHELXS-97. Program for the Solution of Crystal Structures. University of Göttingen, Germany 1997.
- Sheldrick, G. M.: SHELXL-97. Program for the Refinement of Crystal Structures. University of Göttingen, Germany 1997.