Crystal structure of spiro-[2.3'']oxindole-spiro[3.3']-5'-(4-methylsulfanylphenylmethylidene)-1'-benzyl-4'-(1*H*)-pyridinone-4-(4-methylsulfanylphenyl)hexahydro-1*H*-pyrrolizine — dichloromethane, $C_{40}H_{39}N_3O_2S_2 \cdot CH_2Cl_2$

Xun Li^{*I} and Aiting Zheng^{II}

^I Changsha University of Science and Technology, Department of Chemical and Environmental Engineering, Changsha, 410114 Hunan, P. R. China

^{II} Hunan University of Science and Technology, School of Chemistry and Chemical Engineering, Xiangtan 411201, P. R. China

Received October 16, 2009; accepted and available on-line November 3, 2009; CCDC no. 1267/2808

Abstract

$C_{41}H_{41}Cl_2N_3O_2S_2$, triclinic, $P\bar{1}$ (no. 2), $a = 9.1853(9)$ Å, $b = 12.846(1)$ Å, $c = 17.559(2)$ Å, $\alpha = 70.299(9)^\circ$, $\beta = 77.36(1)^\circ$, $\gamma = 71.996(9)^\circ$, $V = 1839.8$ Å³, $Z = 2$, $R_g(F) = 0.044$, $wR_{ref}(F^2) = 0.097$, $T = 113$ K.

Source of material

A mixture of 1-benzyl-3,5-bis(4-(methylthio)benzylidene)-piperidin-4-one (1 mmol), isatin (1 mmol) and proline (1 mmol) were refluxed in methanol (60 ml) until the complete dissolution of the starting material as evidenced by the TLC. After the reaction was over, the solvent was removed in vacuo and the residue was separated by column chromatography (silica gel, petroleum ether/ethylacetate = 5:1) to give the title compound. 20 mg of the title compound was dissolved in 15 ml dichloromethane/hexane mixed solvent; the solution was kept at room temperature for 15 d by natural evaporation to give colorless single crystals suitable for X-ray diffraction analysis.

Experimental details

The H atoms bound to C were included in the riding model approximation with $d(C-H) = 0.95 - 0.99$ Å and $U_{iso}(H) = 1.2 U_{eq}(C)$ thermal parameters related to the atoms to which they were bonded. The H3 atom bound to N3 was found in difference Fourier map and refined freely.

Discussion

In the title compound, the two spiro junctions link a 2-oxindole ring, a hexahydro-1*H*-pyrrolizine ring and a piperidine ring in a chair conformation. Two molecules are connected into a dimer by two N–H...O hydrogen bonds.

Table 1. Data collection and handling.

Crystal:	colorless chip, size 0.10 × 0.20 × 0.24 mm
Wavelength:	Mo K_α radiation (0.71070 Å)
μ :	3.31 cm ^{−1}
Diffractometer, scan mode:	Rigaku Saturn CCD, ϕ/ω
$2\theta_{max}$:	55.82°
$N(hkl)_{measured}$, $N(hkl)_{unique}$:	18478, 8642
Criterion for I_{obs} , $N(hkl)_{gt}$:	$I_{obs} > 2\sigma(I_{obs})$, 5217
$N(param)_{refined}$:	480
Programs:	SHELXS-97 [1], SHELXL-97 [2], SHELXTL [3]

Table 2. Atomic coordinates and displacement parameters (in Å²).

Atom	Site	Occ.	x	y	z	U_{iso}
H(3)	2i		1.022(3)	0.586(2)	0.930(1)	0.045(7)
H(5)	2i		1.2692	0.5402	0.6508	0.027
H(6)	2i		1.2851	0.7305	0.6023	0.031
H(7)	2i		1.1814	0.8517	0.6843	0.033
H(8)	2i		1.0695	0.7831	0.8195	0.028
H(10)	2i		1.1921	0.2560	0.7320	0.021
H(11)	2i		1.1900	0.2092	0.9045	0.025
H(12A)	2i		1.4674	0.3965	0.7581	0.031
H(12B)	2i		1.3773	0.3539	0.7098	0.031
H(13A)	2i		1.5811	0.2130	0.8294	0.040
H(13B)	2i		1.5750	0.1919	0.7449	0.040
H(14A)	2i		1.3892	0.0978	0.8028	0.036
H(14B)	2i		1.4265	0.0932	0.8895	0.036
H(16)	2i		1.0447	0.1665	0.6896	0.025
H(17)	2i		0.9042	0.0321	0.7183	0.029
H(19)	2i		0.8663	−0.0016	0.9584	0.027
H(20)	2i		1.0071	0.1309	0.9299	0.029
H(21A)	2i		0.6639	−0.0255	0.7501	0.048
H(21B)	2i		0.6381	−0.1506	0.7950	0.048
H(21C)	2i		0.8050	−0.1360	0.7498	0.048
H(23A)	2i		0.8093	0.3513	0.8061	0.020
H(23B)	2i		0.8818	0.3293	0.8871	0.020
H(24A)	2i		0.7023	0.6472	0.7544	0.023
H(24B)	2i		0.6590	0.5426	0.7427	0.023
H(26)	2i		0.9762	0.6301	0.5792	0.025
H(28)	2i		0.5834	0.7424	0.6460	0.026
H(29)	2i		0.4519	0.9346	0.6221	0.027

* Correspondence author (e-mail: lixun7212@163.com)

Table 2. Continued.

Atom	Site	Occ.	x	y	z	U _{iso}
H(31)	2i		0.8546	1.0222	0.5231	0.038
H(32)	2i		0.9826	0.8319	0.5427	0.035
H(33A)	2i		0.3178	1.1233	0.5612	0.052
H(33B)	2i		0.3004	1.2337	0.5889	0.052
H(33C)	2i		0.3429	1.1087	0.6520	0.052
H(34A)	2i		0.6784	0.4314	0.9456	0.026
H(34B)	2i		0.5753	0.4675	0.8739	0.026
H(36)	2i		0.3980	0.6548	0.8414	0.031

Table 2. Continued.

Atom	Site	Occ.	x	y	z	U _{iso}
H(37)	2i		0.2700	0.8224	0.8766	0.040
H(38)	2i		0.3621	0.8711	0.9705	0.040
H(39)	2i		0.5838	0.7517	1.0282	0.035
H(40)	2i		0.7136	0.5866	0.9907	0.028
H(41A)	2i	0.507	0.7357	0.4486	0.4889	0.046
H(41B)	2i	0.507	0.8380	0.4885	0.5313	0.046
H(41C)	2i	0.493	0.8055	0.4866	0.4782	0.046
H(41D)	2i	0.493	0.8469	0.4737	0.5654	0.046

Table 3. Atomic coordinates and displacement parameters (in Å²).

Atom	Site	Occ.	x	y	z	U ₁₁	U ₂₂	U ₃₃	U ₁₂	U ₁₃	U ₂₃
S(1)	2i		0.75883(6)	−0.09942(4)	0.87441(3)	0.0262(3)	0.0275(3)	0.0352(3)	−0.0121(2)	0.0015(2)	−0.0102(2)
S(2)	2i		0.55864(6)	1.14480(4)	0.56164(3)	0.0329(3)	0.0196(3)	0.0403(3)	−0.0043(2)	−0.0019(2)	−0.0043(2)
O(1)	2i		1.0313(1)	0.3784(1)	0.96002(7)	0.0255(7)	0.0217(7)	0.0161(7)	−0.0065(5)	−0.0005(6)	−0.0037(5)
O(2)	2i		1.0803(2)	0.4250(1)	0.63371(7)	0.0273(8)	0.0224(7)	0.0186(7)	−0.0031(6)	0.0021(6)	−0.0060(6)
N(1)	2i		1.2805(2)	0.3407(1)	0.82881(9)	0.0150(8)	0.0173(8)	0.0258(9)	−0.0017(6)	−0.0001(7)	−0.0076(7)
N(2)	2i		0.7880(2)	0.4955(1)	0.83381(8)	0.0141(8)	0.0150(7)	0.0141(8)	−0.0016(6)	0.0015(6)	−0.0008(6)
N(3)	2i		1.0525(2)	0.5611(1)	0.8915(1)	0.0191(9)	0.0203(8)	0.0190(9)	−0.0038(6)	0.0001(7)	−0.0087(7)
C(1)	2i		1.1352(2)	0.4304(1)	0.8130(1)	0.0146(9)	0.0178(9)	0.021(1)	−0.0017(7)	−0.0017(8)	−0.0069(8)
C(2)	2i		1.0632(2)	0.4515(2)	0.8970(1)	0.0136(9)	0.0208(9)	0.021(1)	−0.0015(7)	−0.0018(8)	−0.0087(8)
C(3)	2i		1.1062(2)	0.6206(1)	0.8129(1)	0.0116(9)	0.0196(9)	0.022(1)	−0.0022(7)	−0.0038(7)	−0.0048(8)
C(4)	2i		1.1610(2)	0.5477(1)	0.7639(1)	0.0120(9)	0.0174(9)	0.021(1)	−0.0043(7)	−0.0016(7)	−0.0048(8)
C(5)	2i		1.2288(2)	0.5889(2)	0.6850(1)	0.017(1)	0.024(1)	0.025(1)	−0.0057(8)	0.0005(8)	−0.0085(8)
C(6)	2i		1.2375(2)	0.7023(2)	0.6562(1)	0.019(1)	0.028(1)	0.027(1)	−0.0099(8)	−0.0004(8)	−0.0015(9)
C(7)	2i		1.1772(2)	0.7741(2)	0.7054(1)	0.022(1)	0.018(1)	0.038(1)	−0.0070(8)	−0.0031(9)	−0.0010(9)
C(8)	2i		1.1104(2)	0.7343(2)	0.7853(1)	0.016(1)	0.021(1)	0.034(1)	−0.0012(8)	−0.0050(9)	−0.0098(9)
C(9)	2i		1.0298(2)	0.3787(1)	0.7807(1)	0.0144(9)	0.0153(9)	0.017(1)	−0.0029(7)	−0.0002(7)	−0.0054(7)
C(10)	2i		1.1290(2)	0.2544(1)	0.7864(1)	0.018(1)	0.0166(9)	0.017(1)	−0.0025(7)	−0.0007(7)	−0.0071(7)
C(11)	2i		1.2413(2)	0.2316(1)	0.8471(1)	0.019(1)	0.0173(9)	0.025(1)	0.0001(7)	−0.0052(8)	−0.0074(8)
C(12)	2i		1.4117(2)	0.3393(2)	0.7629(1)	0.015(1)	0.029(1)	0.032(1)	−0.0040(8)	0.0024(8)	−0.0126(9)
C(13)	2i		1.5117(2)	0.2178(2)	0.7918(1)	0.021(1)	0.032(1)	0.040(1)	0.0032(9)	−0.0015(9)	−0.014(1)
C(14)	2i		1.3957(2)	0.1446(2)	0.8361(1)	0.024(1)	0.025(1)	0.039(1)	0.0035(9)	−0.0083(9)	−0.0103(9)
C(15)	2i		1.0409(2)	0.1645(1)	0.8058(1)	0.020(1)	0.0135(9)	0.020(1)	−0.0010(7)	−0.0034(8)	−0.0034(8)
C(16)	2i		1.0084(2)	0.1325(2)	0.7447(1)	0.020(1)	0.023(1)	0.019(1)	−0.0039(8)	0.0010(8)	−0.0072(8)
C(17)	2i		0.9241(2)	0.0523(2)	0.7616(1)	0.022(1)	0.027(1)	0.026(1)	−0.0062(8)	−0.0007(8)	−0.0128(9)
C(18)	2i		0.8692(2)	0.0018(1)	0.8417(1)	0.017(1)	0.0157(9)	0.029(1)	−0.0009(7)	−0.0033(8)	−0.0065(8)
C(19)	2i		0.9023(2)	0.0327(1)	0.9033(1)	0.027(1)	0.0181(9)	0.019(1)	−0.0047(8)	−0.0042(8)	−0.0003(8)
C(20)	2i		0.9861(2)	0.1118(1)	0.8864(1)	0.030(1)	0.021(1)	0.020(1)	−0.0052(8)	−0.0056(9)	−0.0049(8)
C(21)	2i		0.7110(2)	−0.1033(2)	0.7818(1)	0.025(1)	0.034(1)	0.043(1)	−0.0119(9)	−0.002(1)	−0.017(1)
C(22)	2i		1.0013(2)	0.4509(1)	0.6937(1)	0.0160(9)	0.0190(9)	0.019(1)	−0.0065(7)	−0.0004(7)	−0.0061(8)
C(23)	2i		0.8698(2)	0.3803(1)	0.8310(1)	0.0167(9)	0.0146(9)	0.016(1)	−0.0034(7)	−0.0006(7)	−0.0032(7)
C(24)	2i		0.7430(2)	0.5666(1)	0.7537(1)	0.0154(9)	0.0187(9)	0.019(1)	−0.0025(7)	−0.0005(8)	−0.0027(8)
C(25)	2i		0.8747(2)	0.5591(1)	0.6856(1)	0.0171(9)	0.0199(9)	0.016(1)	−0.0052(7)	−0.0013(7)	−0.0045(8)
C(26)	2i		0.8920(2)	0.6471(1)	0.6193(1)	0.018(1)	0.022(1)	0.018(1)	−0.0039(8)	−0.0012(8)	−0.0025(8)
C(27)	2i		0.7987(2)	0.7655(2)	0.6005(1)	0.019(1)	0.021(1)	0.019(1)	−0.0035(8)	−0.0022(8)	−0.0016(8)
C(28)	2i		0.6393(2)	0.7991(2)	0.6217(1)	0.021(1)	0.021(1)	0.018(1)	−0.0066(8)	−0.0026(8)	0.0003(8)
C(29)	2i		0.5607(2)	0.9138(2)	0.6081(1)	0.019(1)	0.026(1)	0.019(1)	−0.0050(8)	−0.0015(8)	−0.0040(8)
C(30)	2i		0.6404(2)	0.9983(2)	0.5742(1)	0.025(1)	0.020(1)	0.021(1)	−0.0035(8)	−0.0044(8)	0.0006(8)
C(31)	2i		0.7992(2)	0.9654(2)	0.5490(1)	0.024(1)	0.024(1)	0.038(1)	−0.0096(9)	−0.0013(9)	0.0031(9)
C(32)	2i		0.8753(2)	0.8522(2)	0.5613(1)	0.020(1)	0.027(1)	0.029(1)	−0.0045(8)	0.0024(9)	0.0037(9)
C(33)	2i		0.3571(2)	1.1536(2)	0.5947(1)	0.036(1)	0.028(1)	0.029(1)	0.0022(9)	0.0055(9)	−0.0089(9)
C(34)	2i		0.6488(2)	0.4908(1)	0.8944(1)	0.020(1)	0.0201(9)	0.021(1)	−0.0067(8)	0.0045(8)	−0.0042(8)
C(35)	2i		0.5691(2)	0.6024(1)	0.9130(1)	0.0143(9)	0.0204(9)	0.019(1)	−0.0064(7)	0.0051(7)	−0.0050(8)
C(36)	2i		0.4365(2)	0.6743(2)	0.8793(1)	0.018(1)	0.030(1)	0.030(1)	−0.0042(8)	−0.0032(9)	−0.0116(9)
C(37)	2i		0.3601(2)	0.7739(2)	0.9003(1)	0.020(1)	0.029(1)	0.045(1)	0.0026(9)	−0.004(1)	−0.011(1)
C(38)	2i		0.4147(2)	0.8030(2)	0.9559(1)	0.032(1)	0.027(1)	0.039(1)	−0.0082(9)	0.006(1)	−0.015(1)
C(39)	2i		0.5461(2)	0.7325(2)	0.9898(1)	0.034(1)	0.032(1)	0.029(1)	−0.018(1)	0.0048(9)	−0.0130(9)
C(40)	2i		0.6223(2)	0.6340(2)	0.9678(1)	0.019(1)	0.027(1)	0.022(1)	−0.0089(8)	0.0007(8)	−0.0034(9)
C(41)	2i	0.507(9)	0.744(1)	0.4616(6)	0.5401(8)	0.025(3)	0.041(1)	0.040(2)	−0.008(2)	0.012(2)	−0.011(1)
Cl(1)	2i	0.507	0.5841(4)	0.5669(3)	0.5617(2)	0.020(1)	0.026(1)	0.059(2)	0.0011(8)	−0.0054(9)	−0.0228(9)
Cl(2)	2i	0.507	0.7616(3)	0.3348(2)	0.6167(3)	0.063(1)	0.0311(8)	0.064(2)	−0.0157(7)	−0.022(1)	0.009(1)
C(41')	2i	0.493	0.770(1)	0.4653(6)	0.5375(9)	0.025(3)	0.041(1)	0.040(2)	−0.008(2)	0.012(2)	−0.011(1)
Cl(1')	2i	0.493	0.5932(6)	0.5572(4)	0.5568(3)	0.049(2)	0.071(2)	0.160(4)	0.006(2)	−0.010(2)	−0.070(2)
Cl(2')	2i	0.493	0.7617(3)	0.3253(2)	0.5702(4)	0.071(1)	0.0341(8)	0.071(3)	−0.0132(7)	0.005(1)	−0.0109(9)

Acknowledgments. Financial supports from China Ministry of Science and Technology's Foundation under grant nos. 2008AA05Z405, 2007BAD41B05 and 2006DFA62370 and from Hunan Provincial Bureau of Science and Technology's Foundation under grant nos. 06GK3066 and 05C229 are highly appreciated.

References

1. Sheldrick, G. M.: SHELXS-97. Program for the Solution of Crystal Structures. University of Göttingen, Germany 1997.
2. Sheldrick, G. M.: SHELXL-97. Program for the Refinement of Crystal Structures. University of Göttingen, Germany 1997.
3. Sheldrick, G. M.: SHELXTL. Structure Determination Software Suite. Version 6.14. Bruker AXS, Madison, Wisconsin, USA 2000.