

# On the torsion of linear higher order connections

Ivan Kolář\*

*Department of Algebra and Geometry  
Faculty of Science, Masaryk University  
Janáčkovo nám. 2a, 662 95 Brno, Czech Republic*

Received 6 December 2002; accepted 27 March 2003

---

**Abstract:** For a linear  $r$ -th order connection on the tangent bundle we characterize geometrically its integrability in the sense of the theory of higher order  $G$ -structures. Our main tool is a bijection between these connections and the principal connections on the  $r$ -th order frame bundle and the comparison of the torsions under both approaches.

© Central European Science Journals. All rights reserved.

*Keywords:* linear  $r$ -th order connection on the tangent bundle, principal connection on the  $r$ -th order frame bundle, torsion, integrability of  $G$ -structures

*MSC (2000):* 53C05, 58A20

---

## 1 Introduction

Our starting point was the problem of integrability (in the sense of the theory of higher order  $G$ -structures) of a linear  $r$ -th order connection  $\Gamma$  on the tangent bundle of a manifold  $M$ . This kind of integrability plays an important role in our theory (jointly with A. Cabras) of the flow prolongation of some tangent valued forms, [1]. Our solution of the problem is heavily based on the concept of the torsion of  $\Gamma$ . In particular, we have to compare two different approaches to the torsion in higher order. This leads us to certain new geometric results. In our opinion, the greater part of them is of individual interest.

In Section 2 we recall the definition of the torsion  $\tau_\Gamma$  of  $\Gamma$  by A. Zajtz, [6], in which the jet factorization of the bracket of vector fields is used. Then we deduce that the difference of two torsion free linear  $r$ -th order connections that coincide up to order  $r - 1$  is an arbitrary tensor field of the type  $TM \otimes S^{r+1}T^*M$ . In Section 3 we clarify that  $\Gamma$  is equivalent to a principal connection  $\tilde{\Gamma}$  on the  $r$ -th order frame bundle  $P^rM$ . In the latter case, the torsion was introduced by P.C.Yuen, [5], as the exterior covariant

---

\* E-mail: kolar@math.muni.cz

differential  $D_{\bar{\Gamma}}\Theta_r$  of the canonical  $(\mathbb{R}^m \times \mathfrak{g}_m^{r-1})$ -valued 1-form  $\Theta_r$  on  $P^rM$ . The main result of Section 4 reads that  $\tau_{\Gamma}$  and  $D_{\bar{\Gamma}}\Theta_r$  coincide in fact. In the proof of Lemma 4.3 we use as an essential tool the fact that the linear natural operators commute with the Lie differentiation. Section 5 starts with a modification of the concept of the higher order exponential operator from [3] to the case of higher order linear connections on the tangent bundle. This idea enables us to characterize the integrability of  $\Gamma$  in a transparent way.

All manifolds and maps are assumed to be infinitely differentiable. Unless otherwise specified, we use the terminology and notation from the book [4].

## 2 Linear higher order connections on $TM$

Consider the tangent bundle  $p : TM \rightarrow M$  of an  $m$ -dimensional manifold  $M$  and its  $r$ -th jet prolongation  $J^rTM$ . We recall that  $\pi_k^r$  denotes the canonical projection of  $r$ -jets into  $k$ -jets,  $r \geq k \geq 0$ .

**Definition 2.1.** A linear  $r$ -th order connection on  $M$  means a linear base preserving morphism  $\Gamma : TM \rightarrow J^rTM$  satisfying  $\pi_0^r \circ \Gamma = \text{id}_{TM}$ .

If  $x^i$  are some local coordinates on  $M$ ,  $X^i = dx^i$  are the induced coordinates on  $TM$ , and  $X_{\alpha}^i$  are the jet coordinates on  $J^rTM$ , then the equations of  $\Gamma$  are

$$X_{\alpha}^i = \Gamma_{\alpha j}^i(x)X^j, \quad 1 \leq |\alpha| \leq r, \tag{1}$$

where  $\alpha$  is a multi-index of range  $m$ .

The  $(r - 1)$ -jet at  $x \in M$  of the bracket  $[\xi, \eta]$  of two vector fields  $\xi, \eta$  on  $M$  depends on the  $r$ -jets  $j_x^r\xi$  and  $j_x^r\eta$ . This defines a map

$$[\ , \ ]_{r-1} : J^rTM \times_M J^rTM \rightarrow J^{r-1}TM.$$

A. Zajtzy, [6], introduced the torsion of  $\Gamma$  as a map  $\tau_M : TM \times_M TM \rightarrow J^{r-1}TM$  defined by

$$\tau_{\Gamma}(X, Y) = [\Gamma(X), \Gamma(Y)]_{r-1}, \quad X, Y \in T_xM. \tag{2}$$

For  $r = 1$ , (1) is of the form  $X_k^i = \Gamma_{kj}^i(x)X^j$  and one deduces easily that the coordinate expression of  $\tau_{\Gamma}$  is  $\Gamma_{jk}^i - \Gamma_{kj}^i$ , so that we have the classical concept of torsion.

Let  $\Delta$  be another linear  $r$ -th order connection on  $M$  satisfying  $\pi_{r-1}^r \circ \Delta = \pi_{r-1}^r \circ \Gamma$ . Since the kernel of  $\pi_{r-1}^r : J^rTM \rightarrow J^{r-1}TM$  is  $TM \otimes S^rT^*M$ , the difference of  $\Gamma$  and  $\Delta$  is a section

$$\Gamma - \Delta : M \rightarrow TM \otimes S^rT^*M \otimes T^*M. \tag{3}$$

**Proposition 2.2.** If both  $\Gamma$  and  $\Delta$  are torsion free, then the values of  $\Gamma - \Delta$  lie in  $TM \otimes S^{r+1}T^*M$ .

**Proof.** The only  $r$ -th order terms in  $j^{r-1}[\xi, \eta]$  are

$$\xi^j \frac{\partial^r \eta^i}{\partial x^j \partial \beta x} - \eta^j \frac{\partial^r \xi^i}{\partial x^j \partial \beta x}, \quad |\beta| = r - 1. \tag{4}$$

Let  $\Delta^i_{\alpha j}$  be the Christoffels of  $\Delta$ . Then  $\Gamma - \Delta$  can be interpreted as a map  $TM \times_M TM \rightarrow TM \otimes S^{r-1}T^*M$  of the form

$$(\Gamma^i_{\beta j k} - \Delta^i_{\beta j k})\xi^j \eta^k = 0.$$

If both  $\Gamma$  and  $\Delta$  are torsion free, then (4) implies

$$(\Gamma^i_{\beta j k} - \Gamma^i_{\beta k j} - \Delta^i_{\beta j k} + \Delta^i_{\beta k j})\xi^j \eta^k, \quad |\beta| = r - 1.$$

This proves our claim. □

**Remark 2.3.** One sees directly from the proof that the difference  $\Gamma - \Delta$  is an arbitrary section of  $TM \otimes S^{r+1}T^*M$ . In a different form this result was deduced in [3].

On  $\mathbb{R}^m$  there is a distinguished linear  $r$ -th order connection  $I_r : T\mathbb{R}^m \rightarrow J^r T\mathbb{R}^m$  called the canonical integrable connection. It is defined by

$$I_r(X) = j_x^r \tilde{X},$$

where  $\tilde{X}$  is the constant vector field constructed from  $X$  by means of the translations on  $\mathbb{R}^m$ . In [1] we introduced

**Definition 2.4.** A linear  $r$ -th order connection  $\Gamma$  on  $M$  is said to be integrable, if for every  $x \in M$  there exists a neighbourhood  $U$  and a diffeomorphism  $f : U \rightarrow \mathbb{R}^m$  satisfying  $I \circ Tf = J^r T f \circ (\Gamma|_U)$ .

Clearly, every integrable  $\Gamma$  is torsion free.

### 3 Principal connections on $P^r M$

Consider the  $r$ -th order frame bundle  $P^r M$  of  $M$ . This is an open subset of the space  $T_m^r M$  of all  $(m, r)$ -velocities on  $M$ . We recall that there is a natural exchange diffeomorphism  $\varkappa_M : T_m^r TM \rightarrow TT_m^r M$  with the property that for every vector field  $\xi : M \rightarrow TM$  the flow prolongation  $\mathcal{J}_m^r \xi$  satisfies  $\mathcal{J}_m^r \xi = \varkappa_M \circ T_m^r \xi$ , where  $T_m^r \xi : T_m M \rightarrow T_m^r TM$ , [4]. The flow prolongation  $\mathcal{P}^r \xi$  is the restriction of  $\mathcal{J}_m^r \xi$  to  $P^r M \subset T_m^r M$ .

To establish a bijection between the linear  $r$ -th order connections on  $M$  and the principal connections on  $P^r M$ , we introduce a map

$$i : J^r TM \times_M P^r M \rightarrow TP^r M.$$

For  $X = j_x^r \xi$ , we construct the flow prolongation  $\mathcal{P}^r \xi$ . The value  $\mathcal{P}^r \xi(u)$ ,  $u \in P_x^r M$ , depends on  $j_x^r \xi$  only, so that we can set

$$i(X, u) = \mathcal{P}^r \xi(u). \tag{5}$$

On the other hand, for  $Y \in T_u P^r M$  we have  $\varkappa_M^{-1}(Y) \in T_m^r TM = J_0^r(\mathbb{R}^m, TM)$ . Since  $u \in J_0^r(\mathbb{R}^m, M)$ , we can construct the jet composition  $\varkappa_M^{-1}(Y) \circ u^{-1} \in J_x^r(M, TM)$ . We have  $(T_m^r p)(\varkappa_M^{-1}Y \circ u^{-1}) = (T_m^r p)(\varkappa_M^{-1}Y) \circ u^{-1} = u \circ u^{-1} = j_x^r \text{id}_M$ , so that  $(\varkappa_M^{-1}Y) \circ u \in J_x^r TM$ . Clearly,

$$i((\varkappa_M^{-1}Y) \circ u^{-1}, u) = Y.$$

Hence  $i$  is a diffeomorphism.

Since  $P^r$  is a functor with values in the category of principal bundles,  $\mathcal{P}^r \xi$  is a right invariant vector field on  $P^r M$ , [4]. Hence for every  $\Gamma$  the rule

$$\tilde{\Gamma}(u, X) = \mathcal{P}^r \xi(u), \quad \Gamma(X) = j_x^r \xi, \tag{6}$$

defines the lifting map  $\tilde{\Gamma} : P^r M \times_M TM \rightarrow TP^r M$  of a principal connection  $\tilde{\Gamma}$  on  $P^r M$ . Indeed, the linearity in  $TM$  follows from the basic properties of  $\varkappa_M$ , [4]. Conversely, if  $\Delta : TM \times_M P^r M \rightarrow TP^r M$  is the lifting map of a right invariant connection on  $P^r M$ , then

$$i^{-1} \circ \Delta : TM \times_M P^r M \rightarrow J^r TM \times_M P^r M$$

is of the form  $\Gamma \times_M \text{id}_{P^r M}$ , where  $\Gamma : TM \rightarrow J^r TM$  is a linear splitting. This proves

**Proposition 3.1.** The rule (6) establishes a bijection between the linear  $r$ -th order connections on  $M$  and the principal connections on  $P^r M$ .

Write  $e_{r-1} = j_0^{r-1} \text{id}_{\mathbb{R}^m}$  and  $V_m^{r-1} = \mathbb{R}^m \times \mathfrak{g}_m^{r-1} = T_{e_{r-1}} P^{r-1} \mathbb{R}^m$ . The canonical  $V_m^{r-1}$ -valued 1-form on  $P^r M$  is defined as follows, [4]. Every  $u = j_0^r f$ ,  $f : \mathbb{R}^m \rightarrow M$ , induces  $P^{r-1} f : P^{r-1} \mathbb{R}^m \rightarrow P^{r-1} M$ . The tangent map  $\tilde{u} := T_{e_{r-1}} P^{r-1} f : T_{e_{r-1}} P^{r-1} \mathbb{R}^m \rightarrow T_{u_{r-1}} P^{r-1} M$ ,  $u_{r-1} = \pi_{r-1}^r(u)$  depends on  $u$  only. Then one defines

$$\Theta_r(A) = \tilde{u}^{-1}(T\pi_{r-1}^r(A)), \quad A \in T_u P^r M. \tag{7}$$

P.C.Yuen, [5], introduced the torsion of  $\tilde{\Gamma}$  as the exterior covariant differential  $D_{\tilde{\Gamma}} \Theta_r$  of  $\Theta_r$ .

### 4 The comparison of torsions

Since  $J^{r-1} TM$  is an  $r$ -th order natural bundle, it is a bundle associated to  $P^r M$  with the standard fiber  $U_m^{r-1} := J_0^{r-1} T\mathbb{R}^m$ , [4]. Every  $u = j_0^r h \in P_x^r M$  is interpreted as a frame map  $\bar{u} : U_m^{r-1} \rightarrow J_x^{r-1} TM$  as follows. We construct  $Th : T\mathbb{R}^m \rightarrow TM$  and set  $\bar{u} = J_0^{r-1} Th$ . We have  $J_0^{r-1} T\mathbb{R}^m \subset T_m^{r-1} T\mathbb{R}^m$  and the exchange  $\varkappa_M : T_m^{r-1} TM \rightarrow TT_m^{r-1} M$  maps  $U_m^{r-1}$  into  $V_m^{r-1}$ . Indeed,  $j_0^{r-1} \frac{\partial}{\partial t} \Big|_0 \psi(t, \tau) \in J_0^{r-1} T\mathbb{R}^m$  if and only if  $\psi(0, \tau) = \text{id}_{\mathbb{R}^m}$ . Then  $\frac{\partial}{\partial t} \Big|_0 j_0^{r-1} \psi(t, \tau) \in T_{e_{r-1}} P^{r-1} M$  and vice versa. This yields an identification

$$\varkappa_0 : U_m^{r-1} \rightarrow V_m^{r-1}.$$

**Lemma 4.1.** For every vector field  $\xi$  on  $M$  and every  $u \in P_x^r M$ , we have

$$\Theta_r(\mathcal{P}^r \xi(u)) = \varkappa_0(\bar{u}^{-1}(j_x^{r-1} \xi)). \tag{8}$$

**Proof.** The naturality of  $\varkappa$  on  $h : \mathbb{R}^m \rightarrow M$  yields a commutative diagram

$$\begin{CD} T_m^{r-1} T\mathbb{R}^m @>T_m^{r-1} Th>> T_m^{r-1} TM \\ @V{\varkappa_{\mathbb{R}^m}}VV @VV{\varkappa_M}V \\ TT^{r-1}\mathbb{R} @>TT_m^{r-1} h>> TT_m^{r-1} M \end{CD} \tag{9}$$

The right column of (9) maps  $j_x^{r-1} \xi$  into  $\mathcal{P}_x^{r-1} \xi(u_{r-1})$ ,  $u = j_0^r h$ . Then (8) follows directly from the definitions of  $\tilde{u}$ ,  $\bar{u}$  and  $\varkappa_0$ . □

Since  $D_{\bar{\Gamma}} \Theta_r$  is a horizontal  $V_m^{r-1}$ -valued 2-form on  $P^r M$ , it can be interpreted as a map  $P^r M \rightarrow V_m^{r-1} \otimes \wedge^2 T^* M$ , see e.g. [4], p. 112. Taking into account the identification  $\tilde{u}_1 : \mathbb{R}^m \rightarrow T_x M$ ,  $u_1 = \pi_1^r(u)$ , we construct

$$\overline{D}_{\bar{\Gamma}} \Theta_r : P^r M \rightarrow V_m^{r-1} \otimes \wedge^2 \mathbb{R}^m. \tag{10}$$

On the other hand,  $\tau_\Gamma$  can be interpreted as a section of  $J^{r-1} TM \otimes \wedge^2 T^* M$ . So its frame form, see [4], is a map

$$\overline{\tau}_\Gamma : P^r M \rightarrow U_m^{r-1} \otimes \wedge^2 \mathbb{R}^{m*}.$$

**Proposition 4.2.** Under the identification  $\varkappa_0$ , we have  $\overline{D}_{\bar{\Gamma}} \Theta_r = \frac{1}{2} \overline{\tau}_\Gamma$ .

**Proof.** The most important part of the proof is the following lemma. Let  $\xi, \eta$  be two vector fields on  $M$ . Then  $\Theta_r(\mathcal{P}^r \eta)$  is a  $V_m^{r-1}$ -valued function on  $P^r M$ , so that we can construct its derivation  $(\mathcal{P}^r \xi) \Theta_r(\mathcal{P}^r \eta)$  in the direction of  $\mathcal{P}^r \xi$ , which is another  $V_m^{r-1}$ -valued function on  $P^r M$ .

**Lemma 4.3.** We have  $(\mathcal{P}^r \xi) \Theta_r(\mathcal{P}^r \eta) = \Theta_r(\mathcal{P}^r([\xi, \eta]))$ .

**Proof.** As a direct consequence of the definition,  $\Theta_r$  is natural, i.e. for every local diffeomorphism  $f : M \rightarrow \overline{M}$  we have

$$\Theta_{r,M} = \Theta_{r,\overline{M}} \circ TP^r f.$$

Consider the operator  $\tilde{\Theta}_r$  transforming each vector field on  $M$  into a  $V_m^{r-1}$ -valued function on  $P^r M$

$$\tilde{\Theta}_r(\eta) = \Theta_r(\mathcal{P}^r \eta).$$

Clearly,  $\tilde{\Theta}_r$  is linear and natural. By Proposition 49.5 of [4],  $\tilde{\Theta}_r$  commutes with the Lie differentiation, i.e.

$$\mathcal{L}_\xi \tilde{\Theta}_r(\eta) = \tilde{\Theta}_r(\mathcal{L}_\xi \eta).$$

The geometrical meaning of the Lie derivative on the left hand side is  $(\mathcal{P}^r\xi)\Theta_r(\mathcal{P}^r\eta)$ . On the right hand side, we use the well known fact  $\mathcal{L}_\xi\eta = [\xi, \eta]$ .  $\square$

Consider now  $u \in P_x^rM$  and  $X, Y \in T_xM$ ,  $\Gamma(X) = j_x^r\xi$ ,  $\Gamma(Y) = j_x^r\eta$ . If we interpret  $\overline{D}_{\overline{\Gamma}}\Theta_r$  as a map  $P^rM \times \mathbb{R}^m \times \mathbb{R}^m \rightarrow V_m^{r-1}$ , we have

$$\overline{D}_{\overline{\Gamma}}\Theta_r(u, \tilde{u}_1^{-1}(X), \tilde{u}_1^{-1}(Y)) = d\Theta_r(\mathcal{P}^r\xi(u), \mathcal{P}^r\eta(u)).$$

Applying Lemma 4.3, we obtain

$$\begin{aligned} 2d\Theta_r(\mathcal{P}^r\xi, \mathcal{P}^r\eta) &= (\mathcal{P}^r\xi)\Theta_r(\mathcal{P}^r\eta) - (\mathcal{P}^r\eta)\Theta_r(\mathcal{P}^r\xi) - \Theta_r([\mathcal{P}^r\xi, \mathcal{P}^r\eta]) \\ &= \Theta_r([\mathcal{P}^r\xi, \mathcal{P}^r\eta]). \end{aligned}$$

By Lemma 4.1, the value of the last expression at  $u$  corresponds to  $[\Gamma(X), \Gamma(Y)]_{r-1}$ . This proves Proposition 4.2.

### 5 The integrability of $\Gamma$

Consider a classical torsion free connection  $\Gamma : TM \rightarrow J^1TM$  and denote by  $\exp_x : T_xM \rightarrow M$  its exponential mapping at  $x \in M$ . We define the  $r$ -th exponential prolongation of  $\Gamma$

$$\varepsilon_r(\Gamma) : TM \rightarrow J^rTM$$

as follows. Every  $X \in T_xM$  is extended into a constant vector field  $\tilde{X}$  on  $T_xM$ . This vector field is transformed by  $\exp_x$  into a vector field on  $M$  and we set

$$\varepsilon_r(\Gamma)(X) = j_x^r(T \exp_x \circ \tilde{X} \circ \exp_x^{-1}). \tag{11}$$

The linearity is obvious, so that  $\varepsilon_r(\Gamma)$  is a linear  $r$ -th order connection on  $M$ . The fact  $[\tilde{X}, \tilde{Y}] = 0$  for another  $Y \in T_xM$  implies that  $\varepsilon_r(\Gamma)$  is torsion free.

Clearly, if  $\Gamma$  is integrable, then  $\varepsilon_r(\Gamma)$  is so.

**Remark 5.1.** We remark that the definition of  $\varepsilon_r$  is based on a similar idea as the concept of the exponential operator  $E_r$  from [3]. In that paper we used the fact that the torsion free connections on  $P^rM$  are in bijection with some reductions of  $P^{r+1}M$ . But a complete comparison of both approaches is beyond the scope of the present paper.

Since every integrable linear  $r$ -th order connection  $\Gamma$  on  $M$  is torsion free, the torsion of  $\Gamma$  is the first obstruction to the integrability of  $\Gamma$ . Consider the underlying connections  $\Gamma_k = \pi_k^r \circ \Gamma$ ,  $k = 1, \dots, r$ . Clearly, if  $\Gamma$  is torsion free or integrable, then each  $\Gamma_k$  is so. Assume that  $\Gamma$  is torsion free. By a classical result, the torsion free connection  $\Gamma_1$  is integrable if and only if its curvature vanishes. This is the second obstruction to the integrability of  $\Gamma$ . If the curvature vanishes, each connection  $\varepsilon_k(\Gamma_1)$  is integrable. The difference

$$\Gamma_2 - \varepsilon_2(\Gamma_1)$$

is a tensor field of the type  $TM \otimes S^3T^*M$  that is the third obstruction to the integrability of  $\Gamma$ . Assume by induction that the first up to  $(k+1)$ -st obstructions to the integrability of  $\Gamma$  vanish. Then  $\Gamma_k = \varepsilon_k(\Gamma_1)$  and the tensor field of type  $TM \otimes S^{k+2}T^*M$

$$\Gamma_{k+1} - \varepsilon_{k+1}(\Gamma_1)$$

is the next obstruction to the integrability of  $\Gamma$ . If all the  $r+1$  obstructions vanish, then  $\Gamma = \varepsilon_r(\Gamma_1)$  is integrable. Thus, we have proved

**Proposition 5.2.**  $\Gamma$  is integrable if and only if all the following conditions are satisfied

- a)  $\Gamma$  is torsion free,
- b)  $\Gamma_1$  is curvature free,
- c) all the gradually defined tensor fields  $\Gamma_k - \varepsilon_k(\Gamma_1)$ ,  $k = 2, \dots, r$  vanish. □

## Acknowledgments

The author was supported by a grant of the GAČR No. 201/02/0225.

## References

- [1] A. Cabras and I. Kolář: *Flow prolongation of some tangent valued forms*, to appear.
- [2] M. Elżanowski and S. Prishchepionok: “Connections on higher order frame bundles”, *New Developments in Differential Geometry, Proceedings*, Kluwer, 1996, pp. 131–142.
- [3] I. Kolář: “Torsion free connections on higher order frame bundles”, *New Developments in Differential Geometry, Proceedings*, Kluwer, 1996, pp. 233–241.
- [4] I. Kolář, P.W. Michor, J. Slovák: *Natural Operations in Differential Geometry*, Springer-Verlag, 1993, <http://www.math.muni.cz/EMIS/monographs/index.html>.
- [5] P.C. Yuen: “Higher order frames and linear connections”, *Cahiers Topol. Geom. Diff.*, Vol. 12, (1971), pp. 333–371.
- [6] A. Zajtš: *Foundations of Differential Geometry of Natural Bundles*, Lecture Notes Univ. Caracas, 1984.