

- [94] Yang Y., Kim A.H., Yamada T., Wu B., Bilimoria P.M., Ikeuchi Y., et al., A Cdc20-APC ubiquitin signaling pathway regulates presynaptic differentiation, *Science*, 2009, 326, 575-578
- [95] Puram S.V., Kim A.H., Ikeuchi Y., Wilson-Grady J.T., Merdes A., Gygi S.P., et al., A CaMKII β signaling pathway at the centrosome regulates dendrite patterning in the brain, *Nat. Neurosci.*, 2011, 14, 973-983
- [96] Yang Y., Kim A.H., Bonni A., The dynamic ubiquitin ligase duo: Cdh1-APC and Cdc20-APC regulate neuronal morphogenesis and connectivity, *Curr. Opin. Neurobiol.*, 2010, 20, 92-99
- [97] Stegmuller J., Konishi Y., Huynh M.A., Yuan Z., Dibacco S., Bonni A., Cell-intrinsic regulation of axonal morphogenesis by the Cdh1-APC target SnoN, *Neuron*, 2006, 50, 389-400
- [98] Stroschein S.L., Bonni S., Wrana J.L., Luo K., Smad3 recruits the anaphase-promoting complex for ubiquitination and degradation of SnoN, *Genes Dev.*, 2001, 15, 2822-2836
- [99] Wan Y., Liu X., Kirschner M.W., The anaphase-promoting complex mediates TGF- β signaling by targeting SnoN for destruction, *Mol. Cell*, 2001, 8, 1027-1039
- [100] Liu X., Sun Y., Weinberg R.A., Lodish H.F., Ski/Sno and TGF- β signaling, *Cytokine Growth Factor Rev.*, 2001, 12, 1-8
- [101] Ikeuchi Y., Stegmuller J., Netherton S., Huynh M.A., Masu M., Frank D., et al., A SnoN-Ccd1 pathway promotes axonal morphogenesis in the mammalian brain, *J. Neurosci.*, 2009, 29, 4312-4321
- [102] Bonni S., Wang H.R., Causing C.G., Kavsak P., Stroschein S.L., Luo K., et al., TGF- β induces assembly of a Smad2-Smurf2 ubiquitin ligase complex that targets SnoN for degradation, *Nat. Cell Biol.*, 2001, 3, 587-595
- [103] Stegmuller J., Huynh M.A., Yuan Z., Konishi Y., Bonni A., TGF β -Smad2 signaling regulates the Cdh1-APC/SnoN pathway of axonal morphogenesis, *J. Neurosci.*, 2008, 28, 1961-1969
- [104] Lasorella A., Stegmuller J., Guardavaccaro D., Liu G., Carro M.S., Rothschild G., et al., Degradation of Id2 by the anaphase-promoting complex couples cell cycle exit and axonal growth, *Nature*, 2006, 442, 471-474
- [105] Zachariae W., Schwab M., Nasmyth K., Seufert W., Control of cyclin ubiquitination by CDK-regulated binding of Hct1 to the anaphase promoting complex, *Science*, 1998, 282, 1721-1724
- [106] Huynh M.A., Stegmuller J., Litterman N., Bonni A., Regulation of Cdh1-APC function in axon growth by Cdh1 phosphorylation, *J. Neurosci.*, 2009, 29, 4322-4327
- [107] Su S.C., Tsai L.H., Cyclin-dependent kinases in brain development and disease, *Annu. Rev. Cell Dev. Biology*, 2011, 27, 465-491
- [108] Bermel C., Tonges L., Planchamp V., Gillardon F., Weishaupt J.H., Dietz G.P., et al., Combined inhibition of Cdk5 and ROCK additively increase cell survival, but not the regenerative response in regenerating retinal ganglion cells, *Mol. Cell. Neurosci.*, 2009, 42, 427-437
- [109] Kannan M., Lee S.J., Schwedhelm-Domeyer N., Stegmuller J., The E3 ligase Cdh1-anaphase promoting complex operates upstream of the E3 ligase Smurf1 in the control of axon growth, *Development*, 2012, 139, 3600-3612
- [110] Wang H.R., Zhang Y., Ozdamar B., Ogunjimi A.A., Alexandrova E., Thomsen G.H., et al., Regulation of cell polarity and protrusion formation by targeting RhoA for degradation, *Science*, 2003, 302, 1775-1779
- [111] Kannan M., Lee S.J., Schwedhelm-Domeyer N., Nakazawa T., Stegmuller J., p250GAP is a novel player in the Cdh1-APC/Smurf1 pathway of axon growth regulation, *PLoS One*, 2012, 7, e50735
- [112] Nakazawa T., Watabe A.M., Tezuka T., Yoshida Y., Yokoyama K., Umemori H., et al., p250GAP, a novel brain-enriched GTPase-activating protein for Rho family GTPases, is involved in the N-methyl-D-aspartate receptor signaling, *Mol. Biol. Cell*, 2003, 14, 2921-2934
- [113] Yu P., Zhang Y.P., Shields L.B., Zheng Y., Hu X., Hill R., et al., Inhibitor of DNA binding 2 promotes sensory axonal growth after SCI, *Exp. Neurol.*, 2011, 231, 38-44
- [114] Do J.L., Bonni A., Tuszynski M.H., SnoN facilitates axonal regeneration after spinal cord injury, *PLoS ONE*, 2013, 8, e71906