













- [14] Sirota A., Csicsvari J., Buhl D., Buzsáki G., Communication between neocortex and hippocampus during sleep in rodents, *Proc. Natl. Acad. Sci. USA*, 2003, 100, 2065-2069
- [15] Ji D., Wilson M.A., Coordinated memory replay in the visual cortex and hippocampus during sleep, *Nat. Neurosci.*, 2007, 10, 100-107
- [16] Peyrache A., Khamassi M., Benchenane K., Wiener S.I., Battaglia F.P., Replay of rule-learning related neural patterns in the prefrontal cortex during sleep, *Nat. Neurosci.*, 2009, 12, 919-926
- [17] Gauthier S., Scheltens P., Can we do better in developing new drugs for Alzheimer's disease?, *Alzheimers Dement.*, 2009, 5, 489-491
- [18] Bartus R.T., Dean R.L., Beer B., Lippa A.S., The cholinergic hypothesis of geriatric memory dysfunction, *Science*, 1982, 217, 408-417
- [19] Davies P., Maloney A.J.F., Selective loss of central cholinergic neurons in Alzheimer's disease, *Lancet*, 1976, 2, 1403-1403
- [20] Drachman D.A., Leavitt J., Human memory and cholinergic system - relationship to aging, *Arch. Neurol.*, 1974, 30, 113-121
- [21] Ellis B.W., Johns M.W., Lancaster R., Raptopoulos P., Angelopoulos N., Priest R.G., The St. Mary's Hospital sleep questionnaire: a study of reliability, *Sleep*, 1981, 4, 93-97
- [22] Bodick N.C., Offen W.W., Levey A.I., Cutler N.R., Gauthier S.G., Satlin A., et al., Effects of xanomeline, a selective muscarinic receptor agonist, on cognitive function and behavioral symptoms in Alzheimer disease, *Arch. Neurol.*, 1997, 54, 465-473
- [23] Levey A.I., Immunological localization of m1-m5 muscarinic acetylcholine-receptors in peripheral-tissues and brain, *Life Sci.*, 1993, 52, 441-448
- [24] Roldán G., Bolaños-Badillo E., González-Sánchez H., Quirarte G.L., Prado-Alcalá R.A., Selective M1 muscarinic receptor antagonists disrupt memory consolidation of inhibitory avoidance in rats, *Neurosci. Lett.*, 1997, 230, 93-96
- [25] Ferreira A.R., Fürstenau L., Blanco C., Kornisiuk E., Sánchez G., Daroit D., et al., Role of hippocampal M-1 and M-4 muscarinic receptor subtypes in memory consolidation in the rat, *Pharmacol. Biochem. Behav.*, 2003, 74, 411-415
- [26] Anagnostaras S.G., Murphy G.G., Hamilton S.E., Mitchell S.L., Rahnama N.P., Nathanson N.M., et al., Selective cognitive dysfunction in acetylcholine M-1 muscarinic receptor mutant mice, *Nat. Neurosci.*, 2003, 6, 51-58
- [27] Nissen C., Power A.E., Noftinger E.A., Feige B., Voderholzer U., Kloepfer C., et al., M-1 muscarinic acetylcholine receptor agonism alters sleep without affecting memory consolidation, *J. Cogn. Neurosci.*, 2006, 18, 1799-1807
- [28] Foster N.L., Aldrich M.S., Bluemlein L., White R.F., Berent S., Failure of cholinergic agonist RS-86 to improve cognition and movement in PSP despite effects on sleep, *Neurology*, 1989, 39, 257-261
- [29] Deuschl G., Schade-Brittinger C., Krack P., Volkmann J., Schäfer H., Bötzel K., et al., A randomized trial of deep-brain stimulation for Parkinson's disease, *N. Engl. J. Med.*, 2006, 355, 896-908
- [30] Huys D., Möller M., Kim E.H., Hardenacke K., Huff W., Klosterkötter J., et al., Deep brain stimulation for psychiatric disorders: historical basis, *Nervenarzt*, 2012, 83, 1156-1168
- [31] Laxton A.W., Tang-Wai D.F., McAndrews M.P., Zumsteg D., Wennberg R., Keren R., et al., A phase I trial of deep brain stimulation of memory circuits in Alzheimer's Disease, *Ann. Neurol.*, 2010, 68, 521-534
- [32] Suthana N., Haneef Z., Stern J., Mukamel R., Behnke E., Knowlton B., et al., Memory enhancement and deep-brain stimulation of the entorhinal area, *N. Engl. J. Med.*, 2012, 366, 502-510
- [33] Bethus I., Tse D., Morris R.G.M., Dopamine and memory: modulation of the persistence of memory for novel hippocampal NMDA receptor-dependent paired associates, *J. Neurosci.*, 2010, 30, 1610-1618
- [34] Chowdhury R., Guitart-Masip M., Bunzeck N., Dolan R.J., Düzel E., Dopamine modulates episodic memory persistence in old age, *J. Neurosci.*, 2012, 32, 14193-14204
- [35] Coulthard E.J., Bogacz R., Javed S., Mooney L.K., Murphy G., Keeley S., et al., Distinct roles of dopamine and subthalamic nucleus in learning and probabilistic decision making, *Brain*, 2012, 135, 3721-3734
- [36] Lisman J., Grace A.A., Düzel E., A neoHebbian framework for episodic memory; role of dopamine-dependent late LTP, *Trends Neurosci.*, 2011, 34, 536-547
- [37] Bernabeu R., Bevilacqua L., Ardenghi P., Bromberg E., Schmitz P., Bianchin M., et al., Involvement of hippocampal cAMP/cAMP-dependent protein kinase signaling pathways in a late memory consolidation phase of aversively motivated learning in rats, *Proc. Natl. Acad. Sci. USA*, 1997, 94, 7041-7046
- [38] Hannestad J., Gallezot J.-D., Planeta-Wilson B., Lin S.-F., Williams W.A., van Dyck C.H., et al., Clinically relevant doses of methylphenidate significantly occupy norepinephrine transporters in humans in vivo, *Biol. Psychiatry*, 2010, 68, 854-860
- [39] Volkow N.D., Fowler J.S., Wang G., Ding Y., Gatley S.J., Mechanism of action of methylphenidate: insights from PET imaging studies, *J. Atten. Disord.*, 2002, 6, Suppl. 1, S31-43
- [40] Andersen M.L., Kessler E., Murnane K.S., McClung J.C., Tufik S., Howell L.L., Dopamine transporter-related effects of modafinil in rhesus monkeys, *Psychopharmacology*, 2010, 210, 439-448
- [41] Outram S.M., The use of methylphenidate among students: the future of enhancement?, *J. Med. Ethics*, 2010, 36, 198-202
- [42] Turner D.C., Robbins T.W., Clark L., Aron A.R., Dowson J., Sahakian B.J., Cognitive enhancing effects of modafinil in healthy volunteers, *Psychopharmacology*, 2003, 165, 260-269
- [43] Linsen A.M.W., Vuurman E.F.P.M., Sambeth A., Riedel W.J., Methylphenidate produces selective enhancement of declarative memory consolidation in healthy volunteers, *Psychopharmacology*, 2012, 221, 611-619
- [44] Berridge C.W., Devilbiss D.M., Psychostimulants as cognitive Enhancers: the prefrontal cortex, catecholamines, and attention-deficit/hyperactivity disorder, *Biol. Psychiatry*, 2011, 69, E101-111
- [45] Pierard C., Liscia P., Chauveau F., Coutan M., Corio M., Krazem A., et al., Differential effects of total sleep deprivation on contextual and spatial memory: modulatory effects of modafinil, *Pharmacol. Biochem. Behav.*, 2011, 97, 399-405
- [46] Müller U., Rowe J.B., Rittman T., Lewis C., Robbins T.W., Sahakian B.J., Effects of modafinil on non-verbal cognition, task enjoyment and creative thinking in healthy volunteers, *Neuropharmacology*, 2013, 64, 490-495

- [47] Scammell T.E., Estabrooke I.V., McCarthy M.T., Chemelli R.M., Yanagisawa M., Miller M.S., et al., Hypothalamic arousal regions are activated during modafinil-induced wakefulness, *J. Neurosci.*, 2000, 20, 8620-8628
- [48] Meneses A., Ponce-Lopez T., Tellez R., Gonzalez R., Castillo C., Gasbarri A., Effects of d-amphetamine on short- and long-term memory in spontaneously hypertensive, Wistar-Kyoto and Sprague-Dawley rats, *Behav. Brain Res.*, 2011, 216, 472-476
- [49] Wiig K.A., Whitlock J.R., Epstein M.H., Carpenter R.L., Bear M.F., The levo enantiomer of amphetamine increases memory consolidation and gene expression in the hippocampus without producing locomotor stimulation, *Neurobiol. Learn. Mem.*, 2009, 92, 106-113
- [50] Ballard M.E., Gallo D.A., de Wit H., THC impairs, and amphetamine facilitates, memory encoding preferentially for emotionally salient stimuli, *Soc. Neurosci. Abstr.*, 2011, 41, 752.06
- [51] Advokat C., Scheithauer M., Attention-deficit hyperactivity disorder (ADHD) stimulant medications as cognitive enhancers, *Front. Neurosci.*, 2013, 7, 82
- [52] Sumowski J.F., Chiaravalloti N.D., Erlanger D.M., Kaushik T., Benedict R., DeLuca J., L-amphetamine improves memory capacity among memory-impaired patients with multiple sclerosis, *Neurology*, 2011, 76, Suppl. 4, A482
- [53] Jones S., Kornblum J.L., Kauer J.A., Amphetamine blocks long-term synaptic depression in the ventral tegmental area, *J. Neurosci.*, 2000, 20, 5575-5580
- [54] Del Arco A., González-Mora J.L., Armas V.R., Mora F., Amphetamine increases the extracellular concentration of glutamate in striatum of the awake rat: involvement of high affinity transporter mechanisms, *Neuropharmacology*, 1999, 38, 943-954
- [55] Drevets W.C., Gautier C., Price J.C., Kupfer D.J., Kinahan P.E., Grace A.A., et al., Amphetamine-induced dopamine release in human ventral striatum correlates with euphoria, *Biol. Psychiatry*, 2001, 49, 81-96
- [56] Imperato A., Obinu M.C., Gessa G.L., Effects of cocaine and amphetamine on acetylcholine-release in the hippocampus and caudate nucleus, *Eur. J. Pharmacol.*, 1993, 238, 377-381
- [57] Day J.C., Fibiger H.C., Dopaminergic regulation of septohippocampal cholinergic neurons, *J. Neurochem.*, 1994, 63, 2086-2092
- [58] Ponomarenko A.A., Lin J.S., Selbach O., Haas H.L., Temporal pattern of hippocampal high-frequency oscillations during sleep after stimulant-evoked waking, *Neuroscience*, 2003, 121, 759-769
- [59] Bekinschtein P., Cammarota M., Igaz L.M., Bevilacqua L.R.M., Izquierdo I., Medina J.H., Persistence of long-term memory storage requires a late protein synthesis- and BDNF-dependent phase in the hippocampus, *Neuron*, 2007, 53, 261-277
- [60] Rossato J.I., Bevilacqua L.R.M., Izquierdo I., Medina J.H., Cammarota M., Dopamine controls persistence of long-term memory storage, *Science*, 2009, 325, 1017-1020
- [61] Pencea V., Bingaman K.D., Wiegand S.J., Luskin M.B., Infusion of brain-derived neurotrophic factor into the lateral ventricle of the adult rat leads to new neurons in the parenchyma of the striatum, septum, thalamus, and hypothalamus, *J. Neurosci.*, 2001, 21, 6706-6717
- [62] Nagahara A.H., Merrill D.A., Coppola G., Tsukada S., Schroeder B.E., Shaked G.M., et al., Neuroprotective effects of brain-derived neurotrophic factor in rodent and primate models of Alzheimer's disease, *Nat. Med.*, 2009, 15, 331-337
- [63] Patel N.K., Gill S.S., GDNF delivery for Parkinson's disease, *Acta Neurochir. Suppl.*, 2007, 97, 135-154
- [64] Donlea J.M., Thimman M.S., Suzuki Y., Gottschalk L., Shaw P.J., Inducing sleep by remote control facilitates memory consolidation in *Drosophila*, *Science*, 2011, 332, 1571-1576
- [65] Mathias S., Zihl J., Steiger A., Lancel M., Effect of repeated gaboxadol administration on night sleep and next-day performance in healthy elderly subjects, *Neuropsychopharmacology*, 2005, 30, 833-841
- [66] Faulhaber J., Steiger A., Lancel M., The GABA(A) agonist THIP produces slow wave sleep and reduces spindling activity in NREM sleep in humans, *Psychopharmacology*, 1997, 130, 285-291
- [67] Boyle J., Wolford D., Gargano C., McCreary J., Cummings C., Cerchio K., et al., Next-day residual effects of gaboxadol and flurazepam administered at bedtime: a randomized double-blind study in healthy elderly subjects, *Hum. Psychopharmacol.*, 2009, 24, 61-71
- [68] Dijk D.J., James L.M., Peters S., Walsh J.K., Deacon S., Sex differences and the effect of gaboxadol and zolpidem on EEG power spectra in NREM and REM sleep, *J. Psychopharmacol.*, 2010, 24, 1613-1618
- [69] Massimini M., Ferrarelli F., Esser S.K., Riedner B.A., Huber R., Murphy M., et al., Triggering sleep slow waves by transcranial magnetic stimulation, *Proc. Natl. Acad. Sci. USA*, 2007, 104, 8496-8501
- [70] Marshall L., Helgadottir H., Moelle M., Born J., Boosting slow oscillations during sleep potentiates memory, *Nature*, 2006, 444, 610-613
- [71] Toni G., Riedner B.A., Hulse B.K., Ferrarelli F., Sarasso S., Enhancing sleep slow waves with natural stimuli, *Medicamundi*, 2010, 54, 73-79
- [72] Ngo H.-V.V., Martinetz T., Born J., Moelle M., Auditory closed-loop stimulation of the sleep slow oscillation enhances memory, *Neuron*, 2013, 78, 545-553
- [73] van Praag H., Christie B.R., Sejnowski T.J., Gage F.H., Running enhances neurogenesis, learning, and long-term potentiation in mice, *Proc. Natl. Acad. Sci. USA*, 1999, 96, 13427-13431
- [74] van Praag H., Neurogenesis and exercise: past and future directions, *Neuromolecular Med.*, 2008, 10, 128-140
- [75] Berggren K.L., Kerr A.L., Iles B.W., Nye S.H., Swain R.A., Exercise-induced angiogenesis in the CNS of Dahl Salt-Sensitive and SSBN.13 consomic rats, *Soc. Neurosci. Abstr.*, 2008, 38, 219.10
- [76] Chaddock L., Erickson K.I., Prakash R.S., Kim J.S., Voss M.W., VanPatter M., et al., A neuroimaging investigation of the association between aerobic fitness, hippocampal volume, and memory performance in preadolescent children, *Brain Res.*, 2010, 1358, 172-183
- [77] Erickson K.I., Voss M.W., Prakash R.S., Basak C., Szabo A., Chaddock L., et al., Exercise training increases size of hippocampus and improves memory, *Proc. Natl. Acad. Sci. USA*, 2011, 108, 3017-3022
- [78] Coles K., Tomporowski P.D., Effects of acute exercise on executive processing, short-term and long-term memory, *J. Sports Sci.*, 2008, 26, 333-344



- [79] Gould E., Reeves A.J., Fallah M., Tanapat P., Gross C.G., Fuchs E., Hippocampal neurogenesis in adult Old World primates, *Proc. Nat. Acad. Sci. USA*, 1999, 96, 5263-5267
- [80] Amrein I., Isler K., Lipp H.-P., Comparing adult hippocampal neurogenesis in mammalian species and orders: influence of chronological age and life history stage, *Eur. J. Neurosci.*, 2011, 34, 978-987
- [81] Apostolova L.G., Green A.E., Babakchanian S., Hwang K.S., Chou Y.-Y., Toga A.W., et al., Hippocampal atrophy and ventricular enlargement in normal aging, mild cognitive impairment (MCI), and Alzheimer disease, *Alzheimer Dis. Assoc. Disord.*, 2012, 26, 17-27
- [82] Garcia-Mesa Y., Carlos Lopez-Ramos J., Gimenez-Llort L., Revilla S., Guerra R., Gruart A., et al., Physical exercise protects against Alzheimer's disease in 3xTg-AD mice, *J. Alzheimers Dis.*, 2011, 24, 421-454
- [83] Rodriguez J.J., Noristani H.N., Olabarria M., Fletcher J., Somerville T.D.D., Yeh C.Y., et al., Voluntary running and environmental enrichment restores impaired hippocampal neurogenesis in a triple transgenic mouse model of Alzheimer's disease, *Curr. Alzheimer Res.*, 2011, 8, 707-717
- [84] Nagamatsu L.S., Chan A., Davis J.C., Beattie J.C., Beattie B.L., Graf P., et al., Physical activity improves verbal and spatial memory in older adults with probable mild cognitive impairment: a 6-month randomized controlled trial, *J. Aging Res.*, 2013, 861893
- [85] Lautenschlager N.T., Cox K.L., Flicker L., Foster J.K., van Bockxmeer F.M., Xiao J., et al., Effect of physical activity on cognitive function in older adults at risk for Alzheimer disease: a randomized trial, *JAMA*, 2008, 300, 1027-1037
- [86] Cassilhas R.C., Lee K.S., Fernandes J., Oliveira M.G.M., Tufik S., Meeusen R., et al., Spatial memory is improved by aerobic and resistance exercise through divergent molecular mechanisms, *Neuroscience*, 2012, 202, 309-317
- [87] Chen D.Y., Stern S.A., Garcia-Osta A., Saunier-Rebori B., Pollonini G., Bambah-Mukku D., et al., A critical role for IGF-II in memory consolidation and enhancement, *Nature*, 2011, 469, 491-497
- [88] Dash M.B., Bellesi M., Tononi G., Cirelli C., Sleep/wake dependent changes in cortical glucose concentrations, *J. Neurochem.*, 2013, 124, 79-89
- [89] Jones E.K., Suenram-Lea S.I., Wesnes K.A., Acute ingestion of different macronutrients differentially enhances aspects of memory and attention in healthy young adults, *Biol. Psychol.*, 2012, 89, 477-486
- [90] Messier C., Glucose improvement of memory: a review, *Eur. J. Pharmacol.*, 2004, 490, 33-57
- [91] Smith M.A., Riby L.M., van Eekelen J.A.M., Foster J.K., Glucose enhancement of human memory: a comprehensive research review of the glucose memory facilitation effect, *Neurosci. Biobehav. Rev.*, 2011, 35, 770-783
- [92] Sunram-Lea S.I., Foster J.K., Durlach P., Perez C., Glucose facilitation of cognitive performance in healthy young adults: examination of the influence of fast-duration, time of day and pre-consumption plasma glucose levels, *Psychopharmacology*, 2001, 157, 46-54
- [93] Sunram-Lea S.I., Foster J.K., Durlach P., Perez C., Investigation into the significance of task difficulty and divided allocation of resources on the glucose memory facilitation effect, *Psychopharmacology*, 2002, 160, 387-397
- [94] Sunram-Lea S.I., Foster J.K., Durlach P., Perez C., The effect of retrograde and anterograde glucose administration on memory performance in healthy young adults, *Behav. Brain Res.*, 2002, 134, 505-516
- [95] Varady K.A., Hellerstein M.K., Alternate-day fasting and chronic disease prevention: a review of human and animal trials, *Am. J. Clin. Nutr.*, 2007, 86, 7-13
- [96] Partridge L., Piper M.D.W., Mair W., Dietary restriction in *Drosophila*, *Mech. Ageing Dev.*, 2005, 126, 938-950
- [97] Mattison J.A., Roth G.S., Beasley T.M., Tilmont E.M., Handy A.M., Herbert R.L., et al., Impact of caloric restriction on health and survival in rhesus monkeys from the NIA study, *Nature*, 2012, 489, 318-321
- [98] Weindruch R., Walford R.L., Fligiel S., Guthrie D., The retardation of aging in mice by dietary restriction: longevity, cancer, immunity and lifetime energy-intake, *J. Nutr.*, 1986, 116, 641-654
- [99] Colman R.J., Anderson R.M., Johnson S.C., Kastman E.K., Kosmatka K.J., Beasley T.M., et al., Caloric restriction delays disease onset and mortality in rhesus monkeys, *Science*, 2009, 325, 201-204
- [100] Eckles-Smith K., Clayton D., Bickford P., Browning M.D., Caloric restriction prevents age-related deficits in LTP and in NMDA receptor expression, *Mol. Brain Res.*, 2000, 78, 154-162
- [101] Singh R., Lakhanpal D., Kumar S., Sharma S., Kataria H., Kaur M., et al., Late-onset intermittent fasting dietary restriction as a potential intervention to retard age-associated brain function impairments in male rats, *Age*, 2012, 34, 917-933
- [102] Qin W., Chachich M., Lane M., Roth G., Bryant M., de Cabo R., et al., Calorie restriction attenuates Alzheimer's disease type brain amyloidosis in Squirrel monkeys (*Saimiri sciureus*), *J. Alzheimers Dis.*, 2006, 10, 417-422
- [103] Wu P., Shen Q., Dong S., Xu Z., Tsien J.Z., Hu Y., Calorie restriction ameliorates neurodegenerative phenotypes in forebrain-specific presenilin-1 and presenilin-2 double knockout mice, *Neurobiol. Aging*, 2008, 29, 1502-1511
- [104] Thaler S., Choragiewicz T.J., Rejda R., Fiedorowicz M., Turski W.A., Tulidowicz-Bielak M., et al., Neuroprotection by acetoacetate and beta-hydroxybutyrate against NMDA-induced RGC damage in rat-possible involvement of kynurenic acid, *Graefes Arch. Clin. Exp. Ophthalmol.*, 2010, 248, 1729-1735
- [105] Henderson S.T., Ketone bodies as a therapeutic for Alzheimer's disease, *Neurotherapeutics*, 2008, 5, 470-480
- [106] Reger M.A., Henderson S.T., Hale C., Cholerton B., Baker L.D., Watson G.S., et al., Effects of beta-hydroxybutyrate on cognition in memory-impaired adults, *Neurobiol. Aging*, 2004, 25, 311-314
- [107] Dhurandhar E.J., Allison D.B., van Groen T., Kadish I., Hunger in the absence of caloric restriction improves cognition and attenuates Alzheimer's disease pathology in a mouse model, *PLoS One*, 2013, 8, e60437

- [108] Cao D.H., Kevala K., Kim J., Moon H.S., Jun S.B., Lovinger D., et al., Docosahexaenoic acid promotes hippocampal neuronal development and synaptic function, *J. Neurochem.*, 2009, 111, 510-521
- [109] Bhatia H.S., Agrawal R., Sharma S., Huo Y.-X., Ying Z., Gomez-Pinilla F., Omega-3 fatty acid deficiency during brain maturation reduces neuronal and behavioral plasticity in adulthood, *PLoS One*, 2011, 6, e28451
- [110] Vines A., Delattre A.M., Lima M.M.S., Rodrigues L.S., Suchecki D., Machado R.B., et al., The role of 5-HT1A receptors in fish oil-mediated increased BDNF expression in the rat hippocampus and cortex: a possible antidepressant mechanism, *Neuropharmacology*, 2012, 62, 184-191
- [111] Arsenault D., Julien C., Tremblay C., Calon F., DHA improves cognition and prevents dysfunction of entorhinal cortex neurons in 3xTg-AD mice, *PLoS One*, 2011, 6, e17397