

## RESEARCH PAPER

NONSTANDARD GAUSS–LOBATTO QUADRATURE  
APPROXIMATION TO FRACTIONAL DERIVATIVESShahrokh Esmaeili <sup>1</sup>, Gradimir V. Milovanović <sup>2</sup>

*Dedicated to Professor Ivan Dimovski  
on the occasion of his 80th anniversary*

**Abstract**

A family of nonstandard Gauss-Jacobi-Lobatto quadratures for numerical calculating integrals of the form  $\int_{-1}^1 f'(x)(1-x)^\alpha dx$ ,  $\alpha > -1$ , is derived and applied to approximation of the usual fractional derivative. A software implementation of such quadratures was done by the recent MATHEMATICA package `OrthogonalPolynomials` (cf. [A.S. Cvetković, G.V. Milovanović, Facta Univ. Ser. Math. Inform. **19** (2004), 17–36] and [G.V. Milovanović, A.S. Cvetković, Math. Balkanica **26** (2012), 169–184]). Several numerical examples are presented and they show the effectiveness of the proposed approach.

*MSC 2010:* Primary 65D30; Secondary 33C45, 41A55, 65D32

*Key Words and Phrases:* fractional derivative, quadrature rule, Gaussian quadrature, nodes, weights, software implementation, MATHEMATICA package

**1. Introduction**

Fractional calculus (including the operators of fractional order integration and differentiation) has been often used in modelling many physical and engineering problems. We refer the reader to [2, 3, 21, 22, 25, 36, 38] for

a general theory, as well as a description of the main applications in fields ranging from mechanics, biology to biomechanics, diagnostic imaging, electrochemistry, finance, sustainable environment and renewable energy. A theory of the generalized fractional calculus, as well as the corresponding applications, can be found in a monograph written by Kiryakova [23]. We also mention a few the most cited papers on this subject [17, 26, 27, 5].

The fractional derivative  $D_*^q f(t)$  in the Caputo version [6] and the Riemann–Liouville fractional derivative  ${}_0D_t^q f(t)$ , where  $0 < q < 1$ , are defined by

$$D_*^q f(t) = \frac{1}{\Gamma(1-q)} \int_0^t f'(s)(t-s)^{-q} ds, \quad t > 0, \quad (1.1)$$

$${}_0D_t^q f(t) = \frac{1}{\Gamma(1-q)} \frac{d}{dt} \int_0^t f(s)(t-s)^{-q} ds, \quad t > 0, \quad (1.2)$$

respectively [36]. It is well-known that the fractional derivative of Riemann–Liouville and Caputo type are closely linked by the following relationship:

$${}_0D_t^q f(t) = \frac{f(0)}{t^q \Gamma(1-q)} + D_*^q f(t). \quad (1.3)$$

The operator of fractional derivative is more complicated than the classical one, and its calculation is also more difficult than in the integer order case.

In this paper we construct a quadrature method for approximating the fractional derivative defined by (1.1) or (1.2). Although there is comprehensive literature on the numerical methods for solving equations involving fractional derivatives and integrals (cf. [8, 9, 11, 3, 10, 35]), there seems to exist a few literature on automatic quadrature for the fractional derivatives, see e.g. [20, 24, 39].

Let  $w(x)$  be a nonnegative weight function on the interval  $(a, b)$  that vanishes only at isolated points. We consider an  $n$ -point quadrature formula

$$\int_a^b f(x)w(x) dx = \sum_{k=1}^n A_k f(x_k) + R_n[f], \quad (1.4)$$

where the sum  $Q_n[f] = \sum_{k=1}^n A_k f(x_k)$  provides an approximation to the integral  $\int_a^b f(x)w(x) dx$  and  $R_n[f]$  is the corresponding error. We shall always require  $a \leq x_1 < x_2 < \dots < x_n \leq b$ . In the quadrature sum  $Q_n[f]$ , the points  $x_k$  are called the nodes and  $A_k$  are the weights of the quadrature formula (1.4). The mentioned quadrature rules use the information on the integrand only at some selected points  $x_k$ ,  $k = 1, \dots, n$  (the values of the function  $f$ ). Such quadratures will be called the standard quadrature formulae, see [4, 28, 40].

The quadrature formula (1.4) with the maximal algebraic degree of exactness  $2n - 1$  is called the Gaussian quadrature formula. Indeed, if  $\mathbb{P}_d$  be the set of all algebraic polynomials of degree at most  $d$  then in the Gaussian quadrature formula (1.4) we have  $R_n[p] = 0$  for all  $p \in \mathbb{P}_{2n-1}$ .

This paper is organized as follows. In Section 2 we give a short account on Gauss–Jacobi quadrature rules and the corresponding Lobatto modification. Some remarks on nonstandard Gaussian quadratures are mentioned in Section 4. Main result on the Gauss–Jacobi–Lobatto nonstandard quadrature is proved in Section 4, as well as the corresponding software implementation in MATHEMATICA. Finally, applications to an approximation of fractional derivatives and numerical examples are presented in Sections 5 and 6, respectively. Several numerical examples show the effectiveness of the proposed approach.

## 2. Gauss–Jacobi quadrature rules and the Lobatto modification

Let  $w^{\alpha,\beta}(x) := (1-x)^\alpha(1+x)^\beta$  with parameters  $\alpha, \beta > -1$  be the Jacobi weight function on the interval  $(-1, 1)$ . The Jacobi polynomials, denoted by  $P_n^{(\alpha,\beta)}(x)$ , are mutually orthogonal on the interval  $(-1, 1)$  with respect to the  $w^{\alpha,\beta}(x)$ . This means that

$$\int_{-1}^1 P_n^{(\alpha,\beta)}(x) P_m^{(\alpha,\beta)}(x) w^{\alpha,\beta}(x) dx = 0 \quad n \neq m. \quad (2.1)$$

An  $n$ -point quadrature rule for the weight function  $w^{\alpha,\beta}$  is called a formula of the type

$$\int_{-1}^1 f(x) w^{\alpha,\beta}(x) dx = \sum_{k=1}^n w_k^{\alpha,\beta} f(x_k^{\alpha,\beta}) + R_n[f]. \quad (2.2)$$

A quadrature rule (2.2) with degree of exactness  $n-1$  is called interpolatory. These are precisely those obtained by interpolation, that is, for which

$$w_j^{\alpha,\beta} = \int_{-1}^1 \ell_j(x) w^{\alpha,\beta}(x) dx, \quad j = 1, \dots, n, \quad (2.3)$$

where  $\ell_j \in \mathbb{P}_{n-1}$  are the Lagrange polynomials associated with the nodes  $x_k^{\alpha,\beta}$  such that  $\ell_j(x_k^{\alpha,\beta}) = \delta_{jk}$ , for  $j, k = 1, \dots, n$ .

We now discuss the relation between the Jacobi polynomials and the Gauss–type quadratures. The mechanism of a Gauss–type quadrature is to seek the best numerical approximation of an integral by selecting optimal nodes at which integrand is evaluated. It can be proved that the nodes  $x_k^{\alpha,\beta}$  in a Gauss–Jacobi quadrature are the roots of the Jacobi polynomial  $P_n^{(\alpha,\beta)}$ , that is

$$P_n^{(\alpha,\beta)}(x_k^{\alpha,\beta}) = 0, \quad k = 1, \dots, n. \quad (2.4)$$

All nodes  $x_k^{\alpha,\beta}$  are real, distinct, and contained in the open interval  $(-1, 1)$ . The weights  $w_k^{\alpha,\beta}$  are known in closed form as

$$w_k^{\alpha,\beta} = \frac{2^{\alpha+\beta+1}}{n!\Gamma(n+\alpha+\beta+1)} \cdot \frac{\Gamma(n+\alpha+1)\Gamma(n+\beta+1)}{\left(1 - (x_k^{\alpha,\beta})^2\right) \left[\frac{d}{dx}P_n^{(\alpha,\beta)}(x_k^{\alpha,\beta})\right]^2}, \quad (2.5)$$

where  $k = 1, \dots, n$  (see, for instance, [28, p. 325]). We can thus conclude that the quadrature formula (2.2) with the nodes  $x_k^{\alpha,\beta}$  prescribed by (2.4) and the weights  $w_k^{\alpha,\beta}$  given by (2.3), has degree of exactness  $2n-1$ , the maximum value that can be achieved using interpolatory quadrature formulae with  $n$  nodes. This optimal formula is called the (standard) Gauss–Jacobi quadrature formula.

As  $n$  increases, finding roots of  $P_n^{(\alpha,\beta)}$  become an ill-conditioned and time consuming problem. An alternative and today standard approach for finding the nodes and the weights of a Gaussian quadrature rule is to use the Golub–Welsch algorithm [15, 14].

Let  $k_n^{\alpha,\beta}$  be the leading coefficient of  $P_n^{(\alpha,\beta)}(x)$ , i.e.,

$$k_n^{\alpha,\beta} = \frac{(n+\alpha+\beta+1)_n}{2^n n!}.$$

It is well known that there exists a unique sequence of monic orthogonal polynomials  $\pi_n^{\alpha,\beta}(x) = P_n^{(\alpha,\beta)}(x)/k_n^{\alpha,\beta}$ . They can be generated by the three term recurrence relation [14, 28]

$$\pi_{k+1}^{\alpha,\beta}(x) = (x - \alpha_k^J)\pi_k^{\alpha,\beta}(x) - \beta_k^J\pi_{k-1}^{\alpha,\beta}(x), \quad k \geq 0, \quad (2.6)$$

with starting values  $\pi_{-1}^{\alpha,\beta} = 0$ ,  $\pi_0^{\alpha,\beta} = 1$ , and with the recursion coefficients

$$\alpha_k^J = \frac{\beta^2 - \alpha^2}{(2k + \alpha + \beta)(2k + \alpha + \beta + 2)}, \quad k \geq 0,$$

$$\beta_k^J = \frac{4k(k + \alpha)(k + \beta)(k + \alpha + \beta)}{(2k + \alpha + \beta)^2((2k + \alpha + \beta)^2 - 1)}, \quad k \geq 1.$$

This approach is based on determining the eigenvalues and normalized eigenvectors of the symmetric tridiagonal *Jacobi matrix*

$$\mathbf{J}_n = \text{tridiag}(\boldsymbol{\beta}^J, \boldsymbol{\alpha}^J, \boldsymbol{\beta}^J) \in \mathbb{R}^{n \times n}, \quad (2.7)$$

where

$$\boldsymbol{\alpha}^J = (\alpha_0^J, \alpha_1^J, \dots, \alpha_{n-1}^J)^T \in \mathbb{R}^n, \quad \boldsymbol{\beta}^J = \left(\sqrt{\beta_1^J}, \dots, \sqrt{\beta_{n-1}^J}\right)^T \in \mathbb{R}^{n-1}.$$

The elements of this matrix are obtained from the coefficients of the three-term recurrence relation (2.6). The nodes  $x_k^{\alpha,\beta}$  in (2.2) are the eigenvalues

of the matrix  $\mathbf{J}_n$  and the corresponding weights  $w_k^{\alpha,\beta}$  may be obtained from the first components of the normalized eigenvectors. For more details we refer the reader to [14, 15, 28]. An efficient algorithm for computation of Gauss–Jacobi quadrature nodes and weights for any  $n \geq 100$  has been recently presented in [18].

The  $(n+2)$ -points Gauss–Lobatto quadrature rule for the Jacobi weight function is an interpolatory quadrature rule of the following form

$$\int_{-1}^1 f(x)w^{\alpha,\beta}(x) dx = \lambda_0^{\alpha,\beta} f(-1) + \sum_{k=1}^n \lambda_k^{\alpha,\beta} f(\zeta_k^{\alpha,\beta}) + \lambda_{n+1}^{\alpha,\beta} f(1) + R_n^{\alpha,\beta}[f], \tag{2.8}$$

with the maximal algebraic degree of exactness. The internal nodes  $\zeta_k^{\alpha,\beta}$  are the zeros of the orthogonal polynomial of degree  $n$  with respect to the weight function  $x \mapsto (1-x^2)w^{\alpha,\beta}(x)$  (cf. [28, pp. 330–332]). In this case, the Gauss–Lobatto points  $\zeta_k^{\alpha,\beta}$  are the zeros of the polynomial  $\frac{d}{dx}P_{n+1}^{(\alpha,\beta)}(x)$ . It is well known that the Gauss–Lobatto quadrature (2.8) has degree of exactness  $2n + 1$ , see [14, 28].

### 3. Nonstandard quadratures of Gaussian type

In many cases it is not possible to measure the exact value of the function  $f$  at points  $x_k$ , so that a standard quadrature cannot be applied. Thus, if the information data  $\{f(x_k)\}_{k=1}^n$  in the standard quadrature (1.4) is replaced by  $\{\mathcal{A}^{h_k} f(x_k)\}_{k=1}^n$ , where  $\mathcal{A}^h$  is an extension of some linear operator  $\mathcal{A}^h : \mathbb{P} \rightarrow \mathbb{P}$ ,  $h \geq 0$ , where  $\mathbb{P}$  is the set of all algebraic polynomials, we get a non-standard quadrature formula [33]

$$\int_a^b f(x)w(x) dx = \sum_{k=1}^n w_k(\mathcal{A}^{h_k} f)(x_k) + R_m[f], \tag{3.1}$$

where  $w$  is a positive weight function on  $[a, b]$ .

In the sequel, we consider a non-standard  $m$ -point ( $m > 2$ ) interpolatory quadrature of Gaussian type (precisely, of the Gauss-Lobatto type),

$$\int_a^b g'(x)w(x) dx = \sum_{k=1}^m w_k g(x_k) + R_m[g], \tag{3.2}$$

which is exact on the set  $\mathbb{P}_{2m-3}$ , i.e.,  $R_m(\mathbb{P}_{2m-3}) = 0$ . If we put  $g(x) = \int_c^x f(t) dt$ , where  $c$  is some constant, formula (3.2) reduces to

$$\int_a^b f(x)w(x) dx = \sum_{k=1}^m w_k \int_c^{x_k} f(t) dt + R_m[f],$$

which is a special case of the generalized nonstandard (operator) formulae (3.2) of Gaussian type.

**THEOREM 3.1.** *Given an integer  $k$  with  $0 \leq k \leq m$ , the quadrature formula (3.2) has degree of exactness  $d = m - 1 + k$  if and only if both the following conditions are satisfied:*

- (a) *Formula (3.2) is interpolatory.*
- (b) *The node polynomial  $\ell(x) := \prod_{\nu=1}^m (x - x_\nu)$  satisfies*

$$\int_a^b (\ell(x)p(x))' w(x) dx = 0, \quad (3.3)$$

for all  $p \in \mathbb{P}_{k-1}$ .

**P r o o f.** To prove this, we will follow a manner similar to that used in [14]. We first prove the necessity of (a) and (b). Since, by assumption, the degree of exactness is  $d = m - 1 + k \geq m - 1$ , condition (a) is trivial. condition (b) also follows immediately, since, for any  $p \in \mathbb{P}_{k-1}$ , the product  $\ell(x)p(x)$  is in  $\mathbb{P}_{m-1+k}$ ; hence,

$$\int_a^b (\ell(x)p(x))' w(x) dx = \sum_{\nu=1}^m w_\nu \ell(x_\nu) p(x_\nu) = 0.$$

To prove the sufficiency of (a) and (b), we must show that for any  $p \in \mathbb{P}_{m-1+k}$  we have  $R_m[p] = 0$  in (3.2). Given any such  $p$ , divide it by  $\ell$ , so that

$$p(x) = q(x)\ell(x) + r(x), \quad q \in \mathbb{P}_{k-1}, \quad r \in \mathbb{P}_{m-1},$$

where  $q(x)$  is the quotient and  $r(x)$  is the corresponding remainder. There follows

$$\int_a^b p'(x)w(x) dx = \int_a^b (\ell(x)q(x))' w(x) dx + \int_a^b r'(x)w(x) dx$$

The first integral on the right vanishes by (b), since  $q \in \mathbb{P}_{k-1}$ , whereas the second, by (a), since  $r \in \mathbb{P}_{m-1}$ , equals

$$\sum_{\nu=1}^m w_\nu r(x_\nu)$$

by virtue of (a). But

$$r(x_\nu) = p(x_\nu) - q(x_\nu)\ell(x_\nu) = p(x_\nu),$$

so that indeed

$$\int_a^b p'(x)w(x)dx = \sum_{\nu=1}^m w_\nu p(x_\nu)$$

that is,  $R_n[p] = 0$ . This completes the proof. □

#### 4. Gauss-Jacobi-Lobatto nonstandard quadratures

As an interesting special case for application to fractional derivatives, we consider the nonstandard Gauss-Lobatto quadrature rule, with respect to a special Jacobi weight  $w^{\alpha,0}(x) = (1-x)^\alpha$ ,  $\alpha > -1$ ,

$$\int_{-1}^1 g'(x)w^{\alpha,0}(x) dx = \lambda_0^{\alpha,0}g(-1) + \sum_{k=1}^n \lambda_k^{\alpha,0}g(\xi_k^{\alpha,1}) + \lambda_{n+1}^{\alpha,0}g(1) + R_n^{\alpha,0}[g]. \tag{4.1}$$

Our main result can be stated in the following form:

**THEOREM 4.1.** *Let  $-1 = \xi_0^{\alpha,1} < \xi_1^{\alpha,1} < \dots < \xi_n^{\alpha,1} < \xi_{n+1}^{\alpha,1} = 1$  be the zeros of the polynomial  $(1-x^2)P_n^{(\alpha,1)}(x)$ , and let the weights  $\lambda_k^{\alpha,0}$  be as*

$$\lambda_k^{\alpha,0} = \frac{\alpha w_k^{\alpha,1}}{1 - (\xi_k^{\alpha,1})^2}, \quad k = 1, \dots, n, \tag{4.2}$$

and

$$\lambda_0^{\alpha,0} = -2^\alpha \frac{n^2 + (\alpha + 2)n + 1}{(n + 1)(n + \alpha + 1)}, \quad \lambda_{n+1}^{\alpha,0} = -\lambda_0^{\alpha,0} - \sum_{k=1}^n \lambda_k^{\alpha,0}, \tag{4.3}$$

where  $w_k^{\alpha,1}$ ,  $k = 1, \dots, n$ , are the Christoffel numbers in the standard Gauss-Jacobi quadrature with respect to the weight function  $w^{\alpha,1}(x) = (1-x)^\alpha(1+x)$ ,  $\alpha > -1$ , which correspond to the nodes  $\xi_k^{\alpha,1}$ ,  $k = 1, \dots, n$ .

Then the quadrature formula (4.1) has the maximal degree of precision, i.e., it is exact for each polynomial of degree at most  $2n + 1$ .

**P r o o f.** Suppose that  $\omega(x) = (1-x^2)P_n^{(\alpha,1)}(x)$  and

$$\lambda_k^{\alpha,0} = \int_{-1}^1 \ell_k'(x)w^{\alpha,0}(x) dx, \quad k = 0, 1, \dots, n + 1, \tag{4.4}$$

where  $\ell_k \in \mathbb{P}_{n+1}$  are the Lagrange polynomials associated with the nodes  $\xi_k^{\alpha,1}$ . It is obvious that the quadrature (4.1) with these weights (4.4) is at least interpolatory. According to Theorem 3.1, it is enough to show the orthogonality relation (3.3) for each  $p \in \mathbb{P}_{n-1}$ . To do this, we apply integration by parts to obtain

$$\int_{-1}^1 (\omega(x)p(x))'w^{\alpha,0}(x) dx = \alpha \int_{-1}^1 P_n^{(\alpha,1)}(x)p(x)w^{\alpha,1}(x) dx = \alpha \langle P_n^{(\alpha,1)}, p \rangle.$$

Then, because of orthogonality of  $P_n^{(\alpha,1)}$  to  $\mathbb{P}_{n-1}$  with respect to  $w^{\alpha,1}$  on  $(-1, 1)$ , we conclude that

$$(\forall p \in \mathbb{P}_{n-1}) \int_{-1}^1 (\omega(x)p(x))' w^{\alpha,0}(x) dx = \alpha \langle P_n^{(\alpha,1)}, p \rangle = 0.$$

Now, we denote the “node” polynomial by  $\ell(x) = (x^2 - 1)\pi_n(x)$ , where  $\pi_n \equiv \pi_n^{\alpha,1}$  is the monic Jacobi polynomial with parameters  $\alpha$  and  $\beta = 1$ .

In order to compute the weights  $\lambda_k^{\alpha,0}$ , corresponding to the set of nodes  $\xi_k^{\alpha,1}$ , we use the formulas (4.4).

Since  $\ell'(x) = 2x\pi_n(x) + (x^2 - 1)\pi_n'(x)$ , for the endpoints  $\pm 1$ , we have

$$\ell'(\xi_0^{\alpha,1}) = \ell'(-1) = -2\pi_n(-1), \quad \ell'(\xi_{n+1}^{\alpha,1}) = \ell'(1) = 2\pi_n(1).$$

At the zeros of the Jacobi polynomial  $\pi_n$ , we have

$$\ell'(\xi_k^{\alpha,1}) = ((\xi_k^{\alpha,1})^2 - 1)\pi_n'(\xi_k^{\alpha,1}), \quad k = 1, \dots, n.$$

The corresponding Lagrange polynomials associated with the nodes  $\xi_k^{\alpha,1}$ ,

$$\ell_k(x) = \frac{\ell(x)}{\ell'(\xi_k^{\alpha,1})(x - \xi_k^{\alpha,1})}, \quad k = 0, 1, \dots, n+1, \quad (4.5)$$

can be expressed in the following form

$$\ell_0(x) = \frac{(1-x)\pi_n(x)}{2\pi_n(-1)}, \quad \ell_{n+1}(x) = \frac{(1+x)\pi_n(x)}{2\pi_n(1)}$$

and

$$\ell_k(x) = \frac{1-x^2}{1-(\xi_k^{\alpha,1})^2} \cdot \frac{\pi_n(x)}{\pi_n'(\xi_k^{\alpha,1})(x - \xi_k^{\alpha,1})}, \quad k = 1, \dots, n.$$

Now we need derivatives of the Lagrange polynomials, i.e.,

$$\ell'_0(x) = \frac{(1-x)\pi_n'(x) - \pi_n(x)}{2\pi_n(-1)}, \quad \ell'_{n+1}(x) = \frac{(1+x)\pi_n'(x) + \pi_n(x)}{2\pi_n(1)}$$

and

$$\ell'_k(x) = \frac{1}{1-(\xi_k^{\alpha,1})^2} \left\{ -2x \ell_k^G(x) + (1-x^2) \frac{d}{dx} [\ell_k^G(x)] \right\}, \quad k = 1, \dots, n,$$

where  $\ell_k^G(x)$ ,  $k = 1, \dots, n$ , are Lagrange polynomials associated with the nodes  $\xi_k^{\alpha,1}$  for which

$$\int_{-1}^1 \ell_k^G(x) w^{\alpha,1}(x) dx = w_k^{\alpha,1}, \quad k = 1, \dots, n,$$

where  $w_k^{\alpha,1}$  are the Christoffel numbers in the corresponding Gauss-Jacobi quadrature rule with respect to the weight  $w^{\alpha,1}$  on  $(-1, 1)$ .

Thus, for  $k = 1, \dots, n$ , we get



$$\begin{aligned} \lambda_k^{\alpha,0} &= \int_{-1}^1 \ell'_k(x) w^{\alpha,0}(x) dx \\ &= \frac{1}{1 - (\xi_k^{\alpha,1})^2} \left\{ \int_{-1}^1 [(1-x) - (1+x)] \ell_k^G(x) w^{\alpha,0}(x) dx \right. \\ &\quad \left. + \int_{-1}^1 (1-x^2) \frac{d}{dx} [\ell_k^G(x)] w^{\alpha,0}(x) dx \right\} \\ &= \frac{1}{1 - (\xi_k^{\alpha,1})^2} \left\{ \int_{-1}^1 \ell_k^G(x) w^{\alpha+1,0}(x) dx - \int_{-1}^1 \ell_k^G(x) w^{\alpha,1}(x) dx \right. \\ &\quad \left. + \int_{-1}^1 \frac{d}{dx} [\ell_k^G(x)] w^{\alpha+1,1}(x) dx \right\}. \end{aligned}$$

Using integration by parts, the last integral becomes

$$\begin{aligned} \int_{-1}^1 \frac{d}{dx} [\ell_k^G(x)] w^{\alpha+1,1}(x) dx &= (\alpha + 1) \int_{-1}^1 \ell_k^G(x) w^{\alpha,1}(x) dx \\ &\quad - \int_{-1}^1 \ell_k^G(x) w^{\alpha+1,0}(x) dx, \end{aligned}$$

so that we get

$$\lambda_k^{\alpha,0} = \frac{\alpha}{1 - (\xi_k^{\alpha,1})^2} \int_{-1}^1 \ell_k^G(x) w^{\alpha,1}(x) dx = \frac{\alpha w_k^{\alpha,1}}{1 - (\xi_k^{\alpha,1})^2}, \quad k = 1, \dots, n,$$

i.e., (4.2).

For  $k = 0$  we have

$$\begin{aligned} \lambda_0^{\alpha,0} &= \int_{-1}^1 \ell'_0(x) w^{\alpha,0}(x) dx \\ &= \frac{1}{2\pi_n(-1)} \int_{-1}^1 [(1-x)\pi'_n(x) - \pi_n(x)] w^{\alpha,0}(x) dx \\ &= \frac{1}{2\pi_n(-1)} \left\{ \int_{-1}^1 \pi'_n(x) w^{\alpha+1,0}(x) dx - \int_{-1}^1 \pi_n(x) w^{\alpha,0}(x) dx \right\}. \end{aligned}$$

Since

$$\int_{-1}^1 \pi'_n(x) w^{\alpha+1,0}(x) dx = -\pi_n(-1)2^{\alpha+1} + (\alpha + 1) \int_{-1}^1 \pi_n(x) w^{\alpha,0}(x) dx,$$

we get

$$\lambda_0^{\alpha,0} = \frac{1}{2\pi_n(-1)} \left\{ \alpha \int_{-1}^1 \pi_n(x) w^{\alpha,0}(x) dx - 2^{\alpha+1} \pi_n(-1) \right\}.$$

If we introduce the so-called function of the second kind (cf. [28, p. 117])

$$f_n(z) = \int_{-1}^1 \frac{\pi_n(x)}{z-x} w^{\alpha,1}(x) dx$$

we see that for  $z = -1$ , it reduces to

$$f_n(-1) = - \int_{-1}^1 \frac{\pi_n(x)}{1+x} w^{\alpha,1}(x) dx = - \int_{-1}^1 \pi_n(x) w^{\alpha,0}(x) dx,$$

so that

$$\lambda_0^{\alpha,0} = -\frac{\alpha}{2} \cdot \frac{f_n(-1)}{\pi_n(-1)} - 2^\alpha. \quad (4.6)$$

In order to determine  $f_n(-1)$  we start with the following equality for Jacobi polynomials (cf. [1, p. 304])

$$(2n + \alpha + 1)P_n^{(\alpha,0)}(x) = (n + \alpha + 1)P_n^{(\alpha,1)}(x) + (n + \alpha)P_{n-1}^{(\alpha,1)}(x).$$

Transforming it to monic polynomials, multiplying by  $w^{\alpha,0}(x)$ , and integrating it over  $(-1, 1)$ , we obtain the following relation

$$(2n + \alpha + 1)f_n(-1) + \frac{2n(n + \alpha)}{2n + \alpha} f_{n-1}(-1) = -\frac{2n + \alpha + 1}{\alpha + 1} 2^{\alpha+1} \delta_{n,0}, \quad (4.7)$$

where  $\delta_{i,j}$  is Kronecker's delta. This gives

$$f_0(-1) = -\frac{2^{\alpha+1}}{\alpha + 1}, \quad f_n(-1) = -\frac{2n(n + \alpha)}{(2n + \alpha)(2n + \alpha + 1)} f_{n-1}(-1), \quad n \geq 1,$$

i.e.,

$$f_n(-1) = (-1)^{n+1} \frac{2^{n+\alpha+1} n!}{(n + \alpha + 1)_{n+1}}, \quad n = 0, 1, 2, \dots$$

Since  $\pi_n(-1) = \pi_n^{\alpha,1}(-1) = (-1)^n 2^n (n + 1)! / (n + \alpha + 2)_n$ , from (4.6) we find  $\lambda_0^{\alpha,0}$  in the form given in (4.3).

Finally, because of  $\sum_{k=0}^{n+1} \lambda_k^{\alpha,0} = 0$ , we obtain the second formula in (4.3).  $\square$

**REMARK 4.1.** The weights at the endpoints can be also expressed in the form

$$\lambda_0^{\alpha,0} = -\frac{2^\alpha}{\alpha + 1} - \frac{\alpha}{2} \sum_{k=1}^n \frac{w_k^{\alpha,1}}{1 + \xi_k^{\alpha,1}}, \quad \lambda_{n+1}^{\alpha,0} = \frac{2^\alpha}{\alpha + 1} - \frac{\alpha}{2} \sum_{k=1}^n \frac{w_k^{\alpha,1}}{1 - \xi_k^{\alpha,1}}.$$

Thus, there are simple explicit formulae for the weights  $\lambda_j^{\alpha,0}$ ,  $j = 0, 1, \dots, n+1$ . An alternative way for finding these coefficients are considered in [12].

#### 4.1. Software implementation

The construction of the Gauss-Jacobi-Lobatto nonstandard quadrature (4.1) for an arbitrary  $n \in \mathbb{N}$  and each  $\alpha > -1$  can be realized very easy by our MATHEMATICA package `OrthogonalPolynomials` (see [7] and [34]). Alternatively, for this purpose there is also Gautschi's package `SOPQ` written in MATLAB (cf. [31]). These packages provide many other calculations with orthogonal polynomials and different quadrature rules, and they are downloadable from Web Sites: <http://www.mi.sanu.ac.rs/~gvm/> and <http://www.cs.purdue.edu/archives/2002/wxg/codes/>, respectively.

The corresponding MATHEMATICA code for constructing parameters of the quadrature formula (4.1) is:

```
<< orthogonalPolynomials`
LobattoParam[n_, alpha_, digits_] := Module[{node, lambda, weight, k, a0, an1},
  {node, weight} = aGaussianNodesWeights[n, {aJacobi, alpha, 1},
    WorkingPrecision -> digits+5, Precision->digits];
  a0 = 2^alpha(alpha/((n+1)(n+alpha+1))-1);
  lambda = alpha*weight/(1-node^2);
  an1 = -(a0+Sum[lambda[[k]], {k, 1, n}]);
  lambda=N[Append[Prepend[lambda, a0], an1], digits];
  node=N[Append[Prepend[node, -1], 1], digits]; Return[{node, lambda}];
```

Variable-precision arithmetic enables us to calculate parameters (nodes and weights) in (4.1), named as `node` and `lambda`, i.e.,

$$\text{node} = (\xi_0^{\alpha,1}, \xi_1^{\alpha,1}, \dots, \xi_{n+1}^{\alpha,1}) \quad \text{and} \quad \text{lambda} = (\lambda_0^{\alpha,0}, \lambda_1^{\alpha,0}, \dots, \lambda_{n+1}^{\alpha,0}),$$

with an arbitrary precision. The input parameter `digits` defines the so-called *working precision* in the form `WorkingPrecision -> digits+5`.

For example, taking `digits-> 20` for  $\alpha = -1/2$  and  $n = 5$  we get the following results:

```
In[1] := {node, lambda} = LobattoParam[5, -1/2, 20]
```

```
Out[1] = {{-1.00000000000000000000, -0.78566926929466497066,
  -0.34243721374692749946, 0.19893554984718572955,
  0.68075005442268573279, 0.96270659305743529348,
  1.00000000000000000000},
  {-0.71782052029543460810, -0.072612263768525365535,
  -0.16642116952156041977, -0.37516617602834936907,
  -1.1131007878331247823, -10.292032937247316885,
  12.737153854694311430}}
```

The corresponding MATLAB code for constructing parameters of the quadrature formula (4.1) is:

```
function [node,lambda]=LobattoParam(n,q)
ab=r_jacobi(n,-q,1); xw=gauss(n,ab);
node=xw(:,1); omega=xw(:,2);
lambda(1)=-2^(-q)*(q/((n+1)*(n-q+1))+1);
lambda(2:n+1)=-q*omega(1:n)/(1-node(1:n).^2);
lambda(n+2)=-sum(lambda(1:n+1));
node=[-1 node' 1];
```

### 5. Applications to an approximation of fractional derivatives

Let  $f(s)$  be a sufficiently well-behaved function in  $[-1, 1]$ . If  $f(s)$  is defined in  $[0, t]$  instead of  $[-1, 1]$ , then change of variable  $s = \frac{t}{2}(1+x)$  could transform  $f(s)$  into  $g(x) := f(\frac{t}{2}(1+x))$ , where  $x \in [-1, 1]$ . Rewrite the definition (1.1) in the equivalent form

$$D_*^q f(t) = \frac{2^q}{t^q \Gamma(1-q)} \int_{-1}^1 g'(x)(1-x)^{-q} dx. \quad (5.1)$$

To approximate the integral in (5.1), we use the nonstandard quadrature (4.1) with  $\alpha := -q$ , i.e.,

$$(\forall g \in \mathbb{P}_{2n+1}) \quad \int_{-1}^1 g'(x)w^{-q,0}(x) dx = \sum_{k=0}^{n+1} \lambda_k^{-q,0} g(\xi_k^{-q,1}), \quad (5.2)$$

where  $\xi_k^{-q,1}$  are the zeros of the polynomial  $(1-x^2)P_n^{(-q,1)}(x)$ , and the weights  $\lambda_j^{-q,0}$  are given by (4.2) and (4.3), in which  $\alpha := -q$ . Thanks to the representation (5.1), the following approximation of the Caputo fractional derivative (1.1)

$$D_*^q f(t) \approx D_{*,n}^q f(t) := \frac{2^q}{t^q \Gamma(1-q)} \sum_{k=0}^{n+1} \lambda_k^{-q,0} f(t_k^{-q,1}) \quad (5.3)$$

holds, where

$$t_k^{-q,1} = \frac{t}{2}(\xi_k^{-q,1} + 1).$$

The relationship (1.3) leads to

$${}_0 D_t^q f(t) \approx {}_0 D_{t,n}^q f(t) := \frac{f(0)}{t^q \Gamma(1-q)} + D_{*,n}^q f(t). \quad (5.4)$$



from which we see that the algebraic degree of exactness is  $d = 11$  that is predicted by Theorem 4.1.

Similarly, we can conclude from the second case for  $f(t) = t^\gamma$ , for which

$${}_0D_t^q t^\gamma = \frac{\Gamma(1 + \gamma)}{\Gamma(1 + \gamma - q)} t^{\gamma - q}, \quad \gamma > -1.$$

The corresponding sequence  $\tilde{E}_n^q$  for  $\gamma = 0, 1, \dots, 12$  (the maximum of the errors on  $(0, 1]$ ), defined by (6.3), with the same quadrature ( $n = 5$  and  $q = 1/2$ ), is

$$\{0. \times 10^{-18}, 0. \times 10^{-19}, 0. \times 10^{-19}, 0. \times 10^{-19}, 0. \times 10^{-19}, 0. \times 10^{-19}, 0. \times 10^{-19}, \\ 0. \times 10^{-19}, 0. \times 10^{-19}, 0. \times 10^{-19}, 0. \times 10^{-18}, 0. \times 10^{-18}, 2.55 \times 10^{-7}\}.$$

Using the same quadrature ( $q = 1/2$  and  $n = 5$ ), the maximum of the errors  $\tilde{E}_n^{1/2}$  for  $f(t) = t^\gamma$ , when  $\gamma$  runs over  $[1, 12]$ , is presented in Figure 1 (left) in  $\log_{10}$ -scale. As we can see the quadrature formula is exact in the cases when  $\gamma$  is an positive integer less than 12. In the same figure we give also the corresponding graphics for cases when  $n = 10, n = 15$  and  $n = 20$ . Notice that these quadratures are exact for  $\gamma \in \{0, 1, \dots, 2n + 1\}$ . The case  $q = 9/10$  is corresponding graphics are presented in Figure 1 (right).




FIGURE 1. Maximum of  $\log_{10} \tilde{E}_n^q$  for  $f(t) = t^\gamma$ ,  $q = 1/2$  (left) and  $q = 9/10$  (right), for quadratures with  $n = 5(5)20$  nodes (graphics from top to bottom), when  $\gamma$  runs over interval  $[1, 12]$

Now we consider the error  $e_n^{1/2}(f; t)$ , defined by (6.2), for a simple power function  $f(t) = t^\gamma$  with an algebraic singularity. We take  $\gamma = 1/2, 3/2, 7/2$ , and  $11/2$ , and apply the Gauss-Lobatto quadratures (5.3) with  $n = 5, 10, 15$ , and 20 nodes (plus two fixed nodes at  $\pm 1$ ). The corresponding graphics in log-scale are presented in Figure 2.

As we can see the error  $e_n^{1/2}(f; t)$  for a small  $\gamma$  ( $= 1/2$ ) is almost constant for each  $t$  in the interval  $(0, 1)$ , but the convergence of the quadrature

$\gamma$	$n = 5$	$n = 10$	$n = 15$	$n = 20$	$n = 30$	$n = 60$	$n = 90$	$n = 120$
$\frac{1}{2}$	5.88(-4)	9.03(-5)	2.87(-5)	1.26(-5)	3.86(-6)	5.01(-7)	1.50(-7)	6.38(-8)
$\frac{1}{4}$	2.83(-3)	5.92(-4)	2.28(-4)	1.14(-4)	4.28(-5)	7.80(-6)	2.86(-6)	1.40(-6)
$\frac{1}{8}$	5.89(-3)	1.44(-3)	6.10(-4)	3.30(-4)	1.35(-4)	2.93(-5)	1.19(-5)	6.24(-6)
$\frac{1}{16}$	8.45(-3)	2.23(-3)	9.92(-4)	5.52(-4)	2.39(-4)	5.63(-5)	2.40(-5)	1.31(-5)

TABLE 1. Errors  $e_n^{1/2}(f; 1/2)$  in quadrature sums with  $n = 5(5)20$  and  $n = 30(30)120$ , for  $\gamma = 1/2, 1/4, 1/8,$  and  $1/16$

formulas (5.3) is rather slow, especially when  $\gamma$  is closer to zero. For example, for  $\gamma = 1/2, 1/4, 1/8,$  and  $1/16$ , the errors  $e_n^{1/2}(f; t)$  at  $t = 1/2$  are given in Table 1 for quadrature rules with  $n = 5(5)20$  and  $n = 30(30)120$  nodes. Numbers in parentheses indicate decimal exponents. For  $\gamma > 1$  the convergence is satisfactory.




FIGURE 2. Graphics of  $\log_{10} e_n^q(f; t)$  for  $f(t) = t^\gamma$  and  $t$  runs over  $(0, 1)$ , when  $n = 5(5)20$  (graphics from top to bottom) in the cases: (above)  $\gamma = 1/2$  (left) and  $\gamma = 3/2$  (right); (below)  $\gamma = 7/2$  (left) and  $\gamma = 11/2$  (right)

**REMARK 6.1.** Problems with algebraic or/and logarithmic singularities can be solved using Müntz systems and quadratures of this type (cf. [30, 32, 11]) or using a procedure proposed in [20].

**EXAMPLE 6.2.** For our second experiment we consider fractional differentiation by plotting  $\sin(t)$  on the interval  $[0, 4\pi]$  (already used in [19]), together with its derivatives of order  $q = 1/10, 2/10, \dots, 9/10$ , as drawn in Figure 3. Fractional derivatives have been calculated using the quadrature rule (5.3) with  $n = 5$ . As one can see, each curve is approximately a translation of  $\sin(t)$  by a distance  $q\pi/2$  to the left.




FIGURE 3. The function  $\sin(t)$  on  $[0, 4\pi]$  together with its derivatives of order  $k/10$ ,  $k = 1, 2, \dots, 9$

**EXAMPLE 6.3.** We compute  ${}_0D_t^q f(t)$  for  $f(t) = \sin(\lambda t)$  and  $t \in [0, \pi]$ . The exact value of the fractional derivative of this function is given by [39, 41]

$${}_0D_t^q \sin(\lambda t) = \lambda t^{1-q} E_{2,2-q}(-\lambda^2 t^2),$$

where the Mittag-Leffler function is defined in (6.1).

In Figure 4 we present graphics of the functions  $\sin(\lambda x)$ ,  $\lambda = 1, 2, 3$  (left), and their derivatives of order  $q = 1/2$  (right), obtained by the Gauss-Lobatto quadrature rule (5.3) with  $n = 5$  internal nodes. Since the sine function is smooth, this quadrature rule converges very fast. In order to illustrate this fact we present in Table 2 the numerical results  ${}_0D_{t,n}^q \sin(\lambda t)$  for  $t = \pi/2$ ,  $q = 1/2$ ,  $\lambda = 2$  and  $\lambda = 3$ , taking  $n = 2(1)8$  internal nodes in the Gauss-Lobatto quadrature rule (5.3), as well as the corresponding






FIGURE 4. Graphics of the functions  $\sin(\lambda x)$ ,  $\lambda = 1, 2, 3$  (left), and their derivatives obtained by the numerical quadrature (5.3) with  $n = 5$  (right)

$n$	${}_0D_{t,n}^q \sin(2t)$	$r_n^q(2; t)$	${}_0D_{t,n}^q \sin(3t)$	$r_n^q(3; t)$
2	-1.05 <b><u>6</u></b> 8638589376709	8.69(-4)	-1.26 <b><u>4</u></b> 0813951622687	2.41(-3)
3	-1.0577 <b><u>9</u></b> 33376552489	9.59(-6)	-1.267 <b><u>2</u></b> 323502405542	7.79(-5)
4	-1.0577831 <b><u>2</u></b> 05699668	6.58(-8)	-1.26713 <b><u>1</u></b> 8332287842	1.39(-6)
5	-1.057783190 <b><u>5</u></b> 482818	3.08(-10)	-1.267133 <b><u>6</u></b> 100910347	1.59(-8)
6	-1.05778319022 <b><u>1</u></b> 3884	1.04(-12)	-1.267133589 <b><u>7</u></b> 303999	1.29(-10)
7	-1.0577831902224 <b><u>9</u></b> 60	2.69(-15)	-1.26713358989 <b><u>5</u></b> 1450	7.81(-13)
8	-1.057783190222493 <b><u>2</u></b>	5.41(-18)	-1.26713358989415 <b><u>0</u></b> 1	3.67(-15)

TABLE 2. Gaussian approximations  ${}_0D_{t,n}^q \sin(\lambda t)$  and the corresponding relative errors  $r_n^q(\lambda; t)$  for  $f(t) = \sin(\lambda t)$  and  $q = 1/2$  at  $t = \pi/2$

relative errors

$$r_n^q(\lambda; t) = \left| \frac{{}_0D_{t,n}^q \sin(\lambda t) - {}_0D_t^q \sin(\lambda t)}{{}_0D_t^q \sin(\lambda t)} \right|.$$

In each entry the first digit in error is bolded and underlined.

The maximal absolute errors in Gaussian approximations over the whole interval  $(0, \pi)$  (taking 1000 equidistant points  $t_j = j\pi/1000$ ,  $j = 1, \dots, 1000$ ) are presented in Table 3.

EXAMPLE 6.4. We compute  ${}_0D_t^q f(t)$  for  $f(t) = (t+\nu)^{q-1}$  and  $t \in [0, 1]$ . The exact value of the fractional derivative of this function is given by [39]

$${}_0D_t^q (t + \nu)^{q-1} = \frac{1}{\Gamma(1-q)} \left(\frac{\nu}{t}\right)^q \frac{1}{t + \nu}.$$

$n$	4	6	8	10	12
$\lambda = 1$	4.93(-8)	7.81(-13)	4.05(-18)		
$\lambda = 2$	1.73(-5)	3.42(-9)	2.32(-13)	6.80(-18)	
$\lambda = 3$	1.50(-3)	2.41(-6)	1.13(-9)	2.12(-13)	1.91(-17)

TABLE 3. Maximal absolute errors in Gaussian approximations  ${}_0D_{t,n}^q \sin(\lambda t)$ , when  $t \in (0, \pi]$ ,  $q = 1/2$ ,  $\lambda = 1, 2, 3$  and  $n = 4(2)12$

The actual maximum error  $\tilde{E}_n^q$ , defined by (6.3), is plotted in Figure 5 in a semi-log coordinate system as a function of  $n$  for  $q = 1/2$  and four values of the parameter  $\nu$  ( $= 0.1, 0.2, 0.5$  and  $1.0$ ).




FIGURE 5.  $\log_{10} \tilde{E}_n^q$  versus  $n$  for  $q = 1/2$  with  $\nu = 0.1, 0.2, 0.5, 1.0$

As we can see that  $\tilde{E}_n^q$  decreases rapidly. It is worth noting that the convergence rate is dependent of the regularity of the integrand in the neighborhood of  $t = 0$ . As one can see, in the case that  $\nu$  tends to zero, the quadrature rule converges slowly.

**EXAMPLE 6.5.** For exponential functions  $f_1(t; \lambda) = e^{\lambda t}$  and  $f_2(t; \lambda) = \cosh(\sqrt{\lambda} t)$ ,  $t \in [0, 1]$ , the exact values of the fractional derivative  ${}_0D_t^q f_k(t; \lambda)$ ,  $k = 1, 2$ , are given in terms of the Mittag-Leffler function (6.1) (cf. [36, 41]) as

$${}_0D_t^q e^{\lambda t} = t^{-q} E_{1,1-q}(\lambda t)$$

	$n$	4	6	8	10
$f_1$	$\lambda = 1/2$	1.28(-10)	1.20(-16)		
	$\lambda = 1$	3.32(-7)	4.81(-12)	2.36(-17)	
	$\lambda = 2$	2.36(-3)	4.49(-7)	3.71(-11)	1.20(-15)
$f_2$	$\lambda = 1/2$	3.25(-9)	1.21(-14)	1.51(-20)	
	$\lambda = 1$	1.71(-7)	2.48(-12)	1.22(-17)	
	$\lambda = 2$	1.18(-5)	6.59(-10)	1.27(-14)	1.05(-19)

TABLE 4. Maximal absolute errors  $\tilde{E}_n^q(f_k)$ ,  $k = 1, 2$ ,  $\lambda = 1/2, 1, 2$ , over  $(0, 1]$ , in Gaussian approximations for  $q = 1/2$ , with  $n = 4(2)10$  internal nodes

and

$${}_0D_t^q \cosh(\sqrt{\lambda} t) = t^{-q} E_{2,1-q}(\lambda t^2).$$

The maximal absolute errors in Gaussian approximations  $\tilde{E}_n^q(f_k)$ ,  $k = 1, 2$ , over the whole interval  $(0, 1)$ , defined by (6.3), are presented in Table 4.

EXAMPLE 6.6. Finally, we compute  ${}_0D_t^q f(t)$  for  $f(t) = t^{a/2} J_a(2\sqrt{t})$  and  $t \in [0, 1]$ , where  $J_a(z)$  is the Bessel function of the first kind and order  $a$ , defined as a solution of the so-called Bessel differential equation

$$z^2 \frac{d^2 y}{dz^2} + z \frac{dy}{dz} + (z^2 - a^2)y = 0,$$

which can be expressed in the power form expansion around  $z = 0$ ,

$$J_a(z) = \sum_{k=0}^{\infty} \frac{(-1)^k}{k! \Gamma(k + a + 1)} \left(\frac{z}{2}\right)^{2k+a}.$$

The exact value of the fractional derivative of this function is given by [36, 41]

$${}_0D_t^q f(t) = t^{(a-q)/2} J_{a-q}(2\sqrt{t}),$$

Since the original function

$$f(t) = f(t; a) = t^{a/2} J_a(2\sqrt{t}) = \sum_{k=0}^{\infty} \frac{(-1)^k}{k! \Gamma(k + a + 1)} t^{k+a},$$

we see that  $f$  is an analytic function when  $a \in \mathbb{N}_0$ , so that for such values of the parameter  $a$  we expect fast convergence of the quadrature rule (5.3). However, in other cases this function has an algebraic singularity at  $t = 0$ .

As  $a$  decreases to zero, the convergence of our quadrature rule slows down considerably. In Figure 6 we present graphics of the maximal error  $\tilde{E}_n^q(f(\cdot; a))$  for  $q = 1/2$ , obtained by three different quadrature rules with

$n = 5, 10,$  and  $20$  internal nodes, when the parameter  $a$  in the function runs over  $(0, 3.5)$ . It is clear that the rapidly increasing of accuracy achieved when the parameter  $a$  tends to an integer, i.e., when  $f$  becomes an analytic function.




FIGURE 6. Graphics of  $\log_{10}[\tilde{E}_n^{1/2}(f(\cdot; a))]$  for  $n = 5, 10,$  and  $20$  (from top to bottom), when  $0 < a < 3.5$




FIGURE 7. The values  $\log_{10}[e_n^{1/2}(f; t)], 0 < t < 1,$  for  $n = 2, 3, 4$  internal nodes (graphics from top to bottom), when  $a = 2$  (left) and  $a = 3$  (right)

Thus, for  $a \in \mathbb{N}_0$  the convergence of the quadrature rule (5.3) is very fast! The errors  $e_n^{1/2}(f; t), 0 < t < 1,$  in approximation of fractional derivative  ${}_0D_t^q f(t)$  by quadrature rules with a small number of nodes ( $n = 2, 3, 4$ ) are presented in Figure 7 (in log-scale) for  $a = 2$  (left) and  $a = 3$  (right).

On the other side, taking much more nodes (up to  $n = 120$ ), for a small  $a$ , viz.  $a = 1/2$ , the convergence is very slow (see Figure 8 (left)). Something faster convergence is achieved for  $a = 3/2$  (Figure 8 (right)).




FIGURE 8. The values  $\log_{10}[e_n^{1/2}(f;t)]$ ,  $0 < t < 1$ , for  $n = 10, 20, 30, 60, 120$  internal nodes (graphics from top to bottom), when  $a = 1/2$  (left) and  $a = 3/2$  (right)

However, if  $a$  is sufficiently large the convergence rate becomes quite satisfactory, as is given in Figure 9 (left) for  $a = 7/2$ , where we used  $n = 5(5)20$  and  $n = 30$  internal nodes in the corresponding quadrature rule (5.3).




FIGURE 9. (left) The values  $\log_{10}[e_n^{1/2}(f;t)]$ ,  $0 < t < 1$ ,  $a = 7/2$ , for  $n = 5, 10, 15, 20, 30$  internal nodes (graphics from top to bottom); (right)  $\log_{10}[\tilde{E}_n^q(f(\cdot; a))]$ ,  $a = 3/2$ , versus  $n$  for  $q = 1/10, 1/2$ , and  $9/10$

Finally, in Figure 9 (right) we present the actual maximum error  $\tilde{E}_n^q(f)$ , defined by (6.3), in a semi-log coordinate system as a function of  $n$  for  $a = 3/2$  and three values of  $q$  ( $= 1/10, 1/2$ , and  $9/10$ ). As we can see the behaviour of  $\tilde{E}_n^q(f)$  is very similar with respect to  $q$ .

### Acknowledgements

The second author was supported in part by the Serbian Ministry of Education, Science and Technological Development (No. #OI 174015), as well as under a bilateral agreement between Serbian Academy of Sciences and Arts and Bulgarian Academy of Sciences (“Mathematical modelling via...”, 2012–2014).

### References

- [1] G.E. Andrews, R. Askey, R. Roy, *Special Functions*. Encyclopedia of Mathematics and its Applications, 71, Cambridge University Press, Cambridge (1999).
- [2] T.M. Atanacković, S. Pilipović, B. Stanković, D. Zorica, *Fractional Calculus with Applications in Mechanics: Vibrations and Diffusion Processes*. ISTE, London – Wiley, Hoboken (2014).
- [3] D. Baleanu, K. Diethelm, E. Scalas, J. J. Trujillo, *Fractional Calculus: Models and Numerical Methods*. World Scientific, Singapore (2012).
- [4] H. Brass, K. Petras, *Quadrature Theory: The Theory of Numerical Integration on a Compact Interval*. American Mathematical Soc., Providence, RI. (2011).
- [5] P.L. Butzer, U. Westphal, An introduction to fractional calculus, In: *Applications of Fractional Calculus in Physics*. World Sci. Publ., River Edge, NJ (2000), 1–85.
- [6] M. Caputo, Linear models of dissipation whose  $Q$  is almost frequency independent – II. *Geophysical Journal of the Royal Astronomical Society* **13** (1967), 529–539.
- [7] A.S. Cvetković, G.V. Milovanović, The Mathematica package ”OrthogonalPolynomials”. *Facta Univ. Ser. Math. Inform.* **9** (2014), 17–36.
- [8] K. Diethelm, *The Analysis of Fractional Differential Equations*. Springer-Verlag, Berlin, 2010.
- [9] K. Diethelm, Error bounds for the numerical integration of functions with limited smoothness. *SIAM J. Numer. Anal.* **52**, No 2 (2014), 877–879.
- [10] S. Esmaeili, M. Shamsi, M. Dehghan, Numerical solution of fractional differential equations via a Volterra integral equation approach. *Centr. Eur. J. Phys.* **11** (2013), 1470–1481.
- [11] S. Esmaeili, M. Shamsi, Y. Luchko, Numerical solution of fractional differential equations with a collocation method based on Müntz polynomials. *Comput. Math. Appl.* **62** (2011), 918–929.
- [12] D. Funaro, *Polynomial Approximation of Differential Equations*. Springer-Verlag, Berlin (1992).

- [13] R. Garrappa, M. Popolizio, Evaluation of generalized Mittag-Leffler functions on the real line. *Adv. Comput. Math.* **39** (2013), 205–225.
- [14] W. Gautschi, *Orthogonal Polynomials: Computation and Approximation*. Oxford University Press, New York (2004).
- [15] G.H. Golub, J.H. Welsch, Calculation of Gauss quadrature rules. *Math. Comp.* **23** (1969), 221–230.
- [16] R. Gorenflo, J. Loutchko, Y. Luchko, Computation of the Mittag-Leffler function and its derivatives, *Fract. Calc. Appl. Anal.* **5** (2002), 491–518.
- [17] R. Gorenflo, F. Mainardi, Fractional calculus: integral and differential equations of fractional order. In: *Fractals and Fractional Calculus in Continuum Mechanics (Udine, 1996)*, CISM Courses and Lectures, 378, Springer, Vienna (1997), 223–276.
- [18] N. Hale, A. Townsend, Fast and accurate computation of Gauss–Legendre and Gauss–Jacobi quadrature nodes and weights. *SIAM J. Sci. Comput.* **35** (2013), 652–674.
- [19] N. Hale, L.N. Trefethen, Chebfun and numerical quadrature. *Sci. China Ser. A* **55** (2012), 1749–1760.
- [20] T. Hasegawa, H. Sugiura, Uniform approximation to fractional derivatives of functions of algebraic singularity. *J. Comput. Appl. Math.* **228** (2009), 247–253.
- [21] R. Herrmann, *Fractional Calculus: An Introduction for Physicists*. World Scientific, Singapore (2011).
- [22] A.A. Kilbas, H.M. Srivastava, J.J. Trujillo, *Theory and Applications of Fractional Differential Equations*. North-Holland Mathematics Studies, 204, Elsevier Science B.V., Amsterdam (2006).
- [23] V. Kiryakova, *Generalized Fractional Calculus and Applications*. Pitman Research Notes in Math. Series, 301, Longman Scientific & Technical, Harlow; copubl. by John Wiley & Sons, Inc., New York (1994).
- [24] C. Li, F. Zheng, F. Liu, Spectral approximations to the fractional integral and derivative. *Fract. Calc. Appl. Anal.* **15** (2012), 383–406; DOI: 10.2478/s13540-012-0028-x; <http://link.springer.com/article/10.2478/s13540-012-0028-x>.
- [25] F. Mainardi, *Fractional Calculus and Waves in Linear Viscoelasticity. An Introduction to Mathematical Models*. Imperial College Press, London (2010).
- [26] F. Mainardi, Fractional calculus: some basic problems in continuum and statistical mechanics. In: *Fractals and Fractional Calculus in Continuum Mechanics (Udine, 1996)*, CISM Courses and Lectures, 378, Springer, Vienna (1997), 291–348.

- [27] F. Mainardi, Y. Luchko, G. Pagnini, The fundamental solution of the space-time fractional diffusion equation. *Fract. Calc. Appl. Anal.* **4**, No 2 (2001), 153–192.
- [28] G. Mastroianni, G.V. Milovanović, *Interpolation Processes: Basic Theory and Applications*. Springer–Verlag, Berlin (2008).
- [29] K.S. Miller, B. Ross, *An Introduction to the Fractional Calculus and Fractional Differential Equations*. A Wiley-Interscience Publication, John Wiley & Sons, Inc., New York (1993).
- [30] G.V. Milovanović, Müntz orthogonal polynomials and their numerical evaluation. In: *Applications and Computation of Orthogonal Polynomials* (W. Gautschi, G.H. Golub, and G. Opfer, Eds.), ISNM, Vol. 131, Birkhäuser, Basel (1999), 179–202.
- [31] G.V. Milovanović, Chapter 23: Computer algorithms and software packages. In: *Walter Gautschi: Selected Works and Commentaries*, Volume **3** (C. Brezinski, A. Sameh, Eds.), Birkhäuser, Basel (2014), 9–10.
- [32] G.V. Milovanović, A.S. Cvetković, Gaussian type quadrature rules for Müntz systems. *SIAM J. Sci. Comput.* **27** (2005), 893–913.
- [33] G.V. Milovanović, A.S. Cvetković, Nonstandard Gaussian quadrature formulae based on operator values. *Adv. Comput. Math.* **32** (2010), 431–486.
- [34] G.V. Milovanović, A.S. Cvetković, Special classes of orthogonal polynomials and corresponding quadratures of Gaussian type. *Math. Balkanica* **26**, No 1-2 (2012), 169–184.
- [35] P. Novati, Numerical approximation to the fractional derivative operator. *Numer. Math.* **127** (2014), 539–566.
- [36] I. Podlubny, *Fractional Differential Equations*. Academic Press, San Diego, CA (1999).
- [37] I. Podlubny, M. Kacenak, *The MATLAB mlf code*. MATLAB Central File Exchange (2001–2012), File ID: 8738.
- [38] S.G. Samko, A.A. Kilbas, O.I. Marichev, *Fractional Integrals and Derivatives. Theory and Applications*. Edited and with a foreword by S.M. Nikol’skiĭ, Transl. from the 1987 Russian original, Revised by the authors. Gordon and Breach Science Publishers, Yverdon (1993).
- [39] H. Sugiura, T. Hasegawa, Quadrature rule for Abel’s equations: uniformly approximating fractional derivatives. *J. Comput. Appl. Math.* **223** (2009), 459–468.
- [40] L.N. Trefethen, *Approximation Theory and Approximation Practice*, SIAM, Philadelphia, PA (2013).



- [41] D. Valério, J.J. Trujillo, M. Rivero, J.A.T. Machado, D. Baleanu, Fractional calculus: A survey of useful formulas. *Eur. Phys. J. Special Topics* **222** (2013), 1827–1846.

<sup>1</sup> *Department of Applied Mathematics  
University of Kurdistan  
P.O. Box 416, Sanandaj, IRAN  
e-mail: sh.esmaeili@uok.ac.ir*

*Received: April 16, 2014*

<sup>2</sup> *Mathematical Institute  
Serbian Academy of Sciences and Arts  
Kneza Mihaila 36, 11001 Belgrade, SERBIA  
e-mail: gvm@mi.sanu.ac.rs*

---

Please cite to this paper as published in:

*Fract. Calc. Appl. Anal.*, Vol. **17**, No 4 (2014), pp. 1075–1099;  
DOI:10.2478/s13540-014-0215-z