Time to tell the world about osteopathic medicine

To the Editor:

I would like to address some of the concerns expressed in the letter "Key to our survival lies in honesty," written by Christopher D. Olson, DO, in the September 1998 issue of the *JAOA* (1998;98:475-476). Like Dr Olson, I, too, am concerned about the future of this profession, and many of the decisions the American Osteopathic Association (AOA) has made, and will make, will have a great impact on the future of osteopathic medicine.

I'd like to commend Dr Olson for sending a very thoughtful letter. This is just the type of discourse we encourage from within the profession. As Dr Olson stated, osteopathic medicine is a strong profession with a valuable history. And today, many of the aspects of DO training are indeed similar to allopathic medicine. But, while he argues that aside from manipulation there is no real difference between the two, I believe that the science of osteopathic medicine, as taught in our schools, clearly differentiates between DOs and MDs. And it is more than just manipulation. It's a form of medicine founded on the body's relationship with the musculoskeletal system. It's a form of medicine that has always included the psychosocial component of seeking and maintaining health, not just treating disease. Without clearly defining a distinction between osteopathic and allopathic medicine, the very future of our profession is at stake. The AOA and other affiliated organizations have fought hard to demonstrate our base level of training and our ability to deliver unequaled medical care. And we are making great strides in leveling the playing field and gaining equal regard with the allopathic profession. But just as this country has no need for maintaining separate degrees for other professions

(lawyers, teachers, and accountants) unless we believe, teach, and offer something tangible and unique, we certainly risk further assimilation and the eventual end to this profession.

Osteopathic medicine, including manipulation, and the art of osteopathic principles and practice (OPP), absolutely needs to continue to be developed and taught in our schools. That is why, in part, the AOA established the Committee on Osteopathic History. This committee must explore more effective ways of teaching our rich tradition and history to future students. You never get a second chance to make a first impression, and how our student doctors are exposed to OPP and osteopathic manipulative treatment (OMT) is vital to how they embrace it and eventually practice it years later.

And to that end, we must continue to seek out students who not only have an interest in primary care medicine, but who find the osteopathic medical philosophy appealing and consistent with their own beliefs of how medicine should be practiced. We must continue to more effectively train osteopathic physicians. Our current AOA president, Dr Esper, and I are making an effort to reach out to students. We must create and maintain open lines of communication with them, listen to their concerns, and urge caution about accepting all the negative comments they read and hear online from other students. The last thing we would ever want to do is stifle students' discourse, when their opinions are not in agreement with those of our leaders. Thinking back to my college days, I know that students expressing dissatisfaction with their school situation is quite common.

Dr Olson also addressed continuing medical education (CME). The AOA's

Council on Continuing Medical Education is working on ways to increase the types of programs that can be used to certify osteopathic physicians, including allopathic programs. Conversely, we want our osteopathic CME training to be recognized by the Accreditation Council of Continuing Medical Education (ACCME). This recognition will allow an even greater mix of training that DOs can receive, as well as open the door for AOA certification and membership for those DOs who are unable to get to AOA-approved programs. I don't expect that DOs will in turn abandon their osteopathic CME for other types of training. No matter what the specialty, the underlying osteopathic medical philosophy should remain an important part of CME.

The very foundation of osteopathic medicine's valuable history, lay in the search for an approach that differed from the conventional medicine of the day. Osteopathic medicine began as, and continues to be, both a healthcare and social reform movement. And while we are 100-plus years beyond its origin, we need to continue to maintain and highlight our distinctiveness, so we can effectively argue for having two types of physicians in this country. For much of our history, we have fought, with good success, the battle toward equal recognition with allopathic medicine. Now, with the Campaign for Osteopathic Unity, it is time that we aggressively tell the public what we have known for more than a century, namely what is right, and different, and worth exploring about osteopathic medicine. I came across an article that appeared in April in *USA Today*, written by two MD students. They talked about how doctors are not trained to be caring and compassionate to touch their patients to show them that they care. I wanted to

say yes, we are! That is part of what osteopathic medicine has always been about! Our message needs to be heard, and the time is now!

Eugene A. Oliveri, DO

1998-99 AOA President-elect
Chair, AOA Campaign for Osteopathic
Unity
Milford, Mich

Gerald C. Osborn, DO, MPhil, FACN

Acting Associate Dean
Michigan State University—College of
Osteopathic Medicine
East Lansing, Mich

Osteopathic profession needs leaders with progressive stance

To the Editor:

The pleasure and the encouragement I felt in reading the excellent letter from Christopher Olsen, DO, "Key to our survival lies in honesty," in the September 1998 issue of the *JAOA* (1998;98:475-476) moved me to write this follow-up letter to tender my support. His was a kinder, gentler expression of sentiments that needed to be conveyed. It is my belief that osteopathic physicians who read the *JAOA* also overwhelmingly support Dr Olson's opinions. Yet, I hold some level of trepidation that his letter will result in an onslaught of negative responses similar to that which followed mine (see my letter, "Time to abolish most osteopathic graduate medical education programs," published in the March 1995 *JAOA* (1995;95:155)).

I suspect Dr Olson also harbors some trepidation in that he felt it necessary to preface his opinions by enumerating his many credentials as a model osteopathic physician. That his doing so was probably necessary is truly a shame. I would encourage him to take heart in the belief that the silent majority is with him. The sooner our party line takes on these same notes of reason, the sooner we can move away from the stifling attitudes that hold out osteopathy as some sort of immutable religion. Our profession should be led

by those with Dr Olson's progressive stance, rather than by the zealots who thump on the writings of A.T. Still as if those writings were a bible, shouting down opposing opinions as if they came from heretics. I vigorously applaud the forthright honest of Dr Olson.

Ronald Kienitz, DO

Concentra Medical Centers
Kailua, Hawaii

Revisiting the role of osteopathic manipulation in primary care

To the Editor:

As the lead author of the article, "Evaluation and treatment of sinusitis: Aspects for the managed care environment," in the April 1996 Supplement to the *JAOA* (1996;96(4):S6-S10), I would like to comment on the letter by Dr Dudley, "Sinusitis Supplement missing osteopathic component" in the October issue (*JAOA* 1998;98:539-540). Dr Dudley notes that she is disappointed that osteopathic manipulative treatments for sinusitis are not emphasized.

In talking with and lecturing to primary care physicians, I've found that the concept of sinusitis appears to be confusing and often misunderstood by both physicians and patients. In my published article, my colleague and I attempted to outline the clinical picture of the many "faces" of sinusitis and give rational, reasonable approaches for medical management.

Many cases of sinusitis are viral in origin and self limited (*J Allergy Clin Immunol* 1998;102:403-408). I have no doubts, therefore, that if patients with sinusitis are treated with osteopathic manipulative techniques most will improve, and the clinician will presume the success was the result of this therapy.

Russell J. Hopp, DO

Professor of Pediatrics
Creighton University
School of Medicine
Omaha, Neb

To the Editor:

It is just amazing that the last several *Supplements* have all stimulated great response from osteopathic physicians who are fed up with the exclusion and omission of osteopathic principles and practice as they apply to integrating osteopathic manipulative medicine to common problems in primary care.

I, too, agree with Dr Dudley. I have also been reading the *JAOA* and *The DO* since I was a student at the Kirksville College of Osteopathic Medicine in the 1980s. It is truly a shame that as we approach the new millenium, our organization, which so desperately needs to survive and needs to remain separate from allopathic medicine by its uniqueness, still publishes pieces without the integration of osteopathic principles and practice.

I don't think anyone can argue that even though the Editorial Advisory Board includes several physicians with expertise in manipulation, the information in the *Supplements* and other publications remains largely nonosteopathic in nature. Our students and residents, who are the future of our profession, see these articles and turn to me and others who teach and just shake their heads. Is it no wonder that these publications end up discarded because we can get similar information from allopathic medical journals with a whole seemingly higher value, such as *The New England Journal of Medicine* and *JAMA*.

This is not to say the information contained in the *JAOA* is not stimulating or important. Some excellent articles, with osteopathic components at their core, have been published, and I have used them over the years in teaching. I do think, however, room exists for improvement. I hope our new editor in chief will take these comments to heart and seek to accept no articles unless they have some mention of osteopathic manipulative skills.

I challenge the AOA to call on those of us who have used our clinical skills to help integrate our osteopathic manipulative training into some of these articles on pulmonary, allergic, and other problems. If this profession wants to contin-

ue to thrive, it must exercise its uniqueness and show the public, as well as its own, the uniqueness of our hands-on holistic approach that accentuates the preventative nature of the osteopathic medical profession. The respect we would garner from our allopathic and chiropractic colleagues, as well as the lay public, would be incredible. Currently, people look at us as a cross between chiropractors and MDs, depending on how we practice. It is still ludicrous that we are one of the least understood professions in this country—if not the world—and yet we still concentrate on trying to be mainstream medicine.

David S. Abend, DO
Emerson, NJ

Colposcopy may not be clinically relevant

To the Editor:

I found the article “Natural history of false-negative Papanicolaou smears: A prospective study using screening colposcopy in addition to cytology” by Dr Mannino (*JAOA* 1998;98(10):542-546) to be of academic interest. However, I disagree with the author’s conclusions that the primary screening for cervical cancer with colposcopy is a worthwhile endeavor. The article suggests that women at high risk perhaps would benefit from screening colposcopy; however, the study fails to identify what subgroups of women are to be considered high risk.

Colposcopy is an uncomfortable procedure for the patient, more time consuming for both the patient and physician, and a costly procedure as well, ranging anywhere from \$150 to \$350 per procedure. Furthermore, women who are not compliant with cytologic screening are not anymore likely to be compliant with colposcopic screening.

Several studies¹⁻³ have been done looking at screening colposcopy, and the overall conclusions are that it is an overly sensitive test for primary screening. Dr Mannino’s article states that an average of 25 months (plus or minus 7 months)

is the time it takes for the false-negative Papanicolaou (Pap) smear to mirror colposcopic biopsy results. This may be so; however, the majority of patients in the study remained with low-grade disease, and none of the patients progressed to cervical cancer or carcinoma in situ, thus, making the argument irrelevant that earlier diagnosis of cervical dysplasia by colposcopic biopsy can have an impact on outcome. Earlier treatment would not have made a difference.

In summary, it may be neither convenient nor cost effective to use colposcopy as a primary screening method for cervical cancer, as it seems as if it may be an overly sensitive test in this setting. In the context of a normal Pap smear, colposcopic biopsy results are not likely to be clinically relevant.

Richard R. Terry, DO, CAQ
Adolescent Medicine
Program Director
Wilson Family Practice Residency
Johnson City, New York

References

1. Cecchini S, Bonardi R, Iossa A, Zappa M, Ciatto S. Colposcopy as a primary screening test for cervical cancer. *Tumori* 1997;83(5):810-813.
2. Lonky NM, Mann WJ, Massad LS, Mutch DG, Blanco JS, Vasilev SA, et al. Ability of visual tests to predict underlying cervical neoplasia. Colposcopy and speculscopy. *J Reprod Med* 1995;40(7):530-536.
3. van Niekerk WA, Dunton CJ, Rihart RM, Hilgarth M, Kato H, Kaufman RH, et al. Colposcopy, cervicography, speculscopy and endoscopy. International Academy of Cytology Task Force Summary. Diagnostic Cytology Towards the 21st Century: An International Expert Conference and Tutorial. *Acta Cytologica* 1998;42(1):33-49.

Response

To the Editor:

No controversy exists in saying that the Papanicolaou (Pap) smear is a cost-effective screen for cervical carcinoma that has saved lives. However, the death rate from cervical carcinoma has remained relatively static over the past decade.¹ The impetus for my study, “Natural his-

tory of false-negative Papanicolaou smears: A prospective study using screening colposcopy in addition to cytology” (*JAOA* 1998;98(10):542-546) was a patient who progressed from having a negative Pap smear to invasive carcinoma in a 12-month period. A search of the literature indicated that all the algorithms used for cervical carcinoma were based on cytology, with its inherent errors, rather than on tissue diagnosis. Only in the United States is colposcopy reserved as a follow-up to the abnormal Pap smear.

The problem with the Pap smear is that unless it is used in a serial fashion, its sensitivity is too low, and the number of false-negative results is too high.² In addition, a relatively large segment of the population does not avail themselves of serial cytologic screening.³ What is needed is a more sensitive test that is clinically significant, something analogous to the American Society of Gastroenterology’s recommendation that a negative colonoscopy every 5 years can take the place of yearly sigmoidoscopy as a screen for colon carcinoma.

Without the original research on which my article was based, a cursory review of the current literature would have led me to draw the same conclusions as Dr Terry. However, a careful analysis of that literature, including the articles Dr Terry cites, indicates that the literature does not compare colposcopy with tissue diagnosis, nor is the comparison prospective. Rather, colposcopy is used as a follow-up to the abnormal Pap smear, and the majority of studies are retrospective.⁴⁻⁶ The literature also is replete with articles citing errors in tissue collection and cytology interpretation, yet our knowledge of the progression and regression of cervical dysplasia has been gleaned from this cytology—not tissue.^{7,8}

Dr Terry states that most patients did not progress to a high-grade lesion or to cervical carcinoma. Although this is true, he missed the basic points of my article. Namely, the progression of cervical dysplasia to cervical carcinoma is unknown; even under the best circumstances, the cytologic screen misses tissue-proven dys-

plasia; and prospective studies are needed to ascertain the true worth of increased sensitivity as it relates to decreasing the number of deaths from cervical carcinoma.

Dr Terry does make some valid points in his letter, especially concerning the cost analysis and identification of which subgroup would benefit from the added sensitivity of colposcopy; however, the citations and interpretations he made are not based on prospective, tissue-based research data. I am not advocating the use of colposcopy as a universal adjunct to cytologic screening. Rather, I presented data on 3000 patients that indicated further study is needed to examine the role of increased sensitivity as a means to enhance cervical carcinoma screening.

My article was not intended to be a definitive work on the merits of colposcopy in routine screening. It was meant to be a trend analysis to ascertain if the clinical impact of the increased sensitivity warrants further investigation. With a multicenter study, as was suggested in the article, perhaps the worth of a more sensitive screening procedure will be evident, and then cervical carcinoma could be screened for differently, especially among those women who cannot—or will not—submit to yearly Pap smears. As such, the death rate from cervical carcinoma would decrease. ♦

J. Robert Mannino, DO, FACFP

Department of Family Practice
Cleveland Clinical Florida
Ft Lauderdale, Fla

All opinions expressed in the JAOA are those of the authors and not necessarily those of the editors, the AOA, or the institution with which the authors are affiliated. Letters may be e-mailed to letters@aoa-net.org or they may be mailed to the Letters Editor, AOA, 142 E Ontario St, Chicago, IL 60611. No unsigned letters will be considered for publication.

JAOA

CME quiz

The purpose of this quiz is to provide a convenient means of self-assessment of your reading of the scientific content of this issue of JAOA. Enter your answers to the questions in the spaces provided so that you can easily check them with the answers that will be published next month.

To apply for CME credit, transfer your answers to the application form and answer sheet on page 84 and mail it to the CME office. So that you may complete this self-assessment in privacy, use only your member number to apply for CME credit. The CME office will record only the fact that you have completed the self-assessment test. Any grading will be done by the Editorial Department only for the purpose of planning areas of study which may be helpful to cover in future issues of JAOA.

1. Merckel's diverticulum:

- ___ (a) is congenital.
- ___ (b) is a "true" diverticulum.
- ___ (c) possesses all layers of the bowel wall.
- ___ (d) All of the above.

2. Typical symptoms of jejunal diverticulosis are:

- ___ (a) nausea.
- ___ (b) left epigastric pain.
- ___ (c) intestinal hypermotility.
- ___ (d) All of the above.

3. The so-called acute surgical abdomen

- ___ (a) is usually suggested by mild lower pelvic pain.
- ___ (b) is not associated with peritoneal irritation.
- ___ (c) may be the symptom complex associated with acute appendicitis.
- ___ (d) is not a real entity.

4. The most common complications of jejunal diverticulosis found at the time of laparotomy include:

- ___ (a) inflammation and hemorrhage.
- ___ (b) necrosis.
- ___ (c) peritoneal implants.
- ___ (d) sepsis.

5. Small bowel diverticulosis:

- ___ (a) are usually asymptomatic.
- ___ (b) are associated with ureterolithiasis.
- ___ (c) were first discovered in 1978.
- ___ (d) None of the above.

6. Signs and symptoms of jejunal diverticulosis include:

- ___ (a) diffuse abdominal pain.
- ___ (b) postprandial bloating.
- ___ (c) malabsorption.
- ___ (d) All of the above. ♦