Foreword

Leona Libby is one of the modern pioneers in climatic research. This field has been gaining more impetus in recent years, as it holds considerable promise of predicting climatic trends in future decades and centuries. Thus this book is of great interest not only to the scientist but also to economists, financial experts, politicians, and the military.

In recent years, tree ring-based temperature data have been collected which go far beyond the records available to historians. These data can be analyzed by Fourier transforms which identify certain periodicities. The longer the base data record, the more accurate the predictions are apt to be. It goes without saying that climatic changes detected by tree rings have been checked against historic records of the same time as much as that is possible today. The correspondence is astonishing. Therefore, there is considerable promise that the basic methodology is correct.

At present weather forecasting is becoming more accurate for periods on the order of days, weeks, and months. Climatic prognoses have also been attempted for very long times of tens of thousands of years. But the intermediate range in the decades and centuries so far has been an enigma. It is here where tree ring thermometry plays its trump cards.

The degree of accuracy and verification of modern decade- and century-long forecasts needs to be checked against the actual behavior of climate in the years to come. This makes long-term research projects and their funding mandatory if they are to succeed. The customary support for a few years is obviously not appropriate for this type of research, which need not be a widespread effort, however. Yet its potential is enormous in assessing worldwide crop yields, water inventory, heating requirements, stockpiling policies, and construction planning as well as political and military prospects.

So far, tree rings that have been dendrochronologically, that is, calendrically, dated reach back about 7 to 8,000 years into the past. Moreover, wood fragments or tree logs dated by radiocarbon have been found to be as old as 20 to 30,000 years. Thus there is every possibility of extending our base record far back into the Upper Pleistocene with an attendant increase in our understanding of climatic changes in the past. Using mathematics, we can translate this record into the most valuable predictions on which to base our national and international long-term policies.

Rainer Berger

Professor of Anthropology, Geography, and Geophysics
Chairman, Archaeology Program, UCLA