Preface

Carnivores have always fascinated us, even though they make up only 10% of all mammalian genera and only about 2% of all mammalian biomass. In Greek mythology most of the gods adorned their robes and helmets with depictions of carnivores, and the great hero Hercules' most famous feat was killing the "invulnerable" lion with his bare hands.

Part of our fascination with carnivores stems from fright and intrigue, and sometimes even hatred because of our direct competition with them. Cases of "man-eating" lions, bears, and wolves, as well as carnivores' reputation as killers of livestock and game, provoke communities and governments to adopt sweeping policies to exterminate them. Even President Theodore Roosevelt, proclaimer of a new wildlife protectionism, described the wolf as "the beast of waste and desolation." The sheer presence and power of carnivores is daunting; they can move quickly yet silently through forests, attaining rapid bursts of speed when necessary; their massive muscles are aligned to deliver powerful attacks, their large canines and strong jaws rip open carcasses, and their scissor-like carnassials slice meat. Partly because of our fear of these attributes, trophy hunting of carnivores has been, and to a certain extent still is, a sign of bravery and skill. Among some Alaskan Inuit, for example, a man is not eligible for marriage until he has killed a succession of animals of increasing size and dangerousness, culminating with the most menacing, the polar bear. Carnivore fur, glands, and musk are still treasured even though alternative synthetic materials have been developed. Despite our close relationship and fascination with carnivores, humans are still relatively ignorant of most species in the order. This book synthesizes some of the recent advances in research in the biology of carnivores.

The mammalian order Carnivora is characterized by great morphological, ecological, and behavioral variation. Body sizes range from the 100-g least weasel to the gigantic polar bear, weighing as much as 800 kg. Reproductive rate may be as low as one offspring every five to seven years, as in some black bears, or as high as three litters a year with eight offspring in each litter, as in

some populations of dwarf mongooses. Carnivores inhabit every habitat or vegetational zone, from short grassland (meerkat) to sparse woodland (dwarf mongoose) to desert (fennec fox) to thick forest (banded palm civet) to oceanic waters (sea otter). And the size of the home range may be relatively small (0.55 km²: coati) or extremely large and nondefensible (1500–2000 km²: African hunting dog). In terms of behavior, species range from those that live alone with only brief encounters between adults during breeding (ermine) to those that form monogamous pair bonds (golden jackal) to those that live in large extended packs with as many as 80 individuals (spotted hyena).

Although scientists have emphasized widespread interspecific differences among carnivores, considerable variation and flexibility also occur within species. In wolves, for example, adults weigh from 31 to 78 kg, litter size varies from one to 11, home ranges extend between 103 and 12,950 km², populations are found in every vegetational zone except tropical forests and arid deserts, and adults live as solitaires or within extended packs comprising up to 21 individuals. Within carnivores, therefore, we can look for examples of evolutionary adaptation at the interspecific level as well as investigate natural selection at the intraspecific level.

From the time of R. I. Pocock's and Dwight Davis's classic studies in comparative anatomy and of G. G. Simpson's pioneering work on the fossil record, the carnivores have formed the centerpiece for many studies in comparative and evolutionary biology. But until recently studies of carnivores which required detailed knowledge of their habits and lifestyles in the wild were blocked by our inability to find and track individuals easily. Because of their elusive, nocturnal, fast, solitary, and often dangerous nature, details on most species remained obscure. Today, as a result of improved research techniques (e.g., radiotelemetry and infrared spotting scopes), conservation and captive management programs, and a surge of interest in the ecological and evolutionary significance of carnivore behavior, many species are better known.

Anyone who has tried to compile comparative data on the behavior, ecology, and morphology of carnivores has quickly learned that no volume that critically summarizes and evaluates recent research in carnivore biology has been available. To meet that need, I have assembled this volume; it presents critical reviews in rapidly developing and expanding areas of carnivore behavior, ecology, and evolution. It also elucidates the features of carnivores which distinguish the group from other mammalian lineages so that non-specialists will come to know them better.

I and all the contributors feel a deep gratitude to R. F. Ewer for her monumental volume *The Carnivores* (1973, Cornell University Press), which laid the foundation for modern carnivore studies. It is testimony to the longlasting effect of her work that most of the contributors, though using very different methodologies and theoretical predictions, refer to *The Carnivores* for framing their questions.

Following a general introduction to the carnivores by John F. Eisenberg, the

volume is organized in three parts: (1) behavior—acoustic and olfactory communication; behavioral development; behavioral ecology of hyaenids and canids; modes of solitary living; and group living; (2) ecology—the feeding ecology of the giant panda and Asiatic black bear; adaptations for aquatic living; ecological constraints on predation by felids; the consequences of small size in mustelids; the rate of basal metabolism and food habits; and reproductive output; (3) evolution—morphological constraints on locomotion; dentition and diet; the physiology of delayed implantation; molecular and morphological approaches to phylogeny; and the fossil record. Each part is preceded by a brief introduction outlining the main themes presented in the chapters and explaining why certain subjects are included. Within each part, chapters proceed from specific areas to more general ones; therefore each part first deals with specific mechanisms that drive or constrain general evolutionary systems. At the end of the volume is an appendix by W. Chris Wozencraft which presents a classification of the Recent Carnivora, including the pinnipeds. This listing, which is derived from Honacki et al. (1982, *Mammal Species of the World*, Lawrence, Kans.: Association of Systematic Collections), is not meant to present a new or final word on classification; indeed, some authors have preferred the nomenclature of other taxonomies and have so stated in their text. Rather, the appendix is intended as a reference guide, modeled after that presented in Ewer's volume, for translating species names into common ones (or vice versa), for showing familial membership of species, and for pointing out general taxonomic changes made in recent years.

Because this volume covers a broad taxonomic group and includes many disparate research topics, several editorial decisions were necessary. The first question addressed was general: what is a carnivore, or which species should be included in a book dealing with the Carnivora? There is historical precedent either for combining the terrestrial carnivores and aquatic pinnipeds in the order Carnivora or for splitting off the terrestrial carnivores to form an independent, ordinal group. Studies of the origin and phylogeny of carnivores and pinnipeds continue to produce lively debate, as chapters in this volume attest. The chapters deal mainly with the terrestrial (fissiped) carnivores or species included in the following taxonomic families: Canidae (wolves, wild dogs, jackals, foxes), Procyonidae (raccoons, coatis, kinkajou), Ursidae (bears), Ailuropodidae (giant panda), Ailuridae (red panda), Mustelidae (weasels, martens, fisher, tayra, ratel, badgers, skunks, otters), Viverridae (civets, genets), Herpestidae (mongooses), Hyaenidae (hyenas, aardwolf), and Felidae (wild cats, ocelots, serval, caracal, lynxes, puma, leopards, jaguar, lion, tiger, cheetah). I have thus excluded the pinnipeds from discussion; the behavioral, ecological, and evolutionary features of adaptations for aquatic living set this group apart from the terrestrial carnivores. A further, more practical reason is that other volumes nicely synthesize recent advances in research on the pinnipeds. By considering only the terrestrial carnivores, this book avoids duplicating other publications.

Even when the taxonomic limitations were set, another potential source of confusion remained. The word “carnivore” in everyday language refers to a dietary proclivity for flesh eating. Thus, describing a species as a carnivore could refer to its taxonomic affiliation or to its dietary preference. Because the general focus of this volume is on species taxonomically included as the terrestrial carnivores, “carnivore” refers to the taxonomic usage unless otherwise stated. For example, where diet is analyzed, some ursids are described as being herbivorous carnivorans or simply herbivores.

My primary criteria in making these decisions was consistency and defensibility of argument throughout—that is, questions regarding taxonomy, priority of theoretical questions, method of citations, and other recurring features across chapters were to be presented in the same fashion and as rigorously as possible in each context so that at least ambiguity would be avoided. Because consistency was a goal, however, the problem of overlap between chapters became an issue. Even though repetition was eliminated whenever possible, some overlap had to remain if each chapter was to stand on its own. Such repetition reduces the burden of cross-referencing between chapters. Whenever chapters deal with similar subjects, one chapter serves as the main source for a given set of data or particular theoretical perspective. When other chapters refer to the same material, a brief summary is given, along with cross-references to the chapter(s) where more detailed treatment occurs.

Another editorial decision concerned the presentation of unorthodox views, unpublished information, or disagreement among review chapters, which are meant to present only facts and references. And indeed, the data compilations, descriptions of field or experimental studies, and theoretical discussions in most chapters are based on previously published, refereed work, but some chapters include original data or theoretical arguments that offer a unique perspective. I decided that these views should be aired also, so that future work in carnivore biology might more rapidly test their validity. Nevertheless, in cases where new information is presented, I have required the authors to refer to background literature to introduce the conceptual problem or new methodology and to elucidate why these new data are important in the broad context here, wherein previous studies already established in a particular area are being reviewed.

It is my hope that this collection of chapters brings the excitement and beauty of carnivores closer to those who have not had the opportunity to study them, especially to students in the behavioral, ecological, and evolutionary sciences who are looking for a diverse and intriguing group to work on. And, for my scientific colleagues, it is my intention that the problems and questions raised by taking stock of what we have learned about carnivores will spur us on to discover what we need to know in order to preserve them.

JOHN L. GITTLEMAN

Knoxville, Tennessee