

Contents

Foreword — VII

Author's Preface — VIII

1 Introduction — 1

- 1.1 Origin of EMR — 1
- 1.2 Experimental apparatus — 3
 - 1.2.1 Microwave source — 4
 - 1.2.2 Resonant cavity and coupling system — 5
 - 1.2.3 Magnet — 10
 - 1.2.4 Detection system — 12
 - 1.2.5 Data treatment system — 12
- 1.3 Target of research — 13
- 1.4 Prospects for future — 13
- References — 14

2 Theoretical basics — 16

- 2.1 Phenomenal description of EMR — 16
- 2.2 Angular momentum and magnetic moment — 17
 - 2.2.1 Orbital motion of electron and its magnetic moment — 17
 - 2.2.2 Eigen motion of electrons and its magnetic moment — 20
 - 2.2.3 Spin angular momentum and magnetic moment of atomic nucleus — 23
 - 2.2.4 Electric quadrupole moment of atomic nucleus — 25
- 2.3 Unit of magnetic field — 26
- 2.4 The interaction between external fields and magnetic moment — 28
- 2.5 Interaction of magnetic moment with electromagnetic field in the external magnetic field — 30
- 2.6 Interaction of nuclear magnetic moment with electron magnetic moment in the external magnetic field — 33
 - References — 34
 - Further reading — 34

3 *g*-Tensor theory — 35

- 3.1 Landé factor — 35
- 3.2 Matrix presentation of *g*-tensor — 37
 - 3.2.1 *g*-Tensor of colour center (cubic symmetry and uniaxial symmetry system) — 37
 - 3.2.2 The *g*-tensor of nonaxisymmetric (lower than uniaxial symmetry) system — 39

- 3.3 *g*-Tensor of irregular orientation system — 46
- 3.3.1 *g*-Tensor of axisymmetric system — 46
- 3.3.2 *g*-Tensor of nonaxisymmetric system — 50
- References — 52
- Further reading — 52

- 4 Isotropic hyperfine structure — 53**
 - 4.1 Theoretical exploration of hyperfine interaction — 53
 - 4.1.1 Dipole–dipole interaction — 53
 - 4.1.2 Fermi contact interaction — 54
 - 4.2 Energy operator of isotropic hyperfine interaction — 55
 - 4.2.1 Spin operator and hamiltonians — 55
 - 4.2.2 Zeeman interaction of electrons and nuclei — 57
 - 4.2.3 Spin hamiltonian of isotropic hyperfine interaction — 59
 - 4.3 Spectral isotropic hyperfine structure — 60
 - 4.3.1 System with one magnetic nucleus and one unpaired electron — 60
 - 4.3.2 Multimagnetic nuclei with one unpaired electron system — 65
 - 4.3.3 Hyperfine splitting arising from other magnetic nuclei — 71
 - 4.3.4 Encountered problems in isotropic radical spectra — 75
 - 4.4 Hyperfine structure of organic π -free radical spectrum — 76
 - 4.4.1 Hyperfine coupling constant of organic π -radical — 76
 - 4.4.2 McConnell semiempirical formula — 76
 - 4.4.3 Hückel molecular orbital (HMO) theory — 77
 - 4.4.4 Calculation of probability density distribution of unpaired electron — 79
 - 4.4.5 The Q value of the radical with fully symmetrical structure — 85
 - 4.4.6 Hyperfine coupling constant a value of the even alternant hydrocarbons — 86
 - 4.4.7 Hyperfine coupling constant a value of the even alternant heterocyclic hydrocarbons — 89
 - 4.4.8 Hyperfine coupling constant a value of the odd alternant and nonalternant hydrocarbons — 90
 - 4.5 Mechanism of hyperfine splitting in the spectrum of conjugated systems — 91
 - 4.5.1 “Electronic correlation” effect — 91
 - 4.5.2 The sign of proton hyperfine splitting constant — 93
 - 4.5.3 Negative spin density — 96
 - 4.5.4 About the Q value problem — 96
 - 4.5.5 Hyperfine splitting and hyperconjugation effect of methyl protons — 98
 - 4.6 Hyperfine splitting of other (non-proton) nuclei — 101
 - 4.6.1 Hyperfine splitting of ^{13}C nucleus — 101

- 4.6.2 Hyperfine splitting of ^{14}N nucleus — **103**
- 4.6.3 Hyperfine splitting of ^{19}F nucleus — **104**
- 4.6.4 Hyperfine splittings of ^{17}O and ^{33}S nuclei — **105**
- References — **105**
- Further readings — **106**

- 5 Anisotropic hyperfine structure — 107**
- 5.1 Anisotropic hyperfine interaction — **107**
- 5.2 Matrix interpretation of anisotropic hyperfine interaction — **110**
- 5.3 Example demonstration — **116**
- 5.4 Anisotropic hyperfine coupling tensor and structure of radical — **122**
- 5.4.1 Hyperfine coupling tensor of central atom — **122**
- 5.4.2 Hyperfine coupling tensor of α -hydrogen atom — **127**
- 5.4.3 Hyperfine coupling tensor of β -hydrogen atom — **130**
- 5.4.4 Hyperfine coupling tensor of σ -type organic radicals — **132**
- 5.5 Anisotropy of the combination of g -tensor and A -tensor — **133**
- 5.6 Anisotropy of A tensor in the irregular orientation system — **133**
- References — **135**
- Further readings — **136**

- 6 Fine structure — 137**
- 6.1 Zero-field splitting — **138**
- 6.2 Spin hamiltonian of two-electron interaction — **140**
- 6.2.1 Exchange interaction of electron spin — **140**
- 6.2.2 Dipole interaction of electron–electron — **144**
- 6.3 The triplet molecule ($S = 1$) system — **153**
- 6.3.1 Energy levels and wave functions of triplet molecules under the action of external magnetic field — **153**
- 6.3.2 Examples of triplet state excited by light — **156**
- 6.3.3 Examples of thermal excitation triplet state — **157**
- 6.3.4 Examples of other excited triplet — **159**
- 6.3.5 Examples of ground triplet state — **159**
- 6.4 Triplet system of irregular orientation — **162**
- 6.5 Biradical — **166**
- References — **169**
- Further reading — **170**

- 7 Relaxation and line shape and linewidth — 171**
- 7.1 Model of spin relaxation — **171**
- 7.1.1 Spin temperature and boltzmann distribution — **171**
- 7.1.2 Spin particle transition dynamics — **173**
- 7.1.3 Mechanism of the effect of relaxation time τ_1 on linewidth — **175**

- 7.1.4 Magnetization in static magnetic field — **177**
- 7.1.5 Bloch equation in the static magnetic field — **178**
- 7.1.6 Bloch equation in the static magnetic field coupled with the oscillating magnetic field — **180**
- 7.1.7 Stationary solutions of bloch equation — **181**
- 7.2 Shape, width, and intensity of spectral line — **182**
 - 7.2.1 Line shape function — **182**
 - 7.2.2 Linewidth — **186**
 - 7.2.3 Line broadening — **188**
 - 7.2.4 Line intensity — **188**
- 7.3 Dynamic effects of line shape — **189**
 - 7.3.1 Generalized bloch equation — **189**
 - 7.3.2 Chemical exchange broadening mechanism — **193**
 - 7.3.3 Mechanism of the spectral lines broadening caused by physical motion — **200**
- 7.4 Saturation transfer of spectra — **214**
- 7.5 Intensity of signal dependent on time — **214**
 - 7.5.1 Free radical concentration changes with time — **214**
 - 7.5.2 Chemical-induced dynamic electron polarization (CIDEP) — **215**
- References — **217**
- Further readings — **218**

- 8 Quantitative determination — 220**
 - 8.1 Main factors of influence for quantitative determination — **221**
 - 8.1.1 Factors of instrument — **221**
 - 8.1.2 Influence of operating factors — **228**
 - 8.2 Selection and preparation of standard samples — **235**
 - 8.3 Key parameters and its effect on the intensity of EMR signal — **237**
 - 8.4 Achievable accuracy of quantitative determination — **239**
 - References — **240**

- 9 Paramagnetic gases and inorganic radicals — 242**
 - 9.1 Spectra of paramagnetic gases — **242**
 - 9.1.1 Monoatomic paramagnetic gases — **242**
 - 9.1.2 Diatomic paramagnetic gas — **245**
 - 9.1.3 Gaseous molecules of triatom and polyatom — **256**
 - 9.2 Expanding of EMR technique for study on paramagnetic gas — **257**
 - 9.2.1 Laser electronic magnetic resonance — **257**
 - 9.2.2 Magnetic resonance induced by electron — **257**
 - 9.3 Inorganic radicals — **258**
 - 9.4 Point defects in solid states — **261**
 - 9.5 Spectra of conductor and semiconductor — **265**

- 9.6 Structure of a molecule structure of a molecule estimated from the data of EMR — 268
References — 269
Further readings — 272

10 Ions of transition elements and their complexes — 273

- 10.1 Electron ground state of transition element ion — 273
10.2 Orbital degeneracy is rescinded in ligand field — 275
10.3 Electric potential of ligand field — 278
10.4 Energy-level splitting of transition metal ion in ligand field — 282
10.4.1 *P*-state ion in octahedron Field ($L = 1$) — 282
10.4.2 *D*-state ion — 283
10.4.3 About *F*-state ion — 286
10.5 Spin-orbit coupling and spin hamiltonian — 288
10.6 Ground-state ion with orbital nondegeneracy — 294
10.6.1 *D*-state ions of ground-state orbital nondegenerate — 295
10.6.2 *F*-state ions of ground-state orbital nondegenerate — 298
10.6.3 *S*-state ions of the ground-state orbital nondegenerate — 305
10.7 Ground-state ions with orbital degeneracy — 310
10.7.1 *D*-state ions — 310
10.7.2 *F*-state ions — 316
10.7.3 Jahn-Teller distortion — 320
10.7.4 The palladium group (4d) and platinum group (5d) ions — 321
10.8 EMR spectra of rare earth ions — 321
10.8.1 Lanthanide ion — 321
10.8.2 Actinide ions — 323
10.9 EMR spectra of transition metal complexes — 324
References — 325
Further readings — 327

Appendix 1: Extension and expansion of EMR — 328

Appendix 2: Mathematical preparation — 399

Appendix 3: Angular momentum and stable-state perturbation theory in quantum mechanics — 421

Appendix 4: Fundamental constants and useful conversion actors — 442

Appendix 5: The natural abundance, nuclear spin, nuclear magnetogyric ratio of some magnetic nuclei and their hyperfine coupling parameters — 445

Index — 451

