

Contents

Preface — IX

Introduction — XI

- 1 The 8π -problem in the two dimensional case — XIV
- 2 The parabolic-elliptic Keller–Segel model in higher dimensions — XVIII
- 3 Chemotaxis models with nonlocal diffusion operators — XIX
- 4 Notations — XX
- 5 Preliminaries on semigroups for linear diffusion equations — XXI

1 Local-in-time solutions, small global-in-time solutions — 1

- 1.1 Classical Keller–Segel system — 1
- 1.2 Fractional Keller–Segel model and optimal initial data — 6
- 1.3 Singular stationary solution — 6
- 1.4 Local and global solutions with initial data in Morrey space $M^{d/\alpha}(\mathbb{R}^d)$ — 9
- 1.5 Other simple constructions of solutions in nearly optimal spaces — 13
- 1.6 Notes and complements — 19

2 Large global-in-time solutions to models of chemotaxis — 21

- 2.1 The case of classical diffusion — 21
- 2.2 The case of a nonlocal model of chemotaxis — 30
- 2.3 Statement of results for $\alpha \in (0, 2)$ — 33
- 2.4 Pointwise comparison principle — 35
- 2.5 Averaged comparison principle — 43
- 2.6 Construction of global-in-time solutions for $\alpha \in (0, 1)$ — 49
- 2.7 Construction of the unique global-in-time solutions for $\alpha \in (1, 2)$ — 55
- 2.8 Large global-in-time solutions of the parabolic-parabolic Keller–Segel system on the plane — 58
- 2.9 Comments on special solutions — 66

3 Blowups — 73

- 3.1 Solutions blowing up in a finite time — 73
- 3.2 A novel approach to blowup and concentration of mass in two dimensions — 85
- 3.3 Proof of blowup of radial solutions — 89
- 3.4 Blowup in the plane \mathbb{R}^2 with 8π concentration of mass — 91
- 3.5 Refinements of the proof of blowup of radial solutions in higher dimensions — 104
- 3.6 Blowup of solutions for the two dimensional fractional diffusion Keller–Segel model — 109

3.7	Keller–Segel model with chemoattractant consumption terms —	115
3.8	Blowup of radially symmetric solutions, again —	131
4	Blowups à la Fujita —	137
4.1	Fujita’s idea of the proof of blowup for the system with the classical diffusion —	137
4.2	Blowup à la Fujita of solutions of system with fractional diffusion —	145
5	Interpretations, complements, conjectures, et cetera —	153
5.1	Dichotomy: local–global existence of solutions —	153
5.2	Further comments on solvability of the classical and fractional Keller–Segel system —	154
5.3	Optimal initial data for two dimensional Keller–Segel system —	157
5.4	Blowup of solutions to a general diffusive aggregation model —	165
5.5	Hypercontractivity of the linearization around the singular solution —	177
5.6	Classical diffusion case: super- and subsolutions —	181
5.7	Miscellanea, questions and conjectures, research plans —	188
	Bibliography —	191
	Index —	203