

Ceramides in the skin barrier

Original Paper

Vávrová K.,[✉] Kováčik A., Opálka L.

Charles University, Faculty of Pharmacy in
Hradec Králové, Skin Barrier Research Group,
Hradec Králové, Czech Republic

Received 31 October, 2016, accepted 7 December, 2016

Abstract The skin barrier, which is essential for human survival on dry land, is located in the uppermost skin layer, the stratum corneum. The stratum corneum consists of corneocytes surrounded by multilamellar lipid membranes that prevent excessive water loss from the body and entrance of undesired substances from the environment. To ensure this protective function, the composition and organization of the lipid membranes is highly specialized. The major skin barrier lipids are ceramides, fatty acids and cholesterol in an approximately equimolar ratio. With hundreds of molecular species of ceramide, skin barrier lipids are a highly complex mixture that complicate the investigation of its behaviour. In this minireview, the structures of the major skin barrier lipids, formation of the stratum corneum lipid membranes and their molecular organization are described.

Keywords Skin barrier – stratum corneum – ceramide – sphingolipid

FUNDING

This work was supported by the Czech Science Foundation – German Science Foundation joint project 16-25687J.

INTRODUCTION

Ceramides, that is, *N*-acylsphingosines belong to the family of sphingolipids. Ceramides regulate several cellular processes, such as proliferation, differentiation and apoptosis (Hannun, 1996; Hannun and Obeid, 2008). In human body, ceramides are equally important outside the cell – they form multilamellar lipid membranes between corneocytes of the uppermost skin layer, the stratum corneum (Candi et al., 2005). These stratum corneum lipids prevent excessive water loss from the body and hamper the penetration of undesired substances, allergens and microbes from the environment into the body (Bouwstra and Ponc, 2006; Elias, 2008). In fact, the development of such lipid-based barrier was an essential step in the evolution of life on dry land. In this minireview, the structures of major skin barrier lipids, formation of the stratum corneum lipid membranes and their molecular organization are described.


COMPOSITION OF THE SKIN LIPID BARRIER

Human stratum corneum is composed of approximately 15–25 layers of flattened dead cells, corneocytes, which are surrounded by a lipid matrix (Elias et al., 1977). To provide a competent barrier function, the lipids in the stratum corneum are highly specialized: there are virtually no phospholipids; the major constituents are ceramides, free fatty acids and cholesterol in an approximately 1:1:1 molar ratio (**Fig. 1**). This equimolar ratio seems to be highly important for epidermal homeostasis, as alterations in this lipid composition were associated with a disturbed barrier function (Breiden and Sandhoff, 2014; Feingold and Elias, 2014; Holleran et al., 2006; Rabionet et al., 2014; van Smeden et al., 2014b). Free fatty acids in the skin barrier are 14–34 carbons long, mostly saturated and unbranched. The predominant fatty acids have 24 carbons (lignoceric acid) and 26 carbons (cerotic acid) – these two species comprise approximately 50 weight % of the stratum corneum fatty acids (Norlen et al., 1998; van Smeden et al., 2014a). Unsaturated and shorter fatty acids are present in the stratum corneum lipid membranes only in minor quantities; their concentrations increase in pathological conditions such as atopic dermatitis, probably due to flaggrin

* E-mail: katerina.vavrova@faf.cuni.cz

© European Pharmaceutical Journal

Epidermis:



Major skin barrier lipids:




Figure 1. Schematic representation of epidermal layers, structure of the stratum corneum and the major skin barrier lipids.

deficiency (Jakasa et al., 2011; Jungersted et al., 2010; Vavrova et al., 2014).

Cholesterol is the main sterol in the skin lipid barrier. Cholesterol is also essential for the correct skin barrier function; it contributes to the correct lamellar and lateral structure and lipid fluidity (Feingold and Elias, 2014; Mojumdar et al., 2015a). Previous studies described separation of cholesterol from the lamellar phases (Mojumdar et al., 2015b; Schreiner et al., 2000); however, the physiological relevance of this separation is not known. In contrast, recent study using very high resolution electron microscopy on vitreous skin sections found no cholesterol crystals (Iwai et al., 2012).

Cholesteryl sulphate, that is, ester of sulphuric acid with cholesterol, is a minor component of the stratum corneum lipid matrix; it is present in approximately 5% of the barrier lipids (or approximately 2% of all lipids including surface lipids (Lampe et al., 1983)). The role of cholesteryl sulphate is probably not fully understood, one possible function of this molecule is that it contributes to the stratum corneum cohesion and regulates desquamation (Long et al., 1985).

CHEMISTRY OF CERAMIDES

The dominant lipids in the stratum corneum are ceramides; they constitute approximately one third of lipid molecules or 50% of the lipids by weight (van Smeden et al., 2014b). These simple sphingolipids have a relatively small polar head that does not extensively hydrate and two hydrophobic chains – the sphingoid base chain and *N*-acyl chain (fatty acid chain). The most intriguing fact in skin ceramides is their heterogeneity: 15 ceramide subclasses have been described until the present and together with the different chain lengths, this yields hundreds of distinct chemical structures (Fig. 2). For a recent review, see (Rabionet et al., 2014).

All ceramides are based on a sphingoid base, which is a long chain (usually 18 carbons) amino alcohol. The skin ceramides can contain sphingosine (S) and dihydrosphingosine (dS), which are common in eukaryotic cells, phytosphingosine (P), which is found only in some human tissues, and 6-hydroxysphingosine (H), which is specific for the epidermis (Kováčik et al., 2014). Recently, a novel tetrahydroxylated sphingoid base, which




Figure 3. Lamellar and lateral organization of the stratum corneum lipid membranes.

first lipid layer covalently anchored to the corneocyte surface (Breathnach, 1975; Grayson and Elias, 1982). In contrast to phospholipids, the stratum corneum ceramides have a small polar head and very long, saturated chains with prevailing *all-trans* conformation. The prevalent lateral packing of the skin barrier lipids is orthorhombic, which is a very tight packing with highly limited motional freedom. In addition, some lipids are arranged in a hexagonal subcell with greater rotational freedom, and a minor lipid fraction is liquid (Fig. 3) (Bouwstra et al., 2001; Damien and Boncheva, 2010; Mendelsohn et al., 2006; Mendelsohn and Moore, 2000; Stahlberg et al., 2015). Considering the lamellar arrangement of the stratum corneum lipid multilayers, their unique feature is the formation of the so-called long periodicity lamellar phase with a repeat distance of 11.9–13.1 nm (Hou et al., 1991; Madison et al., 1987; Mojumdar et al., 2015a; Neto et al., 2011; White et al., 1988) (Fig. 3). This lamellar phase was first observed by electron microscopy and later confirmed by X-ray diffraction. In addition, most studies also describe the presence of a short periodicity phase (with repeat distance of 5.3–6.4 nm) (Bouwstra et al., 1995; Bouwstra et al., 1991; de Jager et al., 2004; de Sousa Neto et al., 2011; Mojumdar et al., 2014; White et al., 1988) and phase separated cholesterol (Craven, 1979; Shieh et al., 1981). Nevertheless, recent experiments on native skin barrier using very high resolution cryo-electron microscopy only detected the long periodicity phase, with the repeat distance of approximately 11 nm (Iwai et al., 2012). There is a consensus that acylceramides (that is, ceramides of the EO-subclass) are necessary for the formation of the long periodicity phase (Bouwstra et al., 1991; de Sousa Neto et al., 2011; Kessner et al., 2010; Madison et al., 1987; Mojumdar et al., 2015a; White et al., 1988). However, recent X-ray diffraction

study showed a lamellar phase with 10.6 nm periodicity in ceramide NH membranes without any acylceramide present (Kovacik et al., 2016).

Because of the multilamellar arrangement of lipids in the stratum corneum, both hairpin conformation (with both chains pointing to the same direction) and extended conformation of ceramide (or splayed-chain conformation, with the chains pointing into opposite directions) are theoretically possible (Fig. 4). Although the extended conformation is not common in biological membranes, it could be advantageous for the stratum corneum barrier properties (Corkery, 2002; Iwai et al., 2012). First, it would reduce the packing strain of ceramide because they have a smaller cross-section of the polar head than of the chains and there are no lipids with bulky polar heads to compensate for such unfavourable packing ratio of hairpin ceramide. Second, the extended conformation would connect the adjacent lipid lamellae and prevent permeable boundaries.

STRATUM CORNEUM LIPID MEMBRANE MODELS

The exact lipid organization in the stratum corneum has been under debate for decades (Norlen, 2013). Both lamellar phases and the lateral packing can be reproduced using the in vitro lipid membrane models (Kessner et al., 2008). Numerous previous studies using either lipid membrane models or stratum corneum described how ceramide structure influences their behaviour in membranes and permeability (Janusova et al., 2011; Novotný et al., 2009; Pullmannová et al., 2014; Rerek et al., 2001; Skolova et al., 2014; Skolova et al., 2016; Skolova et al., 2013; Stahlberg et al., 2016; Stahlberg et al., 2015) and proposed models of the arrangement (Bouwstra




Figure 4. Two possible conformations of ceramide in the skin barrier membranes.

et al., 1991; de Jager et al., 2004; de Sousa Neto et al., 2011; Iwai et al., 2012; Mojumdar et al., 2015a; Mojumdar et al., 2015b; Mojumdar et al., 2014). However, no broad consensus has been reached, mainly because of the differences in the experimental setup, method used and, in case of models, lipid composition.

These lipid models always bear a certain level of simplification but some of them showed very good correlations with real skin properties. For example, de Jager *et al.* studied a model lipid membrane composed of ceramides EOS, NS24, NP24, AS24, NP16, AP24, free fatty acid mixture and cholesterol and found similar permeabilities of this model and isolated stratum corneum to *p*-aminobenzoic acid and its esters (de Jager et al., 2006). We compared the effects of ceramide with shortened chains (both acyl and sphingosine) in skin and simple lipid membranes composed of ceramide NS24, lignoceric acid, cholesterol and cholesteryl sulphate and found similar trends in permeabilities to the two model permeants theophylline and indomethacin (Janušová et al., 2011; Školová et al., 2016; Školová et al., 2013). Furthermore, a lipid model composed of ceramide NS24, free fatty acids and cholesterol (with or without cholesteryl sulphate) confirmed the existence of fully extended ceramide with cholesterol molecules associated with the ceramide sphingoid moiety and free fatty acids with the acyl chains (Školová et al., 2014) suggested by very high resolution cryo-electron microscopy of native skin (Iwai et al., 2012). The permeabilities of our recent complex model with ceramides EOS, EoS and EOP, NS, NdS, NP, AS, AdS and AP, free fatty acids, cholesterol and cholesteryl sulphate (the flux values of theophylline and indomethacin were $0.24 \pm 0.02 \mu\text{g}/\text{cm}^2/\text{h}$ and $0.12 \pm 0.02 \mu\text{g}/\text{cm}^2/\text{h}$, (Opálka et al., 2016)) are in very good agreement with those of a membrane model constructed from isolated human skin ceramides (the flux values of theophylline and indomethacin were $0.40 \pm 0.05 \mu\text{g}/\text{cm}^2/\text{h}$ and $0.23 \pm 0.02 \mu\text{g}/\text{cm}^2/\text{h}$, respectively

(Pullmannová et al., 2014)) and also with permeabilities of isolated human epidermis under the same conditions (the flux values of theophylline and indomethacin were $0.15 \pm 0.05 \mu\text{g}/\text{cm}^2/\text{h}$ and $0.05 \pm 0.02 \mu\text{g}/\text{cm}^2/\text{h}$, respectively – Kováčik, unpublished data).

The main drawback of the model lipid membranes is that some ceramides are not commercially available, especially those of the EO-subclass and H-subclass. However, synthetic procedures to these lipids have recently been published (Kovacik et al., 2016; Mori and Matsuda, 1991; Muller and Schmidt, 2000; Opálka et al., 2015) and opened the possibility to further improve the stratum corneum lipid models to better mimic the real skin lipid barrier.

CONCLUSION

Ceramides are essential for our life on dry land. Major disturbances to their biosynthesis or transport could be lethal because of an extensive water loss. Even less pronounced alterations of the stratum corneum lipid composition and organization are associated with skin diseases such as atopic dermatitis, psoriasis and ichthyoses and could greatly affect the quality of life of the patients. Although topical application of skin-identical lipids or their analogues has a strong potential to repair the skin barrier in such diseases and prevent inflammation, deeper understanding of the physical chemistry of ceramides and the molecular arrangement of the skin lipid membranes is necessary for more rational design of such therapies.

ACKNOWLEDGEMENTS

This work was supported by the Czech Science Foundation-German Science Foundation joint project 16-2568J.

References

- [1] Banks-Schlegel S, Green H. Involucrin synthesis and tissue assembly by keratinocytes in natural and cultured human epithelia. *J Cell Biol.* 1981;90:732-737.
- [2] Behne M, Uchida Y, Seki T, de Montellano PO, Elias PM, Holleran WM. *J. Invest. Dermatol.* 2000;114:185.
- [3] Bouwstra JA, Gooris GS, Bras W, Downing DT. Lipid organization in pig stratum corneum. *J. Lipid. Res.* 1995;36:685-695.
- [4] Bouwstra JA, Gooris GS, Dubbelaar FE, Ponc M. Phase behavior of lipid mixtures based on human ceramides: coexistence of crystalline and liquid phases. *J. Lipid. Res.* 2001;42:1759-1770.
- [5] Bouwstra JA, Gooris GS, van der Spek JA, Bras W. Structural investigations of human stratum corneum by small-angle X-ray scattering. *J. Invest. Dermatol.* 1991;97:1005-1012.
- [6] Bouwstra JA, Ponc M. The skin barrier in healthy and diseased state. *Biochim. Biophys. Acta.* 2006;1758:2080-2095.
- [7] Breathnach AS. Aspects of epidermal ultrastructure. *J. Invest. Dermatol.* 1975;65:2-15.
- [8] Breiden B, Sandhoff K. The role of sphingolipid metabolism in cutaneous permeability barrier formation. *Biochim Biophys Acta.* 2014;1841:441-452.
- [9] Candi E, Schmidt R, Melino G. The cornified envelope: a model of cell death in the skin. *Nat. Rev. Mol. Cell Biol.* 2005;6:328-340.
- [10] Corkery RW. The anti-parallel, extended or splayed-chain conformation of amphiphilic lipids. *Colloids Surf B Biointerfaces.* 2002;26:3-20.
- [11] Craven B. Pseudosymmetry in cholesterol monohydrate. *Acta Crystallogr Sect B.* 1979;35:1123-1128.
- [12] Damien F, Boncheva M. The extent of orthorhombic lipid phases in the stratum corneum determines the barrier efficiency of human skin in vivo. *J. Invest. Dermatol.* 2010;130:611-614.
- [13] de Jager M, Gooris G, Ponc M, Bouwstra J. Acylceramide head group architecture affects lipid organization in synthetic ceramide mixtures. *J. Invest. Dermatol.* 2004;123:911-916.
- [14] de Jager M, Groenink W, i Guivernau RB, et al. A novel in vitro percutaneous penetration model: evaluation of barrier properties with p-aminobenzoic acid and two of its derivatives. *Pharmaceut. Res.* 2006;23:951-960.
- [15] de Sousa Neto D, Gooris G, Bouwstra J. Effect of the omega-acylceramides on the lipid organization of stratum corneum model membranes evaluated by X-ray diffraction and FTIR studies (Part I). *Chem Phys Lipids.* 2011;164:184-195.
- [16] Elias PM. Skin barrier function. *Curr. Allergy Asthma Rep.* 2008;8:299-305.
- [17] Elias PM, Goerke J, Friend DS. Mammalian Epidermal Barrier Layer Lipids: Composition and Influence on Structure. *J. Invest. Dermatol.* 1977;69:535-546.
- [18] Elias PM, Gruber R, Crumrine D, et al. Formation and functions of the corneocyte lipid envelope (CLE). *Biochim Biophys Acta.* 2014;1841:314-318.
- [19] Feingold KR, Elias PM. Role of lipids in the formation and maintenance of the cutaneous permeability barrier. *Biochim. Biophys. Acta.* 2014;1841:280-294.
- [20] Grayson S, Elias PM. Isolation and Lipid Biochemical Characterization of Stratum Corneum Membrane Complexes: Implications for the Cutaneous Permeability Barrier. *J. Invest. Dermatol.* 1982;78:128-135.
- [21] Hannun YA. Functions of ceramide in coordinating cellular responses to stress. *Science.* 1996;274:1855-1859.
- [22] Hannun YA, Obeid LM. Principles of bioactive lipid signalling: lessons from sphingolipids. *Nat. Rev. Mol. Cell Biol.* 2008;9:139-150.
- [23] Holleran WM, Takagi Y, Uchida Y. Epidermal sphingolipids: metabolism, function, and roles in skin disorders. *FEBS Lett.* 2006;580:5456-5466.
- [24] Hou SY, Mitra AK, White SH, Menon GK, Ghadially R, Elias PM. Membrane structures in normal and essential fatty acid-deficient stratum corneum: characterization by ruthenium tetroxide staining and x-ray diffraction. *J. Invest. Dermatol.* 1991;96:215-223.
- [25] Iwai I, Han H, den Hollander L, et al. The human skin barrier is organized as stacked bilayers of fully extended ceramides with cholesterol molecules associated with the ceramide sphingoid moiety. *J. Invest. Dermatol.* 2012;132:2215-2225.
- [26] Jakasa I, Koster ES, Calkoen F, et al. Skin barrier function in healthy subjects and patients with atopic dermatitis in relation to filaggrin loss-of-function mutations. *J. Invest. Dermatol.* 2011;131:540-542.
- [27] Janusova B, Zbytovska J, Lorenc P, et al. Effect of ceramide acyl chain length on skin permeability and thermotropic phase behavior of model stratum corneum lipid membranes. *Biochim Biophys Acta.* 2011;1811:129-137.
- [28] Janůšová B, Zbytovská J, Lorenc P, et al. Effect of ceramide acyl chain length on skin permeability and thermotropic phase behavior of model stratum corneum lipid membranes. *BBA-Mol. Cell Biol. L.* 2011;1811:129-137.
- [29] Jennemann R, Rabionet M, Gorgas K, et al. Loss of ceramide synthase 3 causes lethal skin barrier disruption. *Hum Mol Genet.* 2012;21:586-608.
- [30] Jensen JM, Schutze S, Forl M, Kronke M, Proksch E. Roles for tumor necrosis factor receptor p55 and sphingomyelinase in repairing the cutaneous permeability barrier. *J. Clin. Invest.* 1999;104:1761-1770.
- [31] Jungersted JM, Scheer H, Mempel M, et al. Stratum corneum lipids, skin barrier function and filaggrin mutations in patients with atopic eczema. *Allergy.* 2010;65:911-918.
- [32] Kessner D, Brezesinski G, Funari SS, Dobner B, Neubert RH. Impact of the long chain omega-acylceramides on the stratum corneum lipid nanostructure. Part 1: Thermotropic phase behaviour of CER[EOS] and CER[EOP] studied using X-ray powder diffraction and FT-Raman spectroscopy. *Chem Phys Lipids.* 2010;163:42-50.
- [33] Kessner D, Ruettinger A, Kiselev MA, Wartewig S, Neubert RH. Properties of ceramides and their impact on the stratum corneum structure. Part 2: stratum corneum lipid model systems. *Skin Pharmacol. Physiol.* 2008;21:58-74.
- [34] Kovacic A, Opalka L, Silarova M, Roh J, Vavrova K. Synthesis of 6-hydroxyceramide using ruthenium-catalyzed hydrosilylation-protodesilylation. Unexpected formation of a long periodicity lamellar phase in skin lipid membranes. *RSC Adv.* 2016;6:73343-73350.

- [35] Kováčik A, Roh J, Vávrová K. The chemistry and biology of 6 hydroxyceramide, the youngest member of the human sphingolipid family. *ChemBioChem*. 2014;15:1555–1562.
- [36] Lampe MA, Burlingame A, Whitney J, et al. Human stratum corneum lipids: characterization and regional variations. *JLipid Res*. 1983;24:120-130.
- [37] Long SA, Wertz PW, Strauss JS, Downing DT. Human stratum corneum polar lipids and desquamation. *Arch Dermatol Res*. 1985;277:284-287.
- [38] Madison KC, Swartzendruber DC, Wertz PW, Downing DT. Presence of intact intercellular lipid lamellae in the upper layers of the stratum corneum. *J. Invest. Dermatol*. 1987;88:714-718.
- [39] Masukawa Y, Narita H, Shimizu E, et al. Characterization of overall ceramide species in human stratum corneum. *J. Lipid. Res*. 2008;49:1466-1476.
- [40] Mendelsohn R, Flach CR, Moore DJ. Determination of molecular conformation and permeation in skin via IR spectroscopy, microscopy, and imaging. *Biochim Biophys Acta*. 2006;1758:923-933.
- [41] Mendelsohn R, Moore DJ. Infrared determination of conformational order and phase behavior in ceramides and stratum corneum models. *Methods Enzymol*. 2000;312:228-247.
- [42] Mizutani Y, Mitsutake S, Tsuji K, Kihara A, Igarashi Y. Ceramide biosynthesis in keratinocyte and its role in skin function. *Biochimie*. 2009;91:784-790.
- [43] Mojumdar EH, Gooris GS, Barlow DJ, Lawrence MJ, Deme B, Bouwstra JA. Skin lipids: localization of ceramide and fatty acid in the unit cell of the long periodicity phase. *Biophys J*. 2015a;108:2670-2679.
- [44] Mojumdar EH, Gooris GS, Bouwstra J. Phase behavior of skin lipid mixtures: the effect of cholesterol on lipid organization. *Soft matter*. 2015b;11:4326-4336.
- [45] Mojumdar EH, Kariman Z, van Kerckhove L, Gooris GS, Bouwstra JA. The role of ceramide chain length distribution on the barrier properties of the skin lipid membranes. *Biochim Biophys Acta*. 2014;1838:2473-2483.
- [46] Mori K, Matsuda H. Synthesis of sphingosine relatives .10. Synthesis of (2S,3R,4E)-1-O-(beta-D-glucopyranosyl)-N-30'-(linoleoyloxy)triacontanoyl-4-icosasphingenine, a new esterified cerebroside isolated from human and pig epidermis. *Liebigs Ann. Chem*. 1991:529-535.
- [47] Motta S, Monti M, Sesana S, Caputo R, Carelli S, Ghidoni R. Ceramide composition of the psoriatic scale. *Biochim Biophys Acta*. 1993;1182:147-151.
- [48] Muller S, Schmidt RR. Synthesis of two unique compounds, a ceramide and a cerebroside, occurring in human stratum corneum. *J. Prakt. Chem*. 2000;342:779-784.
- [49] Neto DD, Gooris G, Bouwstra J. Effect of the omega-acylceramides on the lipid organization of stratum corneum model membranes evaluated by X-ray diffraction and FTIR studies (Part I). *Chem. Phys. Lipids*. 2011;164:184-195.
- [50] Norlen L. Current understanding of skin barrier morphology. *Skin Pharmacol Physiol*. 2013;26:213-216.
- [51] Norlen L, Nicander I, Lundsjo A, Cronholm T, Forslind B. A new HPLC-based method for the quantitative analysis of inner stratum corneum lipids with special reference to the free fatty acid fraction. *Arch. Dermatol. Res*. 1998;290:508-516.
- [52] Novotný J, Hrabálek A, Vávrová K. Synthesis and structure-activity relationships of skin ceramides. *Curr Med Chem*. 2010;17:2301-2324.
- [53] Novotný J, Janůšová B, Novotný M, Hrabálek A, Vávrová K. Short-chain ceramides decrease skin barrier properties. *Skin Pharmacol. Physiol*. 2009;22:22-30.
- [54] Opálka L, Kováčik A, Maixner J, Vávrová K. Omega-O-Acylceramides in Skin Lipid Membranes: Effects of Concentration, Sphingoid Base, and Model Complexity on Microstructure and Permeability. *Langmuir*. 2016;32:12894-12904.
- [55] Opálka L, Kováčik A, Sochorová M, et al. Scalable Synthesis of Human Ultralong Chain Ceramides. *Org. Lett*. 2015;17:5456-5459.
- [56] Pullmannová P, Staňková K, Pospíšilová M, Školová B, Zbytovská J, Vávrová K. Effects of sphingomyelin/ceramide ratio on the permeability and microstructure of model stratum corneum lipid membranes. *BBA-Biomembranes*. 2014;1838:2115-2126.
- [57] Rabionet M, Bayerle A, Marsching C, et al. 1-O-acylceramides are natural components of human and mouse epidermis. *J. Lipid. Res*. 2013;54:3312-3321.
- [58] Rabionet M, Gorgas K, Sandhoff R. Ceramide synthesis in the epidermis. *Biochim Biophys Acta*. 2014;1841:422-434.
- [59] Rerek ME, Chen H, Markovic B, et al. Phytosphingosine and Sphingosine Ceramide Headgroup Hydrogen Bonding: Structural Insights through Thermotropic Hydrogen/Deuterium Exchange. *J. Phys. Chem. B*. 2001;105:9355 -9362.
- [60] Robson KJ, Stewart ME, Michelsen S, Lazo ND, Downing DT. 6-Hydroxy-4-sphingenine in human epidermal ceramides. *J. Lipid. Res*. 1994;35:2060-2068.
- [61] Shieh H-S, Hoard LG, Nordman CE. The structure of cholesterol. *Acta Crystallogr Sect B*. 1981;37:1538-1543.
- [62] Schreiner V, Pfeiffer S, Lanzendörfer G, et al. Barrier characteristics of different human skin types investigated with X-ray diffraction, lipid analysis, and electron microscopy imaging. *J. Invest. Dermatol*. 2000;114:654-660.
- [63] Skolova B, Hudská K, Pullmannova P, et al. Different phase behavior and packing of ceramides with long (C16) and very long (C24) acyls in model membranes: infrared spectroscopy using deuterated lipids. *J Phys Chem B*. 2014;118:10460-10470.
- [64] Skolova B, Janusova B, Vavrova K. Ceramides with a pentadecaspingosine chain and short acyls have strong permeabilization effects on skin and model lipid membranes. *Biochim Biophys Acta*. 2016;1858:220-232.
- [65] Skolova B, Janusova B, Zbytovska J, et al. Ceramides in the skin lipid membranes: length matters. *Langmuir*. 2013;29:15624-15633.
- [66] Stahlberg S, Lange S, Dobner B, Huster D. Probing the Role of Ceramide Headgroup Polarity in Short-Chain Model Skin Barrier Lipid Mixtures by (2)H Solid-State NMR Spectroscopy. *Langmuir*. 2016;32:2023-2031.
- [67] Stahlberg S, Skolova B, Madhu PK, Vogel A, Vavrova K, Huster D. Probing the role of the ceramide acyl chain length and sphingosine unsaturation in model skin barrier lipid mixtures by (2)H solid-state NMR spectroscopy. *Langmuir*. 2015;31:4906-4915.
- [68] Školová B, Hudská Kr, Pullmannová P, et al. Different phase behavior and packing of ceramides with long (C16) and very long (C24) acyls in model membranes: infrared spectroscopy using deuterated lipids. *J. Phys. Chem. B*. 2014;118:10460-10470.

- [69] Školová B, Janůšová B, Vávrová K. Ceramides with a pentadecaphingosine chain and short acyls have strong permeabilization effects on skin and model lipid membranes. *BBA-Biomembranes*. 2016;1858:220-232.
- [70] Školová B, Janůšová B, Zbytovská J, et al. [Ceramide in the skin lipid membranes: length matters](#). *Langmuir*. 2013;29:15624-15633.
- [71] t'Kindt R, Jorge L, Dumont E, et al. Profiling and characterizing skin ceramides using reversed-phase liquid chromatography-quadrupole time-of-flight mass spectrometry. *Anal Chem*. 2012;84:403-411.
- [72] Uchida Y, Holleran WM. Omega-O-acylceramide, a lipid essential for mammalian survival. *Journal of Dermatological Science*. 2008;51:77-87.
- [73] van Smeden J, Boiten WA, Hankemeier T, Rissmann R, Bouwstra JA, Vreeken RJ. Combined LC/MS-platform for analysis of all major stratum corneum lipids, and the profiling of skin substitutes. *Biochim Biophys Acta*. 2014a;1841:70-79.
- [74] van Smeden J, Hoppel L, van der Heijden R, Hankemeier T, Vreeken RJ, Bouwstra JA. LC/MS analysis of stratum corneum lipids: ceramide profiling and discovery. *J. Lipid Res*. 2011;52:1211-1221.
- [75] van Smeden J, Janssens M, Gooris GS, Bouwstra JA. The important role of stratum corneum lipids for the cutaneous barrier function. *Biochim Biophys Acta*. 2014b;1841:295-313.
- [76] Vasireddy V, Uchida Y, Salem N, et al. *Hum. Mol. Genet*. 2007;16:471.
- [77] Vavrova K, Henkes D, Struver K, et al. Filaggrin Deficiency Leads to Impaired Lipid Profile and Altered Acidification Pathways in a 3D Skin Construct. *J Invest Dermatol*. 2014;134:746-753.
- [78] Wertz PW, Madison KC, Downing DT. [Covalently bound lipids of human stratum corneum](#). *J. Invest. Dermatol*. 1989;92:109-111.
- [79] White SH, Mirejovsky D, King GI. Structure of lamellar lipid domains and corneocyte envelopes of murine stratum corneum. An X-ray diffraction study. *Biochemistry*. 1988;27:3725-3732.